

# Dual Roles of Autophagy and Their Potential Drugs for Improving Cancer Therapeutics

Dong Wook Shin\*

College of Biomedical and Health Science, Konkuk University, Chungju 27478, Republic of Korea

## Abstract

Autophagy is a major catabolic process that maintains cell metabolism by degrading damaged organelles and other dysfunctional proteins via the lysosome. Abnormal regulation of this process has been known to be involved in the progression of pathophysiological diseases, such as cancer and neurodegenerative disorders. Although the mechanisms for the regulation of autophagic pathways are relatively well known, the precise regulation of this pathway in the treatment of cancer remains largely unknown. It is still complicated whether the regulation of autophagy is beneficial in improving cancer. Many studies have demonstrated that autophagy plays a dual role in cancer by suppressing the growth of tumors or the progression of cancer development, which seems to be dependent on unknown characteristics of various cancer types. This review summarizes the key targets involved in autophagy and malignant transformation. In addition, the opposing tumor-suppressive and oncogenic roles of autophagy in cancer, as well as potential clinical therapeutics utilizing either regulators of autophagy or combinatorial therapeutics with anti-cancer drugs have been discussed.

**Key Words:** Autophagy, Cancer, Target, Anti-tumor drug, Combinational therapy

## INTRODUCTION

Cancer is characterized by excessive cell growth and malignancy due to the accumulation of genetic defects or abnormal metabolic processes (Sell *et al.*, 2016; Dang and Kim, 2018). Upregulated catabolism in cancer cells promotes tumor growth and metastasis (Danhier *et al.*, 2017). Moreover, the rapid proliferation of cancer cells is dependent on the recycling of cellular components (Danhier *et al.*, 2017; Momcilovic and Shackelford, 2018). Recent studies have demonstrated that the metabolic processes of tumor cells are correlated with autophagy (Su *et al.*, 2015; Mowers *et al.*, 2017). Autophagy is a mechanism by which abnormal proteins or damaged cellular organelles undergo lysosomal degradation, which provides energy and macromolecular precursors (Yorimitsu and Klionsky, 2005; Mizushima, 2007). Autophagy is involved in the initiation and development of cancer by preventing toxic accumulation of carcinogenic factors (Guo and White, 2016).

Cancer cells meet their high metabolic energy demand by utilizing degraded biomolecules derived from autophagy (Galati *et al.*, 2019). Compared to normal tissues, cancerous tissues are highly dependent on autophagy. Thus, targeting

autophagy may be a potential therapeutic strategy against cancer (White *et al.*, 2015). The role of autophagy in malignant transformation is complex and varies depending on the cancer cell type (Galluzzi *et al.*, 2015). Further elucidation of the interaction between autophagy regulation and the key molecules involved in each cancer type is crucial for the identification of an effective therapeutic.

Autophagy-related genes function as either tumor suppressors or oncogenes. Whereas suppression of autophagy inhibits the growth of cancer cells, the induction of autophagy has also been reported to decrease tumor growth (Kimmelman and White, 2017). Many studies suggest that autophagy may function as a tumor suppressor in the early stages of cancer progression, but exert pro-tumor effects in later stages of cancer (Kimmelman and White, 2017; Kocaturk *et al.*, 2019).

This review summarizes recent findings on the molecular regulation of autophagy in cancer. In addition, the therapeutic potential of various autophagy inducers, and inhibitors alone and in combination with known anti-cancer drugs is discussed.

**Open Access** <https://doi.org/10.4062/biomolther.2020.155>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received Sep 7, 2020 Revised Sep 18, 2020 Accepted Sep 18, 2020  
Published Online Oct 20, 2020

**\*Corresponding Author**

E-mail: biocosmed@kku.ac.kr

Tel: +82-43-840-3693, Fax: +82-43-840-3929

## MOLECULAR MECHANISM OF AUTOPHAGY

Autophagy is classified into the following three types: macroautophagy, microautophagy, and chaperone-mediated autophagy (CMA). Macroautophagy (called autophagy) is a highly evolutionarily conserved pathway for the degradation of dysfunctional proteins (Yorimitsu and Klionsky, 2005; Kuma *et al.*, 2017; Kaushik and Cuervo, 2018). This process is accompanied by evolutionarily well-conserved autophagy-related (Atg) proteins and the formation of a vesicle structure known as an autophagosome. The contents of autophagosomes are degraded by lysosomal hydrolase upon fusion with the lysosome (Yorimitsu and Klionsky, 2005; Kuma *et al.*, 2017; Kaushik and Cuervo, 2018). Microautophagy involves direct uptake of cytosolic components by lysosomes (Li and Hochstrasser, 2020). CMA is a complex proteolytic system involving lysosomal membrane-associated protein 2 A and heat-shock protein 70 (Kaushik and Cuervo, 2018).


Nutrient signaling modulates the formation of autophagosomes via the mammalian target of rapamycin (mTOR) and adenosine monophosphate-activated protein kinase (AMPK) (Tamargo-Gomez and Marino, 2018). Under physiological conditions, mTORC1 inhibits autophagy by phosphorylating unco-51-like kinase (ULK1), which initiates autophagosome biogenesis. In contrast, mTORC1 is inhibited during starvation (Yorimitsu and Klionsky, 2005; Russell *et al.*, 2014). The ULK1 complex, which is activated by autophosphorylation, phosphorylates FIP200, ATG13, ATG101, and other ATG proteins (Park *et al.*, 2016). Nutrient deprivation or starvation causes the depletion of ATP and leads to increased ADP and AMP

levels, which activates AMPK (Tamargo-Gomez and Marino, 2018; Li and Chen, 2019). Further, AMPK contributes to the restoration of cellular energy levels through autophagy by inhibiting mTORC1 and phosphorylating ULK1 (Li and Chen, 2019).

The ULK1 complex triggers autophagosome nucleation by activating a multi-domain complex comprising Beclin 1, ATG14, and VPS34 (Russell *et al.*, 2013). Autophagosomes are modulated by the ATG5-ATG12-ATG16L1 complex, which facilitates the conjugation of LC3-I to phosphatidylethanolamine to form LC3-II (Russell *et al.*, 2014). Meanwhile, misfolded proteins or damaged organelles are marked for degradation through ubiquitination. Polyubiquitinated target proteins destined for degradation are recognized by sequestosome-1 (SQSTM1/p62) (Moscat and Diaz-Meco, 2009; Kuma *et al.*, 2017; Zaffagnini *et al.*, 2018). LC3-II recruits the ubiquitinated p62-cargo complex, which is engulfed by autophagosomes (Zaffagnini *et al.*, 2018). ATG4B mediates the recycling of the LC3-II protein to LC3-I (Zaffagnini *et al.*, 2018). The fully mature autophagosomes fuse with acidic lysosomes, which results in autolysosome formation. Lysosomal hydrolases degrade the target molecules or organelles, which are then used as metabolite substrates or precursors for macromolecule biosynthesis (Fig. 1) (Mizushima, 2007; Russell *et al.*, 2014).

## CORRELATION BETWEEN AUTOPHAGY AND TUMOR FORMATION

Autophagy contributes to the inhibition of tumorigenesis at



**Fig. 1.** Overview of Autophagy Process. The depletion of ATP activates AMPK under fasting conditions. The AMPK activates autophagy by suppressing the activity of mTORC1 and by phosphorylating the ULK1. The ULK1 complex initiates the formation of autophagosome by stimulating Beclin 1/ATG14/VPS34 complex. The Atg5-Atg12-Atg16L1 complex, which regulates the autophagosomes, converts LC3-I into LC3-II. Abnormal organelles or proteins after their ubiquitination (Ub) is recognized by p62. LC3-II interacts with the ubiquitinated p62-cargo complex and is engulfed into the autophagosomes. The autophagosomes fuse with lysosome and leads to the formation of autolysosome. The target molecules or organelles are finally degraded by lysosomal proteases and are utilized as metabolite substrates.

multiple stages (Galluzzi *et al.*, 2015; Amaravadi *et al.*, 2016). ATGs in the autophagy pathway are reported to suppress tumor formation. Deletion of ATG4C has been shown to promote tumorigenesis in mice (Marino *et al.*, 2007). Similarly, ATG5 deficiency induces benign tumor formation in mouse livers (Takamura *et al.*, 2011).

ULK1 (human ATG1) stimulates autophagic flux, which results in the inhibition of cell proliferation during starvation (Jung *et al.*, 2011). In addition, ULK2 is markedly downregulated in glioma. These findings indicate that autophagy inhibition through ULK1/2 downregulation promotes tumor progression (Shukla *et al.*, 2014).

Beclin-1, which is involved in autophagy, functions as a tumor suppressor, and its expression is inhibited in various human cancers, such as prostate and colon cancers (Chen and Karantza-Wadsworth, 2009). Monoallelic Beclin-1 gene deletions result in malignancies in humans (Qu *et al.*, 2003). Further, Beclin-1 overexpression markedly suppresses proliferation and induces apoptosis in human laryngeal squamous carcinoma cells (Wan *et al.*, 2018). Beclin-1-mediated autophagy suppresses cell death and promotes survival under stress. The activity of Beclin-1 is repressed in the presence of the anti-apoptotic proteins BCL-2 and BCL-XL. Moreover, BCL-2, which is upregulated in several cancers (Huang, 2000), negatively regulates Beclin-1-mediated autophagy (Ramakrishnan *et al.*, 2007). The interaction between BCL-2 and Beclin-1 promotes autophagy or apoptosis in various cancer cells (Marquez and Xu, 2012). Caspase or calpain, which plays an important role in apoptosis, inhibits Beclin-1-mediated autophagy (Kang *et al.*, 2011). p53 levels also regulate Beclin-1 in the autophagic flux. Specifically, overexpression of p53 results in the degradation of Beclin-1 and leads to the suppression of autophagy in embryonal carcinoma (Tripathi *et al.*, 2014). Therefore, proteins interacting with Beclin-1 are essential for determining whether cells undergo autophagy or apoptosis.

p62 (or sequestosome-1, SQSTM1) is a signaling scaffold that is upregulated in human cancers (Liu *et al.*, 2012;

Umemura *et al.*, 2016). Autophagy suppression results in p62 accumulation, which promotes oncogenesis by increasing DNA damage and endoplasmic reticulum stress (Moscat and Diaz-Meco, 2009). Benign tumors in ATG5 or ATG7 knockout mice have been shown to exhibit high levels of p62. Loss of p62 in these mice inhibits the growth of tumors (Takamura *et al.*, 2011; Liu *et al.*, 2012). Diethylnitrosamine, a potent liver carcinogen, stimulates the tumorigenic activity of p62 and aggravates hepatocellular carcinoma (Umemura *et al.*, 2016). Therefore, p62 inhibition during autophagy is a potential therapeutic strategy against cancer. These data suggest that autophagy-related gene regulation plays a key role in determining cancer progression.

## AUTOPHAGY INDUCERS IN CANCER THERAPY

Activation of autophagy may be a direct strategy to promote tumor cell death (Table 1). Some tumor cells exhibit resistance to apoptosis. Thus, autophagy may provide an alternative cell death mechanism for cancer cells with defective apoptosis (Tsujiimoto and Shimizu, 2005).

As autophagy is negatively regulated by mTOR, some autophagy-inducing drugs target the mTOR pathway. Rapamycin and its derivative, everolimus, inhibit mTORC1, which promotes the dissociation of the ULK1 complex (Vignot *et al.*, 2005; Cicchini *et al.*, 2015). Rapamycin has been reported to suppress the proliferation of MCF-7 breast cancer (Chang *et al.*, 2007), B16 melanoma (Busca *et al.*, 1996), and PANC-1 pancreatic carcinoma cell lines (Grewe *et al.*, 1999). Similarly, everolimus inhibits the growth of various malignancies (Hare and Harvey, 2017; Bhat *et al.*, 2018; Kocaturk *et al.*, 2019). For example, it suppresses human vascular endothelial cell growth by delaying the cell cycle in lymphoblastic B cells (Neri *et al.*, 2014). Everolimus, although not as effective as rapamycin, exerts similar anti-tumor effects *in vivo*. Further, temsirolimus (CCI-779), a rapamycin analog, activates autophagy by inhibiting mTOR in adenoid cystic carcinoma (Liu *et al.*, 2014).

**Table 1.** Autophagy inducers for improving cancer

| Compound                              | Target or mode of action         | Cancer types  | References   |
|---------------------------------------|----------------------------------|---|--|
| Rapamycin                             | mTOR                             | MCF-7 Breast Cancer, B16 melanoma, Pan-1 pancreatic carcinoma           | Busca <i>et al.</i> , 1996; Grewe <i>et al.</i> , 1999; Chang <i>et al.</i> , 2007 |
| Everolimus                            | mTOR                             | Lymphoblastic B cells   | Neri <i>et al.</i> , 2014  |
| Temsirolimus (CCI-779)                | mTOR                             | Adenoid cystic carcinoma  | Liu <i>et al.</i> , 2014   |
| Metformin                             | AMPK                             | Prostate cancer cells, myeloma  | Sesen <i>et al.</i> , 2015; Wang <i>et al.</i> , 2018; Mishra and Dingli, 2019     |
| Perifosine                            | Akt                              | Multiple myeloma, neuroblastoma, colorectal cancer, neuroblastoma cells | Li <i>et al.</i> , 2010, 2011; Richardson <i>et al.</i> , 2012                     |
| Ibrutinib                             | Bruton's Tyrosine Kinase (BTK)   | Skin cancer cell lines, HS-4 and A431                                   | Sun <i>et al.</i> , 2018   |
| Suberoylanilidehydroxamic acid (SAHA) | Histone deacetylase (HDAC), mTOR | Cutaneous T cell lymphoma, glioblastoma stem cells                      | Gammoh <i>et al.</i> , 2012; Chiao <i>et al.</i> , 2013                            |
| Magnolol                              | LIF/Stat-3/Mcl-1                 | Colorectal cancer cells   | Yu <i>et al.</i> , 2018  |
| Resveratrol                           | SIRT1                            | Prostate cancer cells, MCF-7 cells                                      | El-Mowafy and Alkhalaf, 2003; Li <i>et al.</i> , 2014                              |
| Spermidine                            | AMPK                             | Colon cancer cells  | Morselli <i>et al.</i> , 2011  |

Metformin (250 mg/kg body weight), an anti-diabetic drug, stimulates autophagy through AMPK activation and markedly inhibits prostate cancer cell proliferation in HimYC mice (Ben Saha *et al.*, 2010). Additionally, metformin is reported to inhibit the growth of glioblastoma cells (Sesen *et al.*, 2015). Metformin arrests the cell cycle at the G0/G1 phase and suppresses myeloma cell proliferation by regulating AMPK and mTORC (Wang *et al.*, 2018; Mishra and Dingli, 2019). Furthermore, metformin has been shown to enhance the dual effects of chemotherapy and radiation therapy (Saha *et al.*, 2015; Sesen *et al.*, 2015).

Perifosine induces autophagy through the suppression of Akt, which interferes with the phosphatidylinositol-3-kinase (PI3K) signaling pathway. The therapeutic effect of perifosine on multiple myeloma, neuroblastoma, and colorectal cancer has been evaluated in phase III clinical trials (Li *et al.*, 2010; Richardson *et al.*, 2012). Akt inhibition suppresses TrkB/neurotrophic factor-induced resistance to chemotherapy, which increases the neuroblastoma cell sensitivity to chemotherapy and radiation therapy. Furthermore, perifosine promotes apoptosis in neuroblastoma cells (Li *et al.*, 2010) and suppresses the proliferation of colorectal cancer cells (Li *et al.*, 2011). Ibrutinib, a Bruton's tyrosine kinase inhibitor, stimulates autophagy by upregulating ATG7 and promoting LC3-II formation. Additionally, ibrutinib has been shown to exhibit growth-inhibiting activity against the skin cancer cell lines HS-4 and A431 (Sun *et al.*, 2018). Histone deacetylase (HDAC) inhibitors are also reported to affect the autophagic flux (Bhat *et al.*, 2018). Suberoylanilide hydroxamic acid (SAHA), an HDAC inhibitor, promotes autophagy through mTOR inhibition and enhances apoptosis in cutaneous T cell lymphoma (Gammoh *et al.*, 2012). Another study reported that SAHA induces autophagy, which delays tumor growth in glioblastoma stem cells (Chiao *et al.*, 2013).

In addition, some natural compounds treat cancer via activating the autophagic flux. Magnolin, a bioactive natural compound, activates autophagy and arrests the cell cycle in human colorectal cancer cells by inhibiting the LIF/Stat-3/Mcl-1 pathway (Yu *et al.*, 2018). Resveratrol stimulates autophagy by activating sirtuin 1, which promotes autophagy through ATG5 and ATG7 deacetylation (Lee *et al.*, 2008; Morselli *et al.*, 2011). Further, resveratrol suppresses prostate cancer cell proliferation (Li *et al.*, 2014) and promotes apoptosis in MCF-7 cells (El-Mowafy and Alkhalaf, 2003). Spermidine, a natural polyamine, promotes autophagy through an mTOR-independent pathway. Additionally, spermidine phosphorylates protein tyrosine kinase 2 $\beta$  and cyclin-dependent kinase inhibitor 1B via AMPK (Morselli *et al.*, 2011) and regulates the acetylation status of ATG5 and LC3 in human colon cancer cells (Morselli *et al.*, 2011). Therefore, elucidating the regulatory mechanisms underlying autophagy and cell death in cancer cells will contribute to the development of novel therapeutic strategies.

## AUTOPHAGY INHIBITORS IN CANCER THERAPY

Several autophagy inhibitors effectively inhibit cancer cell proliferation (Table 2). Some inhibitors target kinases in the phagophore or inhibit the fusion of autophagosomes and lysosomes during the autophagic flux (Yang *et al.*, 2013; Yuan *et al.*, 2015; Mauthe *et al.*, 2018).

3-Methyladenine (3-MA) and 2-(4-morpholinyl)-8-phenyl-

chromone (LY294002) inhibit autophagy by suppressing the activity of PI3K, which is involved in the production of phosphatidylinositol (3,4,5)-triphosphate and the nucleation and extension of the phagophore (Yang *et al.*, 2013). Class I PI3Ks negatively regulate autophagy, whereas class III PI3Ks directly promote autophagy (Bilanges *et al.*, 2019). LY294002 and 3-MA inactivate class III PI3Ks, which has been shown to induce caspase-induced death in HeLa cells (Hou *et al.*, 2012).

Bafilomycin A1 promotes the binding of Beclin-1 to BCL-2, which inhibits autophagy (Yuan *et al.*, 2015) and suppresses the invasion and migration of gastric cancer cells, as well as promoting the apoptosis of these cells (Li *et al.*, 2016). Bafilomycin A1 is also a lysosomal H<sup>+</sup>-ATPase inhibitor and impairs autophagy by inhibiting the fusion of autophagosomes with lysosomes (Bhat *et al.*, 2018).

Chloroquine (CQ) and hydroxychloroquine (HCQ) are clinically approved by the U.S. Food and Drug Administration as anti-malarial agents. CQ acts as an autophagy inhibitor by suppressing the fusion process of autophagosomes and lysosomes (Mauthe *et al.*, 2018). CQ also promotes the production of cellular reactive oxygen species and enhances the cytotoxicity of temozolomide by suppressing mitophagy in glioma cells (Hori *et al.*, 2015). In addition, CQ, in combination with different anti-cancer drugs, exhibits an anti-cancer effect in breast (Cave *et al.*, 2018) and colon cancer (Liu *et al.*, 2019).

HCQ enhances anti-tumor immunity and suppresses autophagy by promoting p62 accumulation and upregulating the lysosomal protease cathepsin D in metastatic colorectal cancer (Patel *et al.*, 2016). Additionally, HCQ suppresses autophagy to potentiate the anti-estrogen responsiveness of breast cancer (Cook *et al.*, 2014). Quinacrine, a synthetic anti-malarial drug, has been reported to suppress autophagic flux (Golden *et al.*, 2015). Quinacrine also increases the expression of p21/p27 independent of p53 owing to the downregulation of the p62-Skp2 axis in ovarian cancer (Jung *et al.*, 2018).

BCL-2 proteins, which are representative regulators of apoptosis signaling pathways, regulate autophagy (Xu and Qin, 2019). ABT-737, a BCL-2 inhibitor, has been reported to be an autophagy inhibitor. Moreover, colorectal cancer cell sensitivity to ixazomib, used to treat multiple myeloma, is enhanced upon ABT-737 treatment through MCL-1 downregulation and autophagy inhibition (Yang *et al.*, 2016). Obatoclax, which inhibits Pan-BCL-2, suppresses autophagy in the bladder (Jimenez-Guerrero *et al.*, 2018) and colorectal cancer at a late stage (Koehler *et al.*, 2015).

Autophagy is associated with proteasome pathways. Bortezomib, a proteasome inhibitor, is an approved anti-cancer drug for multiple myeloma. Bortezomib treatment initiates autophagosome formation, induces autophagic flux, and upregulates ATG5 in human prostate cancer cells (Zhu *et al.*, 2010). Bortezomib also regulates both apoptosis and autophagic pathways via mitogen-activated protein kinase activation in osteocarcinoma (Lou *et al.*, 2013). In breast cancer cell lines, bortezomib suppresses autophagy by inhibiting the cathepsin activity and promoting caspase-dependent apoptosis (Periyasamy-Thandavan *et al.*, 2010). Bortezomib also promotes apoptosis in ovarian cancer cells by suppressing autophagic flux (Kao *et al.*, 2014). The treatment combination of bortezomib and 3-MA has been shown to promote apoptosis in the human glioblastoma cell lines, U87 and U251 (Zhang *et al.*, 2014).

**Table 2.** Autophagy inhibitors for treating cancer

| Compound                         | Target or mode of action                             | Cancer types   | References  |
|----------------------------------|--|--|---|
| 3-Methyladenine (3-MA), LY294002 | Class III PI3K                                       | HeLa cells   | Hou <i>et al.</i> , 2012  |
| Bafilomycin A1                   | Beclin-1, lysosomal H <sup>+</sup> -ATPase inhibitor | Gastric cancer cells   | Li <i>et al.</i> , 2016   |
| Chloroquine (CQ)                 | Fusion process of autophagosome and lysosome         | Glioma cells, breast cancer, colon cancer  | Hori <i>et al.</i> , 2015; Cave <i>et al.</i> , 2018; Liu <i>et al.</i> , 2019  |
| Hydroxychloroquine (HCQ)         | Lysosomal cathepsin D                                | Colorectal cancer, breast cancer   | Cook <i>et al.</i> , 2014; Patel <i>et al.</i> , 2016   |
| Quinacrine                       | p21/p27  | Ovarian cancer   | Jung <i>et al.</i> , 2018   |
| ABT-737                          | Bcl-2  | Multiple myeloma, colorectal cancer cells  | Sun <i>et al.</i> , 2018  |
| Obatoclax                        | Bcl-2  | Bladder cancer, colorectal cancer  | Koehler <i>et al.</i> , 2015; Jimenez-Guerrero <i>et al.</i> , 2018   |
| Bortezomib                       | Proteasome   | Multiple myeloma, prostate cancer cells, osteocarcinoma, breast cancer cells, ovarian cancer cells, glioblastoma | Periyasamy-Thandavan <i>et al.</i> , 2010; Zhu <i>et al.</i> , 2010; Lou <i>et al.</i> , 2013; Kao <i>et al.</i> , 2014; Zhang <i>et al.</i> , 2014 |
| Elaiophyllin                     | Inhibition of autophagy flux                         | Ovarian cancer cells, multiple myeloma   | Zhao <i>et al.</i> , 2015; Wang <i>et al.</i> , 2017  |
| 4-Acetylanthroquinonol B         | Inhibition of autophagy flux                         | Epithelial cancer cells  | Liu <i>et al.</i> , 2017  |
| Thymoquinone                     | Permeabilization of the lysosome membrane            | Glioblastoma cells   | Racoma <i>et al.</i> , 2013   |

**Table 3.** Combinatorial treatment by targeting autophagy and anti-tumor drug in clinical trials

| Compounds                  | Cancer types   | References  |
|----------------------------|--|---|
| CQ+temozolomide            | Glioblastoma multiforme, glioblastomas, glioma cells | Hori <i>et al.</i> , 2015; Zanotto-Filho <i>et al.</i> , 2015 |
| HCQ+temozolomide           | Solid tumors and melanoma                            | Rangwala <i>et al.</i> , 2014a                                |
| HCQ+temsirolimus (CCI-779) | Melanoma   | Rangwala <i>et al.</i> , 2014b                                |
| HCQ+bortezomib             | Myeloma  | Vogl <i>et al.</i> , 2014                                     |
| HCQ and vorinostat         | Solid tumors, metastatic colorectal cancer           | Mahalingam <i>et al.</i> , 2014; Patel <i>et al.</i> , 2016   |
| Pantoprazole+doxorubicin   | Solid tumors   | Brana <i>et al.</i> , 2014                                    |

Clarithromycin, a macrolide antibiotic, is used to treat upper and lower respiratory tract and *Helicobacter pylori* infections (Crowe, 2019). Clarithromycin suppresses autophagy by inhibiting the interaction between the hERG1 potassium channel and PI3K in colorectal cancer (Altman and Plataniotis, 2012; Petroni *et al.*, 2020). Elaiophyllin, a macrolide antibiotic with immunosuppressive properties, inhibits the autophagic flux and has anti-tumor activity in ovarian cancer (Zhao *et al.*, 2015) and multiple myeloma with mutant TP53 (Wang *et al.*, 2017).

4-Acetylanthroquinonol B, a novel compound derived from anthroquinonol, enhances the sensitivity of cisplatin in epithelial cancer cells by inhibiting the autophagic flux (Liu *et al.*, 2017). Thymoquinone stimulates the permeabilization of the lysosome membrane and inhibits autophagy, which induces cathepsin-mediated death in glioblastoma cells (Racoma *et al.*, 2013). Further studies are required to elucidate the effect

of autophagy inhibitors on cancer cells, to develop potential therapeutic agents against cancer (Table 2).

## COMBINATION TREATMENT OF AN AUTOPHAGY REGULATOR AND AN ANTI-CANCER DRUG

Studies have suggested that the efficacy of anti-cancer drugs is enhanced upon co-administration of autophagic flux regulators (Table 3).

CQ enhances the efficacy of temozolomide, which is an alkylating agent used to treat brain cancers, such as glioblastoma multiforme and glioblastomas (Zanotto-Filho *et al.*, 2015) by suppressing mitophagy in glioma cells (Hori *et al.*, 2015). Additionally, CQ promotes apoptotic cell death induced by vorinostat, an HDAC inhibitor, in colon cancer (Carew *et al.*, 2010).



HCQ, in combination with temozolomide, is effective in patients suffering from solid tumors and melanoma, in a phase 1 study (Rangwala *et al.*, 2014a). Another phase 1 dose-escalation study has demonstrated that the anti-tumor effect of HCQ and temsirolimus (CCI-779; an mTOR inhibitor) treatment combination is stronger than that of HCQ alone in patients with melanoma (Rangwala *et al.*, 2014b). These studies suggest that autophagy inhibitors markedly enhance the therapeutic effects of anti-cancer drugs. The suppression of autophagy may promote apoptosis, which can enhance the sensitivity of cancer cells to radiation therapy. Vogl *et al.* (2014) demonstrated that the treatment combination of HCQ and bortezomib increases the efficacy of proteasome inhibition in patients with myeloma. Additionally, the efficacy of proteasome inhibitors is strongly dependent on the ability of malignant cells to degrade misfolded proteins (Vogl *et al.*, 2014). Furthermore, Mahalingam *et al.* (2014) demonstrated the pharmacokinetic and pharmacodynamic properties, as well as the safety of the HCQ and vorinostat treatment combination in patients with advanced solid tumors. This combination treatment suppresses autophagy and markedly enhances anti-tumor immunity in metastatic colorectal cancer (Patel *et al.*, 2016).

Similarly, a phase I clinical study revealed that pantoprazole, a proton pump inhibitor, enhances the anti-cancer efficacy of doxorubicin by inhibiting autophagy in patients with solid tumors (Brana *et al.*, 2014). These results suggest that the anti-cancer efficacy of an autophagy inhibitor and anti-cancer drug treatment combination is higher than that of anti-tumor drugs alone (Table 3).

## CONCLUSION

Malignant transformation involves dramatic changes in cellular metabolism owing to the high energy demand. During carcinogenesis, a continuous supply of macromolecule precursors is required to support abnormal cell proliferation. Thus, autophagy regulation has recently emerged as a potential therapeutic strategy against cancer.

Autophagy exhibits either cytoprotective or cytotoxic activity as dual roles during carcinogenesis, depending on the heterogeneity of each tumor. Studies have suggested that autophagy suppresses tumor growth in the early stages, but accelerates tumor progression in the later stages (Kimmelman and White, 2017; Kocaturk *et al.*, 2019).

Autophagy and cancer pathways have been annotated. However, the interactions between these pathways have not been elucidated. The physiological role of autophagy must be analyzed in each tumor type. Various drugs, either alone or in combination, inhibit cancer progression by regulating autophagy (Table 1-3). However, further studies are required to determine the timing, intervals, and dosage of potent autophagy regulator administration in each cancer type. Additionally, clinical studies are required to further examine the anti-cancer efficacy of treatment combinations of autophagy inhibitors and anti-cancer drugs.

## CONFLICT OF INTEREST

The author declares that there are no conflicts of interest.

## ACKNOWLEDGMENTS

This work was supported by the research fund from Konkuk University (2019-A019-0401).

## REFERENCES

- Altman, J. K. and Platanias, L. C. (2012) A new purpose for an old drug: inhibiting autophagy with clarithromycin. *Leuk. Lymphoma* **53**, 1255-1256.
- Amaravadi, R., Kimmelman, A. C. and White, E. (2016) Recent insights into the function of autophagy in cancer. *Genes Dev.* **30**, 1913-1930.
- Ben Sahra, I., Tanti, J. F. and Bost, F. (2010) The combination of metformin and 2-deoxyglucose inhibits autophagy and induces AMPK-dependent apoptosis in prostate cancer cells. *Autophagy* **6**, 670-671.
- Bhat, P., Kriel, J., Shubha Priya, B., Basappa, Shivananju, N. S. and Loos, B. (2018) Modulating autophagy in cancer therapy: advancements and challenges for cancer cell death sensitization. *Biochem. Pharmacol.* **147**, 170-182.
- Bilanges, B., Posor, Y. and Vanhaesebroeck, B. (2019) PI3K isoforms in cell signalling and vesicle trafficking. *Nat. Rev. Mol. Cell Biol.* **20**, 515-534.
- Brana, I., Ocana, A., Chen, E. X., Razak, A. R., Haines, C., Lee, C., Douglas, S., Wang, L., Siu, L. L., Tannock, I. F. and Bedard, P. L. (2014) A phase I trial of pantoprazole in combination with doxorubicin in patients with advanced solid tumors: evaluation of pharmacokinetics of both drugs and tissue penetration of doxorubicin. *Invest. New Drugs* **32**, 1269-1277.
- Busca, R., Bertolotto, C., Ortonne, J. P. and Ballotti, R. (1996) Inhibition of the phosphatidylinositol 3-kinase/p70(S6)-kinase pathway induces B16 melanoma cell differentiation. *J. Biol. Chem.* **271**, 31824-31830.
- Carew, J. S., Medina, E. C., Esquivel, J. A., 2nd, Mahalingam, D., Swords, R., Kelly, K., Zhang, H., Huang, P., Mita, A. C., Mita, M. M., Giles, F. J. and Nawrocki, S. T. (2010) Autophagy inhibition enhances vorinostat-induced apoptosis via ubiquitinated protein accumulation. *J. Cell. Mol. Med.* **14**, 2448-2459.
- Cave, D. D., Desiderio, V., Mosca, L., Ilisso, C. P., Mele, L., Caraglia, M., Cacciapuoti, G. and Porcelli, M. (2018) S-Adenosylmethionine-mediated apoptosis is potentiated by autophagy inhibition induced by chloroquine in human breast cancer cells. *J. Cell. Physiol.* **233**, 1370-1383.
- Chang, S. B., Miron, P., Miron, A. and Iglehart, J. D. (2007) Rapamycin inhibits proliferation of estrogen-receptor-positive breast cancer cells. *J. Surg. Res.* **138**, 37-44.
- Chen, N. and Karantza-Wadsworth, V. (2009) Role and regulation of autophagy in cancer. *Biochim. Biophys. Acta* **1793**, 1516-1523.
- Chiao, M. T., Cheng, W. Y., Yang, Y. C., Shen, C. C. and Ko, J. L. (2013) Suberoylanilide hydroxamic acid (SAHA) causes tumor growth slowdown and triggers autophagy in glioblastoma stem cells. *Autophagy* **9**, 1509-1526.
- Cicchini, M., Karantza, V. and Xia, B. (2015) Molecular pathways: autophagy in cancer--a matter of timing and context. *Clin. Cancer Res.* **21**, 498-504.
- Cook, K. L., Warri, A., Soto-Pantoja, D. R., Clarke, P. A., Cruz, M. I., Zwart, A. and Clarke, R. (2014) Hydroxychloroquine inhibits autophagy to potentiate antiestrogen responsiveness in ER+ breast cancer. *Clin. Cancer Res.* **20**, 3222-3232.
- Crowe, S. E. (2019) Helicobacter pylori Infection. *N. Engl. J. Med.* **380**, 1158-1165.
- Dang, C. V. and Kim, J. W. (2018) Convergence of cancer metabolism and immunity: an overview. *Biomol. Ther. (Seoul)* **26**, 4-9.
- Danhier, P., Banski, P., Payen, V. L., Grasso, D., Ippolito, L., Sonveaux, P. and Porporato, P. E. (2017) Cancer metabolism in space and time: beyond the Warburg effect. *Biochim. Biophys. Acta* **1858**, 556-572.
- El-Mowafy, A. M. and Alkhalaf, M. (2003) Resveratrol activates adenylyl-cyclase in human breast cancer cells: a novel, estrogen

- receptor-independent cytostatic mechanism. *Carcinogenesis* **24**, 869-873.
- Galati, S., Boni, C., Gerra, M. C., Lazzaretti, M. and Buschini, A. (2019) Autophagy: a player in response to oxidative stress and DNA damage. *Oxid. Med. Cell. Longev.* **2019**, 5692958.
- Galluzzi, L., Pietrocola, F., Bravo-San Pedro, J. M., Amaravadi, R. K., Baehrecke, E. H., Cecconi, F., Codogno, P., Debnath, J., Gewirtz, D. A., Karantza, V., Kimmelman, A., Kumar, S., Levine, B., Maiuri, M. C., Martin, S. J., Penninger, J., Piacentini, M., Rubinsztein, D. C., Simon, H. U., Simonsen, A., Thorburn, A. M., Velasco, G., Ryan, K. M. and Kroemer, G. (2015) Autophagy in malignant transformation and cancer progression. *EMBO J.* **34**, 856-880.
- Gammoh, N., Lam, D., Puente, C., Ganley, I., Marks, P. A. and Jiang, X. (2012) Role of autophagy in histone deacetylase inhibitor-induced apoptotic and nonapoptotic cell death. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 6561-6565.
- Golden, E. B., Cho, H. Y., Hofman, F. M., Louie, S. G., Schonthal, A. H. and Chen, T. C. (2015) Quinoline-based antimalarial drugs: a novel class of autophagy inhibitors. *Neurosurg. Focus* **38**, E12.
- Grewé, M., Gansauge, F., Schmid, R. M., Adler, G. and Seufferlein, T. (1999) Regulation of cell growth and cyclin D1 expression by the constitutively active FRAP-p70s6K pathway in human pancreatic cancer cells. *Cancer Res.* **59**, 3581-3587.
- Guo, J. Y. and White, E. (2016) Autophagy, metabolism, and cancer. *Cold Spring Harb. Symp. Quant. Biol.* **81**, 73-78.
- Hare, S. H. and Harvey, A. J. (2017) mTOR function and therapeutic targeting in breast cancer. *Am. J. Cancer Res.* **7**, 383-404.
- Hori, Y. S., Hosoda, R., Akiyama, Y., Sebori, R., Wanibuchi, M., Mikami, T., Sugino, T., Suzuki, K., Maruyama, M., Tsukamoto, M., Mikuni, N., Horio, Y. and Kuno, A. (2015) Chloroquine potentiates temozolomide cytotoxicity by inhibiting mitochondrial autophagy in glioma cells. *J. Neurooncol.* **122**, 11-20.
- Hou, H., Zhang, Y., Huang, Y., Yi, Q., Lv, L., Zhang, T., Chen, D., Hao, Q. and Shi, Q. (2012) Inhibitors of phosphatidylinositol 3'-kinases promote mitotic cell death in HeLa cells. *PLoS ONE* **7**, e35665.
- Huang, Z. (2000) Small molecule inhibitors of Bcl-2 function: modulators of apoptosis and promising anticancer agents. *Curr. Opin. Drug Discov. Devel.* **3**, 565-574.
- Jimenez-Guerrero, R., Gasca, J., Flores, M. L., Perez-Valderrama, B., Tejera-Parrado, C., Medina, R., Tortolero, M., Romero, F., Japon, M. A. and Saez, C. (2018) Obatoclox and paclitaxel synergistically induce apoptosis and overcome paclitaxel resistance in urothelial cancer cells. *Cancers (Basel)* **10**, 490.
- Jung, C. H., Seo, M., Otto, N. M. and Kim, D. H. (2011) ULK1 inhibits the kinase activity of mTORC1 and cell proliferation. *Autophagy* **7**, 1212-1221.
- Jung, D., Khurana, A., Roy, D., Kalogera, E., Bakkum-Gamez, J., Chien, J. and Shridhar, V. (2018) Quinacrine upregulates p21/p27 independent of p53 through autophagy-mediated downregulation of p62-Skp2 axis in ovarian cancer. *Sci. Rep.* **8**, 2487.
- Kang, R., Zeh, H. J., Lotze, M. T. and Tang, D. (2011) The Beclin 1 network regulates autophagy and apoptosis. *Cell Death Differ.* **18**, 571-580.
- Kao, C., Chao, A., Tsai, C. L., Chuang, W. C., Huang, W. P., Chen, G. C., Lin, C. Y., Wang, T. H., Wang, H. S. and Lai, C. H. (2014) Bortezomib enhances cancer cell death by blocking the autophagic flux through stimulating ERK phosphorylation. *Cell Death Dis.* **5**, e1510.
- Kaushik, S. and Cuervo, A. M. (2018) The coming of age of chaperone-mediated autophagy. *Nat. Rev. Mol. Cell Biol.* **19**, 365-381.
- Kimmelman, A. C. and White, E. (2017) Autophagy and tumor metabolism. *Cell Metab.* **25**, 1037-1043.
- Kocaturk, N. M., Akkoc, Y., Kig, C., Bayraktar, O., Gozuacik, D. and Kutlu, O. (2019) Autophagy as a molecular target for cancer treatment. *Eur. J. Pharm. Sci.* **134**, 116-137.
- Koehler, B. C., Jassowicz, A., Scherr, A. L., Lorenz, S., Radhakrishnan, P., Kautz, N., Elssner, C., Weiss, J., Jaeger, D., Schneider, M. and Schulze-Bergkamen, H. (2015) Pan-Bcl-2 inhibitor obatoclox is a potent late stage autophagy inhibitor in colorectal cancer cells independent of canonical autophagy signaling. *BMC Cancer* **15**, 919.
- Kuma, A., Komatsu, M. and Mizushima, N. (2017) Autophagy-monitoring and autophagy-deficient mice. *Autophagy* **13**, 1619-1628.
- Lee, I. H., Cao, L., Mostoslavsky, R., Lombard, D. B., Liu, J., Bruns, N. E., Tsokos, M., Alt, F. W. and Finkel, T. (2008) A role for the NAD-dependent deacetylase Sirt1 in the regulation of autophagy. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 3374-3379.
- Li, J., Chong, T., Wang, Z., Chen, H., Li, H., Cao, J., Zhang, P. and Li, H. (2014) A novel anticancer effect of resveratrol: reversal of epithelial-mesenchymal transition in prostate cancer cells. *Mol. Med. Rep.* **10**, 1717-1724.
- Li, J. and Hochstrasser, M. (2020) Microautophagy regulates proteasome homeostasis. *Curr. Genet.* **66**, 683-687.
- Li, L. Q., Xie, W. J., Pan, D., Chen, H. and Zhang, L. (2016) Inhibition of autophagy by bafilomycin A1 promotes chemosensitivity of gastric cancer cells. *Tumour Biol.* **37**, 653-659.
- Li, Y. and Chen, Y. (2019) AMPK and autophagy. *Adv. Exp. Med. Biol.* **1206**, 85-108.
- Li, Z., Oh, D. Y., Nakamura, K. and Thiele, C. J. (2011) Perifosine-induced inhibition of Akt attenuates brain-derived neurotrophic factor/TrkB-induced chemoresistance in neuroblastoma *in vivo*. *Cancer* **117**, 5412-5422.
- Li, Z., Tan, F., Liewehr, D. J., Steinberg, S. M. and Thiele, C. J. (2010) *In vitro* and *in vivo* inhibition of neuroblastoma tumor cell growth by AKT inhibitor perifosine. *J. Natl. Cancer Inst.* **102**, 758-770.
- Liu, J., Long, S., Wang, H., Liu, N., Zhang, C., Zhang, L. and Zhang, Y. (2019) Blocking AMPK/ULK1-dependent autophagy promoted apoptosis and suppressed colon cancer growth. *Cancer Cell Int.* **19**, 336.
- Liu, L., McKeen, W. L., Wang, F. and Xie, R. (2012) MAP1S enhances autophagy to suppress tumorigenesis. *Autophagy* **8**, 278-280.
- Liu, M., Bamodu, O. A., Huang, W. C., Zucha, M. A., Lin, Y. K., Wu, A. T. H., Huang, C. C., Lee, W. H., Yuan, C. C., Hsiao, M., Deng, L., Tzeng, Y. M. and Yeh, C. T. (2017) 4-Acetyltantroquinonol B suppresses autophagic flux and improves cisplatin sensitivity in highly aggressive epithelial cancer through the PI3K/Akt/mTOR/p70S6K signaling pathway. *Toxicol. Appl. Pharm.* **325**, 48-60.
- Liu, W., Huang, S., Chen, Z., Wang, H., Wu, H. and Zhang, D. (2014) Temsirolimus, the mTOR inhibitor, induces autophagy in adenoid cystic carcinoma: *in vitro* and *in vivo*. *Pathol. Res. Pract.* **210**, 764-769.
- Lou, Z., Ren, T., Peng, X., Sun, Y., Jiao, G., Lu, Q., Zhang, S., Lu, X. and Guo, W. (2013) Bortezomib induces apoptosis and autophagy in osteosarcoma cells through mitogen-activated protein kinase pathway *in vitro*. *J. Int. Med. Res.* **41**, 1505-1519.
- Mahalingam, D., Mita, M., Sarantopoulos, J., Wood, L., Amaravadi, R. K., Davis, L. E., Mita, A. C., Curiel, T. J., Espitia, C. M., Nawrocki, S. T., Giles, F. J. and Carew, J. S. (2014) Combined autophagy and HDAC inhibition: a phase I safety, tolerability, pharmacokinetic, and pharmacodynamic analysis of hydroxychloroquine in combination with the HDAC inhibitor vorinostat in patients with advanced solid tumors. *Autophagy* **10**, 1403-1414.
- Marino, G., Salvador-Montoliu, N., Fueyo, A., Knecht, E., Mizushima, N. and Lopez-Otin, C. (2007) Tissue-specific autophagy alterations and increased tumorigenesis in mice deficient in Atg4C/autophagin-3. *J. Biol. Chem.* **282**, 18573-18583.
- Marquez, R. T. and Xu, L. (2012) Bcl-2:Beclin 1 complex: multiple mechanisms regulating autophagy/apoptosis toggle switch. *Am. J. Cancer Res.* **2**, 214-221.
- Mauthe, M., Orhon, I., Rocchi, C., Zhou, X., Luhr, M., Hijlkema, K. J., Coppes, R. P., Engedal, N., Mari, M. and Reggiori, F. (2018) Chloroquine inhibits autophagic flux by decreasing autophagosome-lysosome fusion. *Autophagy* **14**, 1435-1455.
- Mishra, A. K. and Dingli, D. (2019) Metformin inhibits IL-6 signaling by decreasing IL-6R expression on multiple myeloma cells. *Leukemia* **33**, 2695-2709.
- Mizushima, N. (2007) Autophagy: process and function. *Genes Dev.* **21**, 2861-2873.
- Momcilovic, M. and Shackelford, D. B. (2018) Imaging cancer metabolism. *Biomol. Ther. (Seoul)* **26**, 81-92.
- Morselli, E., Marino, G., Bennetzen, M. V., Eisenberg, T., Megalou, E., Schroeder, S., Cabrera, S., Benit, P., Rustin, P., Criollo, A., Kepp, O., Galluzzi, L., Shen, S., Malik, S. A., Maiuri, M. C., Horio, Y., López-Otín, C., Andersen, J. S., Tavernarakis, N., Madeo, F. and

- Kroemer, G. (2011) Spermidine and resveratrol induce autophagy by distinct pathways converging on the acetylproteome. *J. Cell Biol.* **192**, 615-629.
- Moscat, J. and Diaz-Meco, M. T. (2009) p62 at the crossroads of autophagy, apoptosis, and cancer. *Cell* **137**, 1001-1004.
- Mowers, E. E., Sharifi, M. N. and Macleod, K. F. (2017) Autophagy in cancer metastasis. *Oncogene* **36**, 1619-1630.
- Neri, L. M., Cani, A., Martelli, A. M., Simioni, C., Junghanss, C., Tabellini, G., Ricci, F., Tazzari, P. L., Pagliaro, P., McCubrey, J. A. and Capitani, S. (2014) Targeting the PI3K/Akt/mTOR signaling pathway in B-precursor acute lymphoblastic leukemia and its therapeutic potential. *Leukemia* **28**, 739-748.
- Park, J. M., Jung, C. H., Seo, M., Otto, N. M., Grunwald, D., Kim, K. H., Moriarity, B., Kim, Y. M., Starker, C., Nho, R. S., Voytas, D. and Kim, D. H. (2016) The ULK1 complex mediates MTORC1 signaling to the autophagy initiation machinery via binding and phosphorylating ATG14. *Autophagy* **12**, 547-564.
- Patel, S., Hurez, V., Nawrocki, S. T., Goros, M., Michalek, J., Sarantopoulos, J., Curiel, J. R. and Mahalingam, D. (2016) Vorinostat and hydroxychloroquine improve immunity and inhibit autophagy in metastatic colorectal cancer. *Oncotarget* **7**, 59087-59097.
- Periyasamy-Thanavan, S., Jackson, W. H., Samadder, J. S., Erickson, B., Barrett, J. R., Raney, L., Gopal, E., Ganapathy, V., Hill, W. D., Bhalla, K. N. and Schoenlein, P. V. (2010) Bortezomib blocks the catabolic process of autophagy via a cathepsin-dependent mechanism, affects endoplasmic reticulum stress and induces caspase-dependent cell death in antiestrogen-sensitive and resistant ER+ breast cancer cells. *Autophagy* **6**, 19-35.
- Petroni, G., Bagni, G., Iorio, J., Duranti, C., Lottini, T., Stefanini, M., Kragol, G., Becchetti, A. and Arcangeli, A. (2020) Clarithromycin inhibits autophagy in colorectal cancer by regulating the hERG1 potassium channel interaction with PI3K. *Cell Death Dis.* **11**, 161.
- Qu, X., Yu, J., Bhagat, G., Furuya, N., Hibshoosh, H., Troxel, A., Rosen, J., Eskelinen, E. L., Mizushima, N., Ohsumi, Y., Cattoretti, G. and Levine, B. (2003) Promotion of tumorigenesis by heterozygous disruption of the beclin 1 autophagy gene. *J. Clin. Invest.* **112**, 1809-1820.
- Racoma, I. O., Meisen, W. H., Wang, Q. E., Kaur, B. and Wani, A. A. (2013) Thymoquinone inhibits autophagy and induces cathepsin-mediated, caspase-independent cell death in glioblastoma cells. *PLoS ONE* **8**, e72882.
- Ramakrishnan, S., Nguyen, T. M., Subramanian, I. V. and Kelekar, A. (2007) Autophagy and angiogenesis inhibition. *Autophagy* **3**, 512-515.
- Rangwala, R., Leone, R., Chang, Y. C., Fecher, L. A., Schuchter, L. M., Kramer, A., Tan, K. S., Heitjan, D. F., Rodgers, G., Gallagher, M., Piao, S., Troxel, A. B., Evans, T. L., DeMichele, A. M., Nathanson, K. L., O'Dwyer, P. J., Kaiser, J., Pontiggia, L., Davis, L. E. and Amaravadi, R. K. (2014a) Phase I trial of hydroxychloroquine with dose-intense temozolomide in patients with advanced solid tumors and melanoma. *Autophagy* **10**, 1369-1379.
- Rangwala, R., Chang, Y. C., Hu, J., Algazy, K. M., Evans, T. L., Fecher, L. A., Schuchter, L. M., Torigian, D. A., Panosian, J. T., Troxel, A. B., Tan, K. S., Heitjan, D. F., DeMichele, A. M., Vaughn, D. J., Redlinger, M., Alavi, A., Kaiser, J., Pontiggia, L., Davis, L. E., O'Dwyer, P. J. and Amaravadi, R. K. (2014b) Combined MTOR and autophagy inhibition: phase I trial of hydroxychloroquine and temsirolimus in patients with advanced solid tumors and melanoma. *Autophagy* **10**, 1391-1402.
- Richardson, P. G., Eng, C., Kolesar, J., Hideshima, T. and Anderson, K. C. (2012) Perifosine, an oral, anti-cancer agent and inhibitor of the Akt pathway: mechanistic actions, pharmacodynamics, pharmacokinetics, and clinical activity. *Expert Opin. Drug Metab. Toxicol.* **8**, 623-633.
- Russell, R. C., Tian, Y., Yuan, H., Park, H. W., Chang, Y. Y., Kim, J., Kim, H., Neufeld, T. P., Dillin, A. and Guan, K. L. (2013) ULK1 induces autophagy by phosphorylating Beclin-1 and activating VPS34 lipid kinase. *Nat. Cell Biol.* **15**, 741-750.
- Russell, R. C., Yuan, H. X. and Guan, K. L. (2014) Autophagy regulation by nutrient signaling. *Cell Res.* **24**, 42-57.
- Saha, A., Blando, J., Tremmel, L. and DiGiovanni, J. (2015) Effect of metformin, rapamycin, and their combination on growth and progression of prostate tumors in HiMyc mice. *Cancer Prev. Res. (Phila.)* **8**, 597-606.
- Sell, S., Nicolini, A., Ferrari, P. and Biava, P. M. (2016) Cancer: a problem of developmental biology; scientific evidence for reprogramming and differentiation therapy. *Curr. Drug Targets* **17**, 1103-1110.
- Sesen, J., Dahan, P., Scotland, S. J., Saland, E., Dang, V. T., Lemarie, A., Tyler, B. M., Brem, H., Toulas, C., Cohen-Jonathan Moyal, E., Sarry, J. E. and Skuli, N. (2015) Metformin inhibits growth of human glioblastoma cells and enhances therapeutic response. *PLoS ONE* **10**, e0123721.
- Shukla, S., Patric, I. R., Patil, V., Shwetha, S. D., Hegde, A. S., Chandramouli, B. A., Arivazhagan, A., Santosh, V. and Somasundaram, K. (2014) Methylation silencing of ULK2, an autophagy gene, is essential for astrocyte transformation and tumor growth. *J. Biol. Chem.* **289**, 22306-22318.
- Su, Z., Yang, Z., Xu, Y., Chen, Y. and Yu, Q. (2015) Apoptosis, autophagy, necroptosis, and cancer metastasis. *Mol. Cancer* **14**, 48.
- Sun, F. D., Wang, P. C., Shang, J., Zou, S. H. and Du, X. (2018) Ibrutinib presents antitumor activity in skin cancer and induces autophagy. *Eur. Rev. Med. Pharmacol. Sci.* **22**, 561-566.
- Takamura, A., Komatsu, M., Hara, T., Sakamoto, A., Kishi, C., Waguri, S., Eishi, Y., Hino, O., Tanaka, K. and Mizushima, N. (2011) Autophagy-deficient mice develop multiple liver tumors. *Genes Dev.* **25**, 795-800.
- Tamargo-Gomez, I. and Marino, G. (2018) AMPK: regulation of metabolic dynamics in the context of autophagy. *Int. J. Mol. Sci.* **19**, 3812.
- Tripathi, R., Ash, D. and Shaha, C. (2014) Beclin-1-p53 interaction is crucial for cell fate determination in embryonal carcinoma cells. *J. Cell. Mol. Med.* **18**, 2275-2286.
- Tsujimoto, Y. and Shimizu, S. (2015) Another way to die: autophagic programmed cell death. *Cell Death Differ.* **12**, 1528-1534.
- Umehara, A., He, F., Taniguchi, K., Nakagawa, H., Yamachika, S., Font-Burgada, J., Zhong, Z., Subramanian, S., Raghunandan, S., Duran, A., Linares, J. F., Reina-Campos, M., Umehara, S., Valasek, M. A., Seki, E., Yamaguchi, K., Koike, K., Itoh, Y., Diaz-Meco, M. T., Moscat, J. and Karin, M. (2016) p62, upregulated during preneoplasia, induces hepatocellular carcinogenesis by maintaining survival of stressed HCC-initiating cells. *Cancer Cell* **29**, 935-948.
- Vignot, S., Faivre, S., Aguirre, D. and Raymond, E. (2005) mTOR-targeted therapy of cancer with rapamycin derivatives. *Ann. Oncol.* **16**, 525-537.
- Vogl, D. T., Stadtmauer, E. A., Tan, K. S., Heitjan, D. F., Davis, L. E., Pontiggia, L., Rangwala, R., Piao, S., Chang, Y. C., Scott, E. C., Paul, T. M., Nichols, C. W., Porter, D. L., Kaplan, J., Mallon, G., Bradner, J. E. and Amaravadi, R. K. (2014) Combined autophagy and proteasome inhibition: a phase 1 trial of hydroxychloroquine and bortezomib in patients with relapsed/refractory myeloma. *Autophagy* **10**, 1380-1390.
- Wan, B., Zang, Y. and Wang, L. (2018) Overexpression of Beclin1 inhibits proliferation and promotes apoptosis of human laryngeal squamous carcinoma cell Hep-2. *Onco Targets Ther.* **11**, 3827-3833.
- Wang, G., Zhou, P., Chen, X., Zhao, L., Tan, J., Yang, Y., Fang, Y. and Zhou, J. (2017) The novel autophagy inhibitor elaiophyllin exerts antitumor activity against multiple myeloma with mutant TP53 in part through endoplasmic reticulum stress-induced apoptosis. *Cancer Biol. Ther.* **18**, 584-595.
- Wang, Y., Xu, W., Yan, Z., Zhao, W., Mi, J., Li, J. and Yan, H. (2018) Metformin induces autophagy and G0/G1 phase cell cycle arrest in myeloma by targeting the AMPK/mTORC1 and mTORC2 pathways. *J. Exp. Clin. Cancer Res.* **37**, 63.
- White, E., Mehnert, J. M. and Chan, C. S. (2015) Autophagy, metabolism, and cancer. *Clin. Cancer Res.* **21**, 5037-5046.
- Xu, H. D. and Qin, Z. H. (2019) Beclin 1, bcl-2 and autophagy. *Adv. Exp. Med. Biol.* **1206**, 109-126.
- Yang, L., Wan, J., Xiao, S., Barkhouse, D., Zhu, J., Li, G., Lu, B. and Zhang, Z. (2016) BH3 mimetic ABT-737 sensitizes colorectal cancer cells to ixazomib through MCL-1 downregulation and autophagy inhibition. *Am. J. Cancer Res.* **6**, 1345-1357.
- Yang, Y. P., Hu, L. F., Zheng, H. F., Mao, C. J., Hu, W. D., Xiong, K. P., Wang, F. and Liu, C. F. (2013) Application and interpretation of



- current autophagy inhibitors and activators. *Acta Pharmacol. Sin.* **34**, 625-635.
- Yorimitsu, T. and Klionsky, D. J. (2005) Autophagy: molecular machinery for self-eating. *Cell Death Differ.* **12**, 1542-1552.
- Yu, H., Yin, S., Zhou, S., Shao, Y., Sun, J., Pang, X., Han, L., Zhang, Y., Gao, X., Jin, C., Qiu, Y. and Wang, T. (2018) Magnolin promotes autophagy and cell cycle arrest via blocking LIF/Stat3/Mcl-1 axis in human colorectal cancers. *Cell Death Dis.* **9**, 702.
- Yuan, N., Song, L., Zhang, S., Lin, W., Cao, Y., Xu, F., Fang, Y., Wang, Z., Zhang, H., Li, X., Wang, Z., Cai, J., Wang, J., Zhang, Y., Mao, X., Zhao, W., Hu, S., Chen, S. and Wang, J. (2015) Bafilomycin A1 targets both autophagy and apoptosis pathways in pediatric B-cell acute lymphoblastic leukemia. *Haematologica* **100**, 345-356.
- Zaffagnini, G., Savova, A., Danieli, A., Romanov, J., Tremel, S., Ebner, M., Peterbauer, T., Sztacho, M., Trapanone, R., Tarafder, A. K., Sachse, C. and Martens, S. (2018) p62 filaments capture and present ubiquitinated cargos for autophagy. *EMBO J.* **37**, e98308.
- Zanotto-Filho, A., Braganhol, E., Klafke, K., Figueiro, F., Terra, S. R., Paludo, F. J., Morrone, M., Bristot, I. J., Battastini, A. M., Forcelini, C. M., Bishop, A. J. R., Gelain, D. P. and Moreira, J. C. F. (2015) Autophagy inhibition improves the efficacy of curcumin/temozolomide combination therapy in glioblastomas. *Cancer Lett.* **358**, 220-231.
- Zhang, X., Li, W., Wang, C., Leng, X., Lian, S., Feng, J., Li, J. and Wang, H. (2014) Inhibition of autophagy enhances apoptosis induced by proteasome inhibitor bortezomib in human glioblastoma U87 and U251 cells. *Mol. Cell. Biochem.* **385**, 265-275.
- Zhao, X., Fang, Y., Yang, Y., Qin, Y., Wu, P., Wang, T., Lai, H., Meng, L., Wang, D., Zheng, Z., Lu, X., Zhang, H., Gao, Q., Zhou, J. and Ma, D. (2015) Elaiophyllin, a novel autophagy inhibitor, exerts anti-tumor activity as a single agent in ovarian cancer cells. *Autophagy* **11**, 1849-1863.
- Zhu, K., Dunner, K., Jr. and McConkey, D. J. (2010) Proteasome inhibitors activate autophagy as a cytoprotective response in human prostate cancer cells. *Oncogene* **29**, 451-462.