



ARTICLE

Antioxidant Properties and Diet-Related α -Glucosidase and Lipase Inhibitory Activities of Yogurt Supplemented with Safflower (*Carthamus tinctorius* L.) Petal Extract

Heeok Hong^{1,†}, Jeong Min Lim^{1,†}, Damini Kothari¹, So Hee Kwon¹,
Hyuk Cheol Kwon², Sung-Gu Han², and Soo-Ki Kim^{1,*}



OPEN ACCESS

Received April 10, 2020

Revised September 11, 2020

Accepted November 4, 2020

¹Department of Animal Science and Technology, Konkuk University, Seoul 05029, Korea

²Department of Food Science and Biotechnology of Animal Resource, Konkuk University, Seoul 05029, Korea

*Corresponding author : Soo-Ki Kim
Department of Animal Science and
Technology, Konkuk University,
Seoul 05029, Korea
Tel: +82-2-450-3728
Fax: +82-2-458-3728
E-mail: sookikim@konkuk.ac.kr

*ORCID
Heeok Hong
<https://orcid.org/0000-0001-6315-7097>
Jeong-Min Lim
<https://orcid.org/0000-0001-5071-961X>
Damini Kothari
<https://orcid.org/0000-0003-3627-2377>
So Hee Kwon
<https://orcid.org/0000-0002-6066-543X>
Hyuk Cheol Kwon
<https://orcid.org/0000-0001-6234-2530>
Sung-Gu Han
<https://orcid.org/0000-0002-1485-861X>
Soo-Ki Kim
<https://orcid.org/0000-0003-3499-3330>

† These authors contributed equally to this work.

Abstract Recently, yogurt has been extensively studied to further enhance its functions using edible plant extracts. This study was conducted to investigate whether safflower petal (SP) as a natural food additive can be used to develop functional yogurt with improved health benefits. SPs were extracted with ethanol (SPE) and hot water (SPW), and then safflower yogurt was prepared by adding 0%–1.0% of those extracts to plain yogurt. With an increase in the fermentation duration, the pH of SPE and SPW yogurt samples was decreased, whereas titratable acidity and microbial counts were increased. The concentration of total polyphenols and total flavonoids, the activity of antioxidants, and the inhibitory effect on reactive oxygen species (ROS) were higher in SPW yogurt than SPE yogurt. Furthermore, α -glucosidase and lipase activity inhibitory effects of SPW yogurt were higher than those of SPE yogurt. In particular, free radical-scavenging activities, ROS inhibitory effect, and α -glucosidase activity inhibitory effects were significantly increased in SPW yogurt in a dose-dependent manner. Overall, these results suggest that SP extract possesses antioxidant activities and that it can downregulate α -glucosidase and lipase activities. The SP extract may have potential benefits as a natural food additive for the development of functional yogurt.

Keywords yogurt, safflower petal, α -glucosidase, lipase, antioxidant activity

Introduction

Diabetes mellitus is a serious chronic metabolic disorder, associated with life-threatening complications (Elbashir et al., 2018). It is characterized by a state of chronically elevated blood glucose levels (hyperglycemia) with altered nutrient metabolism, such as carbohydrates, fats, and proteins, in the body (American Diabetes Association, 2014). Moreover, the oxidative stress state associated with the generation of reactive oxygen species (ROS) increases in both type I and type II diabetes (Omari

et al., 2019; Sabu and Kuttan, 2002). ROS generated during oxidative metabolism can cause various disorders.

α -Glucosidase enzyme inhibitors, as antidiabetic drugs, can lower blood glucose levels by reducing the intestinal absorption of carbohydrates (Buchholz and Melzig, 2016). Additionally, hyperlipidemia is a common feature in patients with diabetes mellitus and is one of the factors that is associated with considerably increases the risk of premature atherosclerosis (Schofield et al., 2016). It is known that pancreatic lipase regulates the digestion and absorption of dietary lipids (Mhatre et al., 2019). Therefore, inhibition of this enzyme may help prevent complications by regulating lipid levels in the blood of diabetic patients through reducing dietary fat absorption into the body.

Despite the growing interest in functional foods that modulate physiological activities by inhibiting these enzymes, the possibility of developing successful and targeted natural products to safely manage these diseases remains unexplored.

Yogurt, obtained by fermenting milk using lactic acid bacteria, contains proteins, vitamins such as vitamins A and B, and minerals such as calcium and manganese. Recently, yogurt has been extensively studied to further enhance its functions using bioactive substances from edible plants. Numerous studies have indicated that the addition of these substances increases the concentration of polyphenols and flavonoids, thus increasing the antioxidant and antimicrobial activities of yogurt (Halal and Nayra, 2011; Kang et al., 2018; Lim, 2018).

Carthamus tinctorius L., commonly known as safflower, belonging to the Asteraceae family (Delshad et al., 2018), is an annual herbaceous crop. It has been cultivated in India, Iran, and China for a long time (Asgarpanah and Kazemivash, 2013). Safflower is used as a traditional herbal medicine to promote blood circulation, regulate menstruation, and alleviate pain of joints (Delshad et al., 2018). Some studies have reported that safflower has more than 200 compounds including flavonoids, phenylethanoid glycosides, coumarins, fatty acids, steroids, and polysaccharides (Asgarpanah and Kazemivash, 2013). Furthermore, in *in vitro* and *in vivo* studies, hydroxysafflower yellow A, a major bioactive compound in safflower, has been found to have a strong oxidative radical-scavenging effect (Tian et al., 2008; Zhao et al., 2018).

The present study has focused on whether the addition of safflower extract containing various phytochemicals to yogurt is effective in the antioxidant activity and inhibition of α -glucosidase and pancreatic lipase. Therefore, this study was conducted to investigate the biochemical, microbiological, sensory characteristics, and physiological effects including α -glucosidase and pancreatic lipase activity inhibitory effects of yogurt supplemented with safflower petal (SP) extract in order to enhance the functionality of yogurt.

Materials and Methods

Plant materials and extract preparation

Dried SPs were purchased from a local market (Daejodong, Seoul, Korea). For ethanol extraction (SPE), 100 g of dried SPs was precipitated with 900 mL of 99.9% ethanol in a shaking incubator at room temperature for 15 h. For water extraction (SPW), 100 g of dried SPs was soaked in 900 mL of 100°C hot water for 1 h using a reflux condenser (Reflux-A, Reservoir; LK Lab Korea, Namyangju, Korea). These extracts were centrifuged at 1,889×g for 15 min; then, the supernatant was collected and passed through qualitative filter paper No. 1 (Toyo Roshi Kaisha, Tokyo, Japan) and 0.22 μ m syringe filter (Sartorius, Goettingen, Germany). The aqueous extract was concentrated using a vacuum freeze-dryer at -80°C (Unifreez™ FD-8; Daihan Scientific, Wonju, Korea). The concentration of SPE and SPW was 50 mg/mL.

Preparation of yogurt supplemented with SP extract

SPE and SPW were added to sterilized milk (Maeil Dairies, Seoul, Korea) at concentrations of 0%–1.0% and homogenized

using Homogenizer T 25 (Ika, Staufen, Germany) at 25°C for 10 min, and then pasteurized at 85°C for 15 min. Thereafter, the samples were immediately cooled to 43°C and inoculated with a commercial yogurt starter culture (0.02%, v/v) of lactic acid bacteria (Lyofast YAB 450 AB; Sacco s.r.l., Codaragok, Italy) containing *Streptococcus thermophilus*, *Lactobacillus delbrueckii* spp. *bulgaricus*, *Lactobacillus acidophilus*, and *Bifidobacterium animalis* spp. *lactis*. Yogurt added with 0% safflower extract was used as control. These yogurt samples were fermented at 43°C for 6 h and stored at 4°C until further analysis.

Proximate composition, pH, titratable acidity, and viscosity

Moisture and total ash content was measured using the methods of Association of Official Analytical Chemists (2000). The concentration of proteins, crude fats, lactose, and total solids was measured using Mikloscan (Milkoscan Minor 78110; Foss, Hillerød, Denmark). The pH of yogurt samples was measured using a digital pH meter (Model 735P; IS Technology, Incheon, Korea). Titratable acidity (TA) was determined by adding 9 mL of dH₂O to 1 g of yogurt sample and then titrating the yogurt mixture to pH 8.3 with 0.1 N NaOH. The amount of acid produced was calculated as follows:

$$\text{TA (\% lactic acid)} = \text{dilution factor} \times V_{\text{NaOH}} \times 0.1 \text{ N} \times 0.009 \times 100$$

where, V_{NaOH} is the volume of NaOH required to neutralize the acid.

Viscosity of yogurt supplemented with SP was measured using a viscometer (Model LVDV 1+; Brookfield Engineering Laboratories, Middleboro, MA, USA) with spindle No. 63 at 50 rpm for 5–8 min.

Viable cell count

Viable cell count in yogurt was determined on a *Streptococcus thermophilus* agar (Sigma-Aldrich, St. Louis, MO, USA) plate after incubating at 37°C for 48 h. The results are expressed as CFU/mL.

2,2-Diphenyl-2-picrylhydrazyl radical-scavenging activity assay

2,2-Diphenyl-2-picrylhydrazyl (DPPH) inhibition was determined as described previously (Kang et al., 2018). Briefly, each yogurt extract (16 µL) was mixed with 320 µL of 0.15 mM DPPH solution (Sigma-Aldrich). The mixture was shaken and incubated at room temperature for 15 min. The absorbance of the solution was measured at 515 nm using an ELISA reader (Synergy 2; BioTek Instruments, Winooski, VT, USA). Inhibition of DPPH oxidation (%) was calculated as follows:

$$\% \text{ Inhibition} = (A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}} \times 100$$

where, A is the absorbance at 515 nm.

2,2'-Azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) radical-scavenging activity assay

2,2'-Azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) radical-scavenging activity was determined using the method of Kang et al. (2018). The stock solution comprised 7.4 mM ABTS (Sigma-Aldrich) and 2.5 mM potassium persulfate solutions. The mixture was prepared by mixing the two stock solutions at a ratio of 1:1 (v/v) and allowed to react

for 16 h in dark. The ABTS solution was diluted with tertiary dH₂O to the appropriate absorbance (7.00±0.05), which was measured at 734 nm. The ABTS solution (180 µL) was mixed with 20 µL of yogurt extract, and the mixture was covered with aluminum foil and placed in the dark at room temperature for 30 min. The absorbance of the mixture was measured at 734 nm using an ELISA reader (Synergy 2; BioTek Instruments). Each sample was measured in triplicate, and percent inhibition was calculated using the following equation:

$$\% \text{ Inhibition} = (A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}} \times 100$$

where, *A* is the absorbance at 734 nm.

Total polyphenol and total flavonoid content

Total phenolic content (TPC) was determined using the method of Wei et al. (2011) with modifications. Briefly, each yogurt extract (100 µL) was mixed with 100 µL of 1 N Folin–Ciocalteu reagent (Sigma-Aldrich); this mixture was thoroughly mixed and incubated at room temperature for 3 min. Na₂CO₃ (200 µL, 1 N) was added to the mixture, which was allowed to react at room temperature for 90 min. The developed color in the mixture was measured using an ELISA reader (Synergy 2; BioTek Instruments) at 725 nm. The TPC is expressed as gallic acid equivalent (Sigma-Aldrich). The regression line of the gallic acid standard was used to determine the TPC in yogurt extract as micrograms of gallic acid equivalent per milliliter (µg GAE/mL).

Total flavonoid content (TFC) was measured using the method of Abeysinghe et al. (2007) with modifications. Briefly, each yogurt extract (100 µL) was mixed with diethylene glycol (500 µL) and NaOH (50 µL, 1 N). The mixture was incubated at 37°C for 60 min. The developed color in the mixture measured using an ELISA reader (Synergy 2; BioTek Instruments) at 420 nm. The TFC is expressed as quercetin equivalent (Sigma-Aldrich). A standard curve was generated using the quercetin standard solution, and the results are expressed as quercetin equivalent per milliliter.

Cell lines and cell culture

The human colorectal cell line (HT-29) used in this study was purchased from the American Type Culture Collection (Manassas, VA, USA). HT-29 cells were cultured in RPMI 1640 medium (Lonza, Walkersville, MD, USA) supplemented with 10% fetal bovine serum (Atlas Biologicals, Fort Collins, CO, USA), 100 U/mL penicillin (Gibco, Grand Island, NY, USA), and 100 µg/mL streptomycin (Gibco) at 37°C under 5% CO₂. The cells were re-fed with the medium every 2 d until 80% confluence.

Measurement of ROS

ROS were measured using the stain 2',7'-dichlorofluorescein diacetate (DCFDA, Sigma-Aldrich). Briefly, the cells were seeded in six-well plates, incubated overnight, and then treated with diluted yogurt extract for 16 h. Thereafter, the cells were treated with 1 µg/mL lipopolysaccharide (Sigma-Aldrich)-containing medium for 6 h. The cells were then incubated with DCFDA at a final concentration of 10 µM for 30 min and washed twice with cold PBS. The plates were subsequently measured using the fluorescence microscope (Model IX71, Olympus Optical, Tokyo, Japan). Photographs were captured using the Olympus DP71 camera and DP controller software (Olympus Optical). Finally, ROS were quantified using Image J

software (National Institute of Health, Bethesda, Rockville, MD, USA).

Measurement of α -glucosidase inhibition

α -Glucosidase activity inhibitory effect was determined using a modified assay reported by Kwon et al. (2006). Briefly, 50 μ L of yogurt extract was added to 100 μ L of 0.1 M phosphate buffer (pH 6.9) containing α -glucosidase solution (1 U/mL, Sigma-Aldrich) in 96-well plates. The plates were incubated at 25°C for 10 min. After pre-incubation, 50 μ L of 5 mM *p*-nitrophenyl- α -D-glucopyranoside solution in 0.1 M phosphate buffer (pH 6.9) was added to each well. The mixture was incubated at 25°C for 5 min. The absorbance of the mixture was measured at 405 nm before and after incubation using an ELISA reader (Synergy 2; BioTek Instruments). The absorbance of yogurt extract was compared with that of the control, which was 50 μ L of buffer solution. The α -glucosidase activity inhibition is expressed as percent inhibition and was calculated using the following equation:

$$\% \text{ Inhibition} = [(A_{\text{control}} - A_{\text{extract}}) / A_{\text{control}}] \times 100$$

where, A is the absorbance at 405 nm.

Measurement of pancreatic lipase inhibition

Pancreatic lipase inhibition was determined using the modified method of Kim et al. (2007) using *p*-nitrophenyl butyrate (NPB). Briefly, the enzyme solution was prepared by adding 30 μ L of lipase solution from porcine pancreas Type VI-S (Sigma-Aldrich). This solution was composed of 10 mM morpholinepropanesulfonic acid and 1 mM ethylenediaminetetraacetic acid (pH 6.8) in 850 μ L of Tris buffer (100 mM Tris-HCl and 5 mM CaCl₂, pH 7.0). Yogurt extract (100 μ L) was mixed with an enzyme substrate solution (880 μ L) containing 10 mM *p*-NPB and incubated for 15 min at 37°C. The absorbance of the sample was measured at 400 nm using an ELISA reader (Synergy 2; BioTek Instruments) and was compared with that of the control buffer solution, without the extract. The lipase activity inhibition is expressed as percent inhibition and was calculated using the following equation:

$$\% \text{ Inhibition} = [(A_{\text{control}} - A_{\text{extract}}) / A_{\text{control}}] \times 100$$

where, A is the absorbance at 400 nm.

Statistical analysis

All data are expressed as mean \pm SD of three experiments. Statistical analysis of the data was performed using the one-way analysis of variance (ANOVA; SPSS 25, SPSS, Chicago, IL, USA) followed by Duncan's multiple range test and Student *t*-test for comparison of means. Statistical significance was set at $p < 0.05$.

Results and Discussion

Proximate composition of yogurt supplemented with SP extract

The nutritional composition of yogurt supplemented with SPE and SPW is shown in Table 1. The moisture content of SPE

Table 1. Proximate analysis of yogurt supplemented with safflower petal extract

Component	Concentration of safflower petal extract (%)	Total amount (%)	
		SPE	SPW
Moisture	0.0	83.19±0.34 ^b	83.19±0.34 ^b
	0.1	82.13±0.14 ^a	82.59±0.36 ^b
	0.5	81.99±0.53 ^a	81.41±0.05 ^a
	1.0	81.67±0.32 ^a	81.42±0.06 ^a
Ash	0.0	0.80±0.02	0.80±0.02
	0.1	0.80±0.03	0.81±0.03
	0.5	0.81±0.02	0.82±0.02
	1.0	0.81±0.03	0.81±0.03
Protein	0.0	3.95±0.11 ^a	3.95±0.11
	0.1	4.33±0.15 ^{bB}	3.93±0.25 ^A
	0.5	4.37±0.10 ^{bB}	4.01±0.15 ^A
	1.0	4.40±0.10 ^{bB}	3.98±0.06 ^A
Fat	0.0	3.63±0.06	3.63±0.06 ^b
	0.1	3.63±0.15	3.40±0.10 ^a
	0.5	3.70±0.26	3.73±0.16 ^b
	1.0	3.73±0.21	3.73±0.23 ^b
Lactose	0.0	11.77±0.06 ^a	11.77±0.06 ^a
	0.1	12.01±0.26 ^a	12.03±0.06 ^b
	0.5	13.02±0.06 ^b	13.07±0.12 ^c
	1.0	13.11±0.06 ^{bB}	13.07±0.06 ^{cA}
Total solids	0.0	15.75±0.45	15.75±0.45
	0.1	15.89±0.36	15.76±1.01
	0.5	15.95±0.59	16.04±0.68
	1.0	16.05±0.44	16.02±0.21

Values are mean±SD (n=3).

Different small letters in the same column and capitalized letters in the same row are significantly different (p<0.05).

SPE, safflower petal ethanol extract; SPW, safflower petal hot water extract.

yogurt increased with an increase in the concentration of SPE added. The moisture content of SPE yogurt was 81.67%–82.13%, which was significantly lower than that of plain yogurt (p<0.05). In SPE yogurt, the concentration of proteins, fats, lactose, and total solids increased with an increase in the concentration of SPE. The lactose concentrations in 0.5% and 1.0% SPE yogurt were 13.02% and 13.11%, which were significantly higher than those in 0% and 0.1% SPE (p<0.05). The concentration of nutrients in SPE yogurt tended to be higher than that in SPW yogurt. In particular, the protein concentrations in SPE yogurt were 4.33%–4.40%, which were significantly higher than those of SPW yogurt (p<0.05). The supplementation of yogurt with red ginseng extract changed the composition of yogurt, that is, with an increase in red ginseng extract concentration, the moisture content and protein and lactose concentrations increased (Jung et al., 2016). These findings are similar to our data. According to Pires et al. (2018), the supplementation of various plant petal extracts to yogurt marginally

increased the nutritional composition such as protein and lactose concentrations, except the moisture content. Overall, yogurt composition was affected by the addition of SP extract.

Changes in pH, TA, microbial count, and viscosity of yogurt with SP extract

As shown in Fig. 1A and 1D, the pH of SPE and SPW yogurt decreased significantly during fermentation ($p < 0.05$). In contrast, the TA level in SPE and SPW yogurt was significantly increased during fermentation as shown in Fig. 1B and 1E ($p < 0.05$). The pH and TA of all yogurt samples were similar during the initial and final stages of fermentation. These results are consistent with those of previous studies, that is, the pH decreased but the TA level increased due to acid production in




Fig. 1. Changes in pH, titratable acidity, viable cell counts, and viscosity during fermentation of yogurt supplemented with safflower petal extract. (A, D) pH, (B, E) titratable acidity, (C, F) viable cell counts, and (G, H) viscosity. (A–C) ethanol extracts: ●, 0%; ○, 0.1%; ▼, 0.5%; △, 1.0%. (D–F) hot water extracts: ●, 0%; □, 0.1%; ■, 0.5%; ◇, 1.0%. Different uppercase letters indicate a significant difference ($p < 0.05$).

red ginseng extract- and ginseng extract-supplemented yogurt (Jang et al., 2018; Jung et al., 2016).

The viable cell counts in SPE and SPW yogurt during fermentation are shown in Fig. 1C and 1F. They ranged from 7.49 to 9.77 Log₁₀ (CFU/g) and 7.43 to 9.99 Log₁₀ (CFU/g), respectively. The number of viable cells in SPE was not significantly different from that in SPW yogurt. However, the number of viable cells in SPE and SPW yogurt was slightly lower than that in plain yogurt at the end of fermentation. It has been reported that in yogurt supplemented with cinnamon ethanol extract, the viable cell counts were marginally decreased at the end of fermentation (Choi et al., 2016). However, yogurt fermented with Korean traditional plant extracts showed an increase in lactic acid bacteria count during fermentation (Joung et al., 2016; Jung et al., 2016).

The viscosity of yogurt supplemented with SP extracts at different concentrations is shown in Fig. 1G and H. The viscosity of yogurt with SPE and SPW tended to be slightly lower than that of plain yogurt. The viscosity of 1.0% SPW and 0.5% SPE yogurt was significantly decreased compared with plain yogurt ($p < 0.05$). It is known that viscosity is related to the aggregation of casein micelles (Joyce et al., 2017). Sung et al. (2015) reported that the viscosity of yogurt supplemented with 5% freeze-dried mulberry fruit juice was significantly lower than that of the control. Tseng and Zhao (2013) reported that the viscosity of yogurt supplemented with wine grape pomace was decreased; they suggested that this was probably because the addition of a high concentration of wine grape pomace into yogurt resulted in the breakdown of the coagulated milk. Furthermore, it has been reported that carthamin yellow from safflower significantly decreased whole blood viscosity, plasma viscosity, and erythrocyte aggregation index, in an animal study (Li et al., 2009).

Antioxidant activity, TPC, and TFC

As shown in Table 2, the DPPH radical-scavenging activity of SPW yogurt was 5.81%–10.66%, which was considerably higher than that of SPE yogurt, which was 1.03%–3.24%. Moreover, the ABTS radical-scavenging activity of SPW yogurt was 40.00%–49.67%, but the activity was not detected in SPE yogurt. These results are consistent with those of Kang et al. (2018), who reported that the antioxidant activity of yogurt supplemented with water extract of fermented pepper was higher than that of methanol extract. Here, the antioxidant activities of yogurt seemed to increase with the concentration of SPE and SPW. In particular, the antioxidant activity of 1.0% SPW yogurt was the highest, and it was significantly higher than that of plain yogurt ($p < 0.05$). These results can be attributed to the TPC and TFC in the SP. Similar to the antioxidant effects, the TPC and TFC in yogurt increased with the increase in the concentration of SPW and SPE. The TPC and TFC in yogurt supplemented with SPW 1.0% were 393.13 and 320.90 µg/mL, respectively, which were the highest among the yogurt samples ($p < 0.05$). The TPC in yogurt supplemented with SPE and SPW was 93.55–115.12 and 282.52–393.19 µg/mL, respectively, and the TFC was 52.53–77.76 and 180.97–320.90 µg/mL, respectively. It is known that edible safflower yellow pigments, which belong to the C-glucosylquinochalcone family of flavonoids, are water soluble (Li et al., 2018). This suggests that hot water extracted more bioactive substances such as polyphenols and flavonoids from SP, in this study. Kruawan and Kangsadalampai (2006) reported that the water extract of *Carthamus tinctorius* L. presented high antioxidant activity and TPC. In summary, the results demonstrated that SP extracted with hot water had greater antioxidant capacity, as evidenced by an increase in the TPC and TFC.

Inhibition of ROS production in LPS-stimulated human colon adenocarcinoma

The effect of SPE yogurt on intracellular ROS levels was determined in order to elucidate its antioxidant activity, through which SPE can protect cells against LPS-induced damage. LPS, a major component of bacterial cell walls, induces the

Table 2. Antioxidant activity of, and total polyphenol content (TPC) and total flavonoid content (TFC) in yogurt supplemented with safflower petal extract

	Concentration of safflower petal extract (%)	Total amount	
		SPE	SPW
TPC ($\mu\text{g/mL}$)	0.0	115.12 \pm 28.47	282.52 \pm 4.47 ^a
	0.1	93.55 \pm 37.28	380.52 \pm 8.60 ^b
	0.5	94.49 \pm 30.49	381.46 \pm 8.61 ^b
	1.0	103.86 \pm 37.17	393.19 \pm 0.81 ^b
TFC ($\mu\text{g/mL}$)	0.0	52.53 \pm 10.00 ^a	180.97 \pm 30.97 ^a
	0.1	57.12 \pm 3.98 ^{ab}	210.80 \pm 2.29 ^b
	0.5	73.17 \pm 4.59 ^{bc}	270.44 \pm 11.92 ^c
	1.0	77.76 \pm 3.97 ^c	320.90 \pm 2.29 ^d
Antioxidant activity			
DPPH (%)	0.0	3.24 \pm 0.62	5.81 \pm 0.61 ^a
	0.1	1.03 \pm 0.58	6.69 \pm 0.78 ^a
	0.5	2.60 \pm 0.20	7.17 \pm 3.94 ^a
	1.0	2.79 \pm 0.85	10.66 \pm 1.21 ^b
ABTS (%)	0.0	ND	40.00 \pm 1.54 ^a
	0.1	ND	48.00 \pm 3.00 ^b
	0.5	ND	48.33 \pm 1.76 ^b
	1.0	ND	49.67 \pm 0.88 ^b

Values are mean \pm SD (n=3).

Different superscripts in the same column are significantly different (p<0.05).

SPE, safflower petal ethanol extract; SPW, safflower petal hot water extract; DPPH, 2,2-diphenyl-2-picrylhydrazyl; ABTS, 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid); ND, not detected.

production of inflammatory mediators such as ROS (Lee et al., 2014). ROS play an important role in oxidative stress associated with the development of chronic diseases such as cancer and type II diabetes (Lee et al., 2014). Therefore, this study was focused on LPS-induced ROS production *in vitro*. As shown in Fig. 2, SPE and SPW yogurt significantly reduced ROS production in LPS-induced HT-29 cells (p<0.001). Ma et al. (2016), based on their study in LPS-induced hepatocellular (HepG2) cells, reported that SP yellow B possesses a high ROS-scavenging activity. As mentioned above, edible safflower yellow pigments are water soluble, and therefore, the ROS-scavenging ability of SPW yogurt with a high TPA and TFC was higher than that of SPE yogurt. Thus, these results demonstrated the antioxidant effect of SP extract-supplemented yogurt *in vitro*.

α -Glucosidase and lipase activity inhibitory effects

It is well known that α -glucosidase and pancreatic lipase are involved in the digestion and absorption of carbohydrates and lipids, respectively (Buchholz and Melzig, 2016; Elbashir et al., 2018). In this study, the activities of these enzymes were measured to determine whether SP could regulate enzymes related to diabetic complications such as hyperglycemia and hyperlipidemia. As shown Fig. 3B, SPW yogurt significantly inhibited the α -glucosidase activity in a dose-dependent manner (p<0.05). Furthermore, the inhibitory effect of SPW yogurt on pancreatic lipase activity was the highest at a concentration of 0.5% (Fig. 3D). However, these enzyme inhibitory effects of SPE yogurt were significantly lower than those of plain yogurt,




Fig. 2. Antioxidant effects of yogurt supplemented with safflower petal extract on human colorectal cells. Fluorescence microscopic image (A). The bar figure was calculated using Image J software (B, C). SPE, safflower petal ethanol extract; SPW, safflower petal hot water extract.




Fig. 3. α-Glucosidase and porcine pancreatic lipase activity inhibitory effects of yogurt supplemented with different concentrations of safflower petal extract. (A, C) safflower petal ethanol extract, (B, D) safflower petal hot water extract. Different uppercase letters indicate a significant difference ($p < 0.05$).

except the α -glucosidase activity inhibitory effect of 0.1% SPE yogurt (Fig. 3A and C; $p < 0.05$). The α -glucosidase activity inhibitory effect of 0.1% SPE yogurt was significantly increased compared with yogurt samples with other concentrations of SPE ($p < 0.05$). Several studies have suggested that various plants inhibit the activities of α -glucosidase and pancreatic lipase (Elbashir et al., 2018; Gholamhoseinian et al., 2008; Yoshioka et al., 2019). Elbashir et al. (2018) suggested that the strong α -glucosidase activity inhibitory effect of these plants is related to their antioxidant capacities due to the presence of bioactive compounds including polyphenols, flavonoids, and tannins. In this study, SPW yogurt had more TPC and TFC, and showed higher free radical-scavenging activities than SPE yogurt. Therefore, SP extract-supplemented yogurt might alleviate hyperglycemia and hyperlipidemia by downregulating the digestion and absorption of carbohydrates and lipids.

Taken together, the number of viable LAB in safflower yogurt was similar to that the plain yogurt, which was more than Korea Food Standard (1×10^7 CFU/mL) and the pH was 4.26–4.31, which were similar to that of the plain yogurt (pH 4.3). These results indicated that SP extract-supplemented yogurt had no negative effects on yogurt characteristics compared with plain yogurt. The addition of SP extract resulted in an increase in the TPC and TFC, which increased the antioxidant activity of SP extract-supplemented yogurt. The inhibitory effect on enzymes that are involved in carbohydrate and lipid digestion and absorption resulted in a positive effect on the regulation of hyperglycemia and hyperlipidemia. In addition, although the changes in the storage characteristics of SP extract-supplemented yogurt were not evaluated in this study, they were considered stable because there were no significant negative changes in the quality of yogurt stored at 4°C for 21 d. In conclusion, SP extract has potential application in the production of high value-added products, as it can enhance health benefits of various foods including yogurt. Further animal studies are needed to confirm the action mechanisms and *in vivo* effects to validate the potential benefits of yogurt supplemented with SP extract in managing weight and blood glucose.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This paper was supported by Konkuk University in 2018.

Author Contributions

Conceptualization: Hong H, Lim JM, Kwon SH, Kwon HC, Han SG, Kim SK. Data curation: Hong H, Lim JM, Kwon SH, Kwon HC, Han SG, Kim SK. Formal analysis: Lim JM, Kwon SH, Kwon HC. Methodology: Hong H, Lim JM, Kothari D, Kwon SH. Software: Lim JM, Kwon SH. Validation: Hong H, Kim SK. Investigation: Hong H, Lim JM, Kothari D, Kwon SH, Kwon HC, Han SG, Kim SK. Writing - original draft: Hong H, Lim JM, Kim SK. Writing - review & editing: Hong H, Lim JM, Kothari D, Kwon SH, Kwon HC, Han SG, Kim SK.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Abeysinghe DC, Li X, Sun C, Zhang W, Zhou C, Chen K. 2007. Bioactive compounds and antioxidant capacities in different edible tissues of citrus fruit of four species. *Food Chem* 104:1338-1344.
- American Diabetes Association. 2014. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 37:S81-S90.
- AOAC. 2000. Official methods of analysis of AOAC International. 17th ed. AOAC International, Gaithersburg, MD, USA.
- Asgarpanah J, Kazemivash N. 2013. Phytochemistry, pharmacology and medicinal properties of *Carthamus tinctorius* L. *Chin J Integr Med* 19:153-159.
- Buchholz T, Melzig MF. 2016. Medicinal plants traditionally used for treatment of obesity and diabetes mellitus-screening for pancreatic lipase and α -amylase inhibition. *Phytother Res* 30:260-266.
- Choi YJ, Jin HY, Yang HS, Lee SC, Huh CK. 2016. Quality and storage characteristics of yogurt containing *Lactobacillus sakei* ALI033 and cinnamon ethanol extract. *J Anim Sci Technol* 58:16.
- Delshad E, Yousefi M, Sasannezhad P, Rakhshandeh H, Ayati Z. 2018. Medical uses of *Carthamus tinctorius* L. (Safflower): A comprehensive review from traditional medicine to modern medicine. *Electron Physician* 10:6672-6681.
- Elbashir SMI, Devkota HP, Wada M, Kishimoto N, Moriuchi M, Shuto T, Misumi S, Kai H, Watanabe T. 2018. Free radical scavenging, α -glucosidase inhibitory and lipase inhibitory activities of eighteen Sudanese medicinal plants. *BMC Complement Altern Med* 18:282.
- Gholamhoseinian A, Fallah H, Sharifi-far F, Mirtajaddini M. 2008. The inhibitory effect of some Iranian plants extracts on the alpha glucosidase. *Iranian J Basic Med Sci* 11:1-9.
- Halah MF, Nayra SM. 2011. Use of natural plant antioxidant and probiotic in the production of novel yogurt. *J Evol Biol Res* 3:12-18.
- Jang HJ, Jung J, Yu HS, Lee NK, Paik HD. 2018. Evaluation of the quality of yogurt using ginseng extract powder and probiotic *Lactobacillus plantarum* NK181. *Korean J Food Sci Anim Resour* 38:1160-1167.
- Joung JY, Lee JY, Ha YS, Shin YK, Kim Y, Kim SH, Oh NS. 2016. Enhanced microbial, functional and sensory properties of herbal yogurt fermented with Korean traditional plant extracts. *Korean J Food Sci Anim Resour* 36:90-99.
- Joyce AM, Brodkorb A, Kelly AL, O'Mahony JA. 2017. Separation of the effects of denaturation and aggregation on whey-casein protein interactions during the manufacture of a model infant formula. *Dairy Sci Technol* 96:787-806.
- Jung J, Paik HD, Yoon HJ, Jang, HJ, Jeewanthi RKC, Jee HS, Li X, Lee NK, Lee SK. 2016. Physicochemical characteristics and antioxidant capacity in yogurt fortified with red ginseng extract. *Korean J Food Sci Anim Resour* 36:412-420.
- Kang SH, Yu MS, Kim JM, Park SK, Lee CH, Lee HG, Kim SK. 2018. Biochemical, microbiological, and sensory characteristics of stirred yogurt containing red or green pepper (*Capsicum annuum* cv. Chungyang) juice. *Korean J Food Sci Anim Resour* 38:451-467.
- Kim JH, Kim HJ, Park HW, Youn SH, Choi DY, Shin CS. 2007. Development of inhibitors against lipase and α -glucosidase from derivatives of monascus pigment. *FEMS Microbiol Lett* 276:93-98.
- Kruawan K, Kangsadalampai K. 2006. Antioxidant activity, phenolic compound contents and antimutagenic activity of some water extract of herbs. *Thai J Pharm Sci* 30:28-35.
- Kwon YI, Vatter DA, Shetty K. 2006. Evaluation of clonal herbs of Lamiaceae species for management of diabetes and hypertension. *Asia Pac J Clin Nutr* 15:107-118.
- Lee KP, Sudjarwo GW, Kim JS, Dirgantara S, Maeng WJ, Hong H. 2014. The anti-inflammatory effect of Indonesian *Areca*

- catechu* leaf extract *in vitro* and *in vivo*. Nutr Res Pract 8:267-271.
- Li HX, Han SY, Wang XW, Ma X, Zhang K, Wang L, Ma ZZ, Tu PF. 2009. Effect of the carthamins yellow from *Carthamus tinctorius* L. on hemorheological disorders of blood stasis in rats. Food Chem Toxicol 47:1797-1802.
- Li X, Gao Z, Li T, Sarker SK, Chowdhury S, Jiang Z, Mu Z. 2018. Effects of pH values on physicochemical properties and antioxidant potential of whey protein isolated-safflower yellow complexes. Food Sci Technol Res 24:475-484.
- Lim ES. 2018. Preparation and functional properties of probiotic and oat-based synbiotic yogurts fermented with lactic acid bacteria. Appl Biol Chem 61:25-37.
- Ma Z, Li C, Qiao Y, Lu C, Li J, Song W, Sun J, Zhai X, Niu J, Ren Q, Wen A. 2016. Safflower yellow B suppresses HepG2 cell injury induced by oxidative stress through the AKT/Nrf2 pathway. Int J Mol Med 37:603-612.
- Mhatre SV, Bhagit AA, Yadav RP. 2019. Proteinaceous pancreatic lipase inhibitor from the seed of *Litchi chinensis*. Food Technol Biotechnol 57:113-118.
- Omari NE, Sayah K, Fettach S, Blidi OE, Bouyahya A, Faouzi MEA, Kamal R, Barkiyou M. 2019. Evaluation of *in vitro* antioxidant and antidiabetic activities of *Aristolochia longa* extracts. Evid Based Complement Alternat Med 2019: 7384735.
- Pires TC, Dias MI, Barros L, Barreira JC, Santos-Buelga C, Ferreira IC. 2018. Incorporation of natural colorants obtained from edible flowers in yogurts. LWT-Food Sci Technol 97:668-675.
- Sabu M, Kuttan R. 2002. Anti-diabetic activity of medicinal plants and its relationship with their antioxidant property. J Ethnopharmacol 81:155-160.
- Sung JM, Kim YB, Kum JS, Choi YS, Seo DH, Choi HW, Park JD. 2015. Effects of freeze-dried mulberry on antioxidant activities and fermented characteristics of yogurt during refrigerated storage. Korean J Food Sci Anim Resour 35:807-814.
- Tian J, Li G, Liu Z, Fu F. 2008. Hydroxysafflor yellow A inhibits rat brain mitochondrial permeability transition pores by a free radical scavenging action. Pharmacology 82:121-126.
- Tseng A, Zhao Y. 2013. Wine grape pomace as antioxidant dietary fibre for enhancing nutritional value and improving storability of yogurt and salad dressing. Food Chem 138:356-365.
- Wei X, Luo M, Xu L, Zhang Y, Lin X, Kong P, Liu H. 2011. Production of fibrinolytic enzyme from *Bacillus amyloliquefaciens* by fermentation of chickpeas, with the evaluation of the anticoagulant and antioxidant properties of chickpeas. J Agric Food Chem 59:3957-3963.
- Yoshioka Y, Yoshimura N, Matsumura S, Wada H, Hoshino M, Makino S, Morimoto M. 2019. α -Glucosidase and pancreatic lipase inhibitory activities of diterpenes from ndian mango ginger (*Curcuma amada* Roxb.) and its derivatives. Molecules 24:4071.
- Zhao Y, Sun H, Li X, Zha Y, Hou W. 2018. Hydroxysafflor yellow A attenuates high glucose-induced pancreatic β -cells oxidative damage via inhibiting JNK/c-jun signaling pathway. Biochem Biophys Res Commun 505:353-359.