









Original Article  
Microbiology



# Colistin resistance and plasmid-mediated *mcr* genes in *Escherichia coli* and *Salmonella* isolated from pigs, pig carcass and pork in Thailand, Lao PDR and Cambodia border provinces

Chanika Pungpian <sup>1</sup>, Scarlett Lee <sup>2</sup>, Suthathip Trongjit <sup>1</sup>, Nuananong Sinwat <sup>3</sup>, Sunpetch Angkititrakul <sup>4</sup>, Rangsiya Prathan <sup>1</sup>, Songsak Srisanga <sup>1</sup>, Rungtip Chuanchuen <sup>1,\*</sup>

<sup>1</sup>Research Unit in Microbial Food Safety and Antimicrobial Resistance, Department of Veterinary Public Health, Faculty of Veterinary Science, Chulalongkorn University, Bangkok, Thailand

<sup>2</sup>Department of Microbiology and Immunology, College of Veterinary Medicine, Cornell University, Ithaca, New York, United States

<sup>3</sup>Department of Farm Resources and Production Medicine, Faculty of Veterinary Medicine Kasetsart University, Kamphangsae Nakornpathom, Thailand

<sup>4</sup>Department of Veterinary Public Health, Faculty of Veterinary Medicine, Khon Kaen University, Khon Kaen, Thailand

 OPEN ACCESS

Received: Apr 10, 2021

Revised: Jun 20, 2021

Accepted: Jul 5, 2021

Published online: Aug 3, 2021

\*Corresponding author:

Rungtip Chuanchuen

Faculty of Veterinary Science, Chulalongkorn University, Pathumwan, Bangkok 10330, Thailand.

E-mail: chuanchuen.r@gmail.com


© 2021 The Korean Society of Veterinary Science

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.


ORCID iDs

Chanika Pungpian 

<https://orcid.org/0000-0002-0335-6761>

Scarlett Lee 

<https://orcid.org/0000-0002-0486-5621>

Suthathip Trongjit 

<https://orcid.org/0000-0003-3346-6299>

Nuananong Sinwat 

<https://orcid.org/0000-0003-3445-4206>

<https://vetsci.org>

## ABSTRACT

**Background:** Colistin and carbapenem-resistant bacteria have emerged and become a serious public health concern, but their epidemiological data is still limited.

**Objectives:** This study examined colistin and carbapenem resistance in *Escherichia coli* and *Salmonella* from pigs, pig carcasses, and pork in Thailand, Lao PDR, and Cambodia border provinces.

**Methods:** The phenotypic and genotypic resistance to colistin and meropenem was determined in *E. coli* and *Salmonella* obtained from pigs, pig carcasses, and pork (n = 1,619). A conjugative experiment was performed in all isolates carrying the *mcr* gene (s) (n = 68). The plasmid replicon type was determined in the isolates carrying a conjugative plasmid with *mcr* by PCR-based replicon typing (n = 7). The genetic relatedness of *mcr*-positive *Salmonella* (n = 11) was investigated by multi-locus sequence typing.

**Results:** Colistin resistance was more common in *E. coli* (8%) than *Salmonella* (1%). The highest resistance rate was found in *E. coli* (17.8%) and *Salmonella* (1.7%) from Cambodia. Colistin-resistance genes, *mcr-1*, *mcr-3*, and *mcr-5*, were identified, of which *mcr-1* and *mcr-3* were predominant in *E. coli* (5.8%) and *Salmonella* (1.7%), respectively. The *mcr-5* gene was observed in *E. coli* from pork in Cambodia. Two colistin-susceptible pig isolates from Thailand carried both *mcr-1* and *mcr-3*. Seven *E. coli* and *Salmonella* isolates contained *mcr-1* or *mcr-3* associated with the IncF and IncI plasmids. The *mcr*-positive *Salmonella* from Thailand and Cambodia were categorized into two clusters with 94%–97% similarity. None of these clusters was meropenem resistant.

**Conclusions:** Colistin-resistant *E. coli* and *Salmonella* were distributed in pigs, pig carcasses, and pork in the border areas. Undivided-One Health collaboration is needed to address the issue.

**Keywords:** Drug resistance; swine; Thailand; Laos; Cambodia

Sunpetch Angkitittrakul   
<https://orcid.org/0000-0002-1701-907X>  
 Rangsiya Prathan   
<https://orcid.org/0000-0003-1242-7493>  
 Songsak Srisanga   
<https://orcid.org/0000-0002-1710-7424>  
 Rungtip Chuanchuen   
<https://orcid.org/0000-0002-9714-9199>

#### Funding

This work was supported by TRF Basic Research Grant BRG6080014 co-funded by Thailand Research Fund (TRF), the Faculty of Veterinary Science, and Chulalongkorn University. The work was also partially supported by the 90<sup>th</sup> anniversary-Ratchadaphiseksomphot Endowment Fund and Chulalongkorn University-the One Health Research Cluster. CP is a recipient of a RGJ Ph.D. program PHD/0054/2558, co-funded by the TRF and Chulalongkorn University

#### Conflict of Interest

The authors declare no conflicts of interest.

#### Author Contributions

Conceptualization: Chuanchuen R, Angkitittrakul S, Pungpian C; Data curation: Pungpian C, Lee S, Trongjit S, Sinwat N; Formal analysis: Chuanchuen R, Pungpian C; Funding acquisition: Chuanchuen R; Investigation: Chuanchuen R, Angkitittrakul S, Pungpian C, Lee S, Trongjit S, Sinwat N, Prathan R, Srisanga S; Methodology: Chuanchuen R, Angkitittrakul S, Pungpian C, Lee S, Trongjit S, Sinwat N, Prathan R, Srisanga S; Project administration: Chuanchuen R; Resources: Chuanchuen R, Angkitittrakul S; Software: Chuanchuen R, Pungpian C; Supervision: Chuanchuen R; Validation: Chuanchuen R, Angkitittrakul S, Pungpian C, Lee S, Trongjit S, Sinwat N, Prathan R, Srisanga S; Visualization: Chuanchuen R; Writing - original draft: Chuanchuen R, Pungpian C; Writing - review & editing: Chuanchuen R, Angkitittrakul S, Pungpian C, Lee S, Trongjit S, Sinwat N, Prathan R, Srisanga S.

## INTRODUCTION

Antimicrobial resistance (AMR) is an ongoing One Health problem of global dimensions. The emergence and rapid spread of multidrug resistance (MDR) have complicated the AMR situation. Last-line antibiotics, e.g., polymyxins and carbapenems, have been used to treat infections involved in MDR Gram-negative bacterial pathogens. On the other hand, resistance to last-resort treatments has been increasingly reported [1], which has raised the alarm over the potentially limited treatment options in the near future.

Carbapenems are critically essential antimicrobials used to treat infections caused by MDR, including ESBL-producing Enterobacteriaceae [1]. Carbapenem-resistant *Enterobacteriaceae* (CRE) infections with high mortality have increased rapidly in hospitals [2], and CRE has also been observed in food animals [1]. Carbapenem resistance is commonly mediated by carbapenemase genes (e.g., *bla*<sub>KPC</sub>, *bla*<sub>NDM</sub>, *bla*<sub>VIM</sub>, *bla*<sub>IMP</sub>, and *bla*<sub>OXA</sub>), which are located on large conjugative plasmids carrying other resistance genes [3]. The coexistence of carbapenem and colistin resistance genes was reported in *E. coli* in patients and animals [1,4].

Colistin or polymyxin B is a last-resort antibiotic against MDR infections, especially CRE in humans [4]. The drug is used widely in pigs to treat intestinal infections caused by *Escherichia coli*, particularly post-weaning diarrhea [5]. Colistin resistance has been identified in pathogens from various sources, e.g., *E. coli* from pigs in China [5]; poultry in Argentina [6]; patients, pigs, and poultry in Cambodia [2]; and *Salmonella* from patients in Denmark [7]. Since the report of plasmid-mediated *mcr-1* in *E. coli* and *Klebsiella pneumoniae* from animals and patients in China in 2016 [8], several *mcr* variants have been discovered [9]. Currently, the detection of plasmid-mediated colistin resistance is included in various national AMR monitoring and surveillance programs globally [10].

The coexistence of *bla*<sub>NDM-1</sub> and *mcr-1* was detected in *E. coli* from patients in China and Taiwan and CRE from broilers and pigs in China [1]. The cotransfer of carbapenem and colistin resistance genes located on different plasmids has been demonstrated [3]. These could contribute to the emergence and spread of superbugs no longer susceptible to any commercially available antibiotic.

Pig and pork production is one of the largest markets worldwide, and several antimicrobials are used to improve the quantity and quality of products [11]. This phenomenon also occurs in Southeast Asia border trade areas, where imports and exports of live animals and products are an important contributor to increased cross-border trade. As a borderless One Health threat, AMR in one region may pose a risk to public health, animal health, and environmental health within or outside the region. AMR monitoring and surveillance is the foundation for assessing the burden of AMR and providing information for strategic action to control and prevent threats and evaluate the strategic interventions. Many developing countries still lack national AMR surveillance. Epidemiological data on AMR and data on last-line antibiotics are insufficient, particularly in these areas. Therefore, this study examined the prevalence and genetics underlying the resistance to carbapenems and colistin in *E. coli* and *Salmonella* isolated from pigs, pig carcasses, and pork in Thailand, Lao PDR, and Cambodia border provinces.

## MATERIALS AND METHODS

### Sample collection and bacterial isolation

A total of 988 *E. coli* and 631 *Salmonella* isolates from pigs (n = 377 and 139), pig carcasses (n = 328 and 174), and pork (n = 283 and 318) were included. They were isolated previously as part of AMR epidemiological studies in food animals and products in Southeast Asia between 2014 and 2018 [12-15]. The *E. coli* and *Salmonella* isolates were obtained from Thailand (n = 508 and 276), Lao PDR (n = 351 and 237), and Cambodia (n = 129 and 118) (Table 1). The sampling sites were located in the provinces with the two largest border-crossing points between Thailand-Lao PDR (i.e., Nong Khai-Vientiane and Mukdahan-Savannakhet) and Thailand-Cambodia (i.e., Sa Kaeo-Banteay Meanchey).

In the Thailand-Lao PDR border areas, the sampling sites included one municipal pig slaughterhouse and one municipal fresh market in each province. The slaughterhouses have large-scale facilities with a daily throughput of 50–200 or more pigs, and the municipal markets are located in the same area. In the Thailand-Cambodia border provinces, the samples were collected from three private pig slaughterhouses and one municipal fresh market in Sa Kaeo. The pig slaughterhouses are small-scale facilities producing pork for local consumption only. For the sampling sites in Cambodia, the samples were obtained from one municipal pig slaughterhouse and one municipal fresh market in Banteay Meanchey.

The rectal swabs were collected from the dead pigs immediately after bleeding but before the scalding process at the slaughterhouses. The pig carcass samples were obtained by swabbing the inner side of a half pig carcass from the neck to the bottom after the first whole carcass was cut lengthwise and divided into two halves. The pork samples were collected by swabbing at least 50 cm<sup>2</sup> of raw meat at the retail fresh markets. Sample collection at the slaughterhouses and retail markets was performed by the authors' sample collection team consisting of laboratory staff and veterinarians trained in the same sampling protocol [16]. All samples collected were stored in an icebox and transported to the laboratory within 24 h after sampling.

**Table 1.** Source and number of *E. coli* and *Salmonella* isolates (n = 1,619)

Bacteria	Country	Provinces	Sampling location	Sample type	No. of isolates	Total
<i>E. coli</i> (n = 988)	Thailand (n = 508)	Nong Khai, Mukdahan, Sa Kaeo	Slaughterhouse	Pig rectal swab	180	508
			Slaughterhouse	Pig carcass swab	185	
			Retail meat market	Pork	143	
	Lao PDR (n = 351)	Vientiane, Savannakhet	Slaughterhouse	Pig rectal swab	115	351
			Slaughterhouse	Pig carcass swab	132	
			Retail meat market	Pork	104	
	Cambodia (n = 129)	Banteay Meanchey, Siem Reap	Slaughterhouse	Pig rectal swab	82	129
			Slaughterhouse	Pig carcass swab	11	
			Retail meat market	Pork	36	
<i>Salmonella</i> (n = 631)	Thailand (n = 276)	Nong Khai, Mukdahan, Sa Kaeo	Slaughterhouse	Pig rectal swab	58	276
			Slaughterhouse	Pig carcass swab	63	
			Retail meat market	Pork	155	
	Lao PDR (n = 237)	Vientiane, Savannakhet	Slaughterhouse	Pig rectal swab	59	237
			Slaughterhouse	Pig carcass swab	82	
			Retail meat market	Pork	96	
	Cambodia (n = 118)	Banteay Meanchey, Siem Reap	Slaughterhouse	Pig rectal swab	22	118
			Slaughterhouse	Pig carcass swab	29	
			Retail meat market	Pork	67	
Grand total						1,619

The *E. coli* strains were isolated and confirmed biochemically [17]. The *E. coli* isolates were selected on MacConkey agar and Eosin Methylene Blue agar and confirmed by an indole test. One *E. coli* colony was collected from each positive sample. The *Salmonella* strains were isolated according to ISO6579:2002 (E) [18] and subjected to serotyping by slide agglutination [19]. A single colony of each serotype was collected from each positive sample. Forty-five serovars were included, of which the serovars Typhimurium and Rissen were most common (**Table 2**).

**Table 2.** *Salmonella* serovars from pig, pig carcass, and pork samples in Thailand, Lao PDR, and Cambodia border provinces in this study (n = 631)

<i>Salmonella</i> serovars	No. of isolates												Total
	Thailand				Lao PDR				Cambodia				
	Pig	Carcass	Pork	Sub total	Pig	Carcass	Pork	Sub total	Pig	Carcass	Pork	Sub total	
Afula					2			2					2
Agona							1	1					1
Anatum	3		15	18	9	25	16	50	1	1			70
Braenderup											1	1	1
Chincol											1	1	1
Corvallis			21	21									21
Derby			4	4	4	3	10	17	3	6		9	30
Dessau					2			2					2
Doncaster										1		1	1
Duesseldorf			4	4									4
Eindegi					1	6	2	9					9
Enteritidis	1			1						1		1	2
Fufu									1			1	1
Give	1	2	1	4	1		1	2					6
Havana							2	2					2
Hvittingfoss			2	2		1		1					3
Indians											1	1	1
Kedougou	11	4	11	26									26
Kentucky		1		1									1
Lexington									3		2	5	5
London						1		1					1
Mbandaka											1	1	1
Menden	1			1									1
Monschau			2	2									2
Mowanjum											1	1	1
Newmexico									2	1		3	3
Norwich										1		1	1
Panama		1	2	3	1			1					4
Paratyphi		1	1	2					3	3	13	19	21
Preston							2	2					2
Readind					1			1					1
Rissen	11	20	16	47	7	11	17	35	4		42	46	128
Saintpaul		6	7	13									13
Schleissheim										1	1	2	2
Schwarzengrund			1	1									1
Singapore										2	1	3	3
Stanley	1	6	9	16	11	10	13	34		1	1	2	52
Stuttgart			1	1									1
Tsevie							2	2					2
Typhimurium	27	16	52	95	15	22	19	56	3	8		11	162
Urbana			2	2									2
Wandsworth										1		1	1
Warragul									1	1		2	2
Welterreden	1	6	4	11	5	3	1	9	1			1	21
Worthington	1			1			10	10		1	2	3	14
Total	58	63	154	276	59	82	96	237	22	29	67	118	631

### Determination of colistin and meropenem susceptibility (n=1,619)

The minimum inhibitory concentrations (MICs) of colistin were determined using the agar dilution method [20]. The joint CLSI-EUCAST Polymyxin Breakpoints Working Group recommended the broth microdilution method for determining the colistin MIC. On the other hand, the plastic binding feature of colistin and the synergistic effects of polysorbate 80 with colistin have been observed [21]. The agar dilution method generates a comparable colistin MIC value to the broth microdilution method [21], but is superior in reproducibility, robustness, and user-friendliness. Therefore, the agar dilution method was used to determine the colistin MIC in this study. The colistin concentration range was 0.125–64 µg/mL with an MIC breakpoint of 2 µg/mL [22]. *E. coli* ATCC25922, *Staphylococcus aureus* ATCC29213, and *Pseudomonas aeruginosa* ATCC27853 served as the quality control strains.

Carbapenemase production was screened by the disk diffusion method using meropenem disks (10 µg) [20] and confirmed using the modified Hodge test (MHT) with meropenem (10 µg). *E. coli* ATCC25922 was the control strain.

### PCR and DNA sequencing analysis

The total DNA prepared by whole-cell boiling was used as a PCR template [23]. **Table 3** lists all the primers used. The PCR products were purified using Nucleospin Gel and PCR clean up (Macherey-Nagel, Germany) and submitted for DNA sequencing using the PCR primers (First Base Laboratories, Selangor Darul Ehsan, Malaysia). The nucleotide sequences were analyzed by a comparison to the published sequences available at the NCBI website.

All bacterial isolates (n = 1,619) were screened for colistin-resistance encoding genes, including *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4*, and *mcr-5* using multiplex PCR with the specific primers [24]. Representatives of PCR amplicons were purified and submitted for DNA sequencing to confirm the specificity of PCR amplification. The DNA sequences obtained were deposited

**Table 3.** Primers used in this study

Target	Primer	Primer sequenc (5'-3')	Product size (bp)	Reference
<b>MCR</b>				
Multiplex	<i>mcr1</i> -F	AGTCCGTTTGTCTTGTGGC	320	[24]
	<i>mcr1</i> -R	AGATCCTTGGTCTCGGCTTG		
	<i>mcr2</i> -F	CAAGTGTGTTGGTCGCAGTT	715	[24]
	<i>mcr2</i> -R	TCTAGCCCGACAAGCATAACC		
	<i>mcr3</i> -F	AAATAAAAATTGTTCCGCTTATG	929	[24]
	<i>mcr3</i> -R	AATGGAGATCCCCGTTTTT		
	<i>mcr4</i> -F	TCACTTTCATCACTGCGTTG	1116	[24]
	<i>mcr4</i> -R	TTGGTCCATGACTACCAATG		
	<i>mcr5</i> -F	ATGCGGTTGTCTGCATTATC	1644	[24]
	<i>mcr5</i> -R	TCATTGTGGTTGTCTTTTCTG		
<b>Carbapenemase genes</b>				
Multiplex	<i>bla<sub>IMP</sub></i> -F	GGAATAGAGTGGCTTAAYTCTC	232	[25]
	<i>bla<sub>IMP</sub></i> -R	GGTTTAAAYAAAACAACCACC		
	<i>bla<sub>VIM</sub></i> -F	GATGGTGTGTTGGTCGCATA	390	[25]
	<i>bla<sub>VIM</sub></i> -R	CGAATGCGCAGCACCAG		
	<i>bla<sub>OXA</sub></i> -F	GCGTGGTTAAGGATGAACAC	477	[25]
	<i>bla<sub>OXA</sub></i> -R	CATCAAGTTCAACCCAACCG		
	<i>bla<sub>NDM</sub></i> -F	GGTTTGCGCATCTGGTTTTTC	621	[25]
	<i>bla<sub>NDM</sub></i> -R	CGGAATGGCTCATCACGATC		
	<i>bla<sub>KPC</sub></i> -F	CGTCTAGTTCTGCTGTCTTG	798	[25]
	<i>bla<sub>KPC</sub></i> -R	CTTGTCATCTTGTAGGCG		

(continued to the next page)

**Table 3.** (Continued) Primers used in this study

Target	Primer	Primer sequenc (5'-3')	Product size (bp)	Reference
<b>PBRT</b>				
Multiplex 1	H11-F	GGAGCGATGGATTACTTCAGTAC	471	[27]
	H11-R	TGCCGTTTCACCTCGTGAGTA		
	H12-F	TTTCTCCTGAGTCACCTGTTAACAC	644	[27]
	H12-R	GGCTCACTACCGTTGTCATCCT		
	I1-F	CGAAAAGCCGGACGGCAGAA	139	[27]
	I1-V	TCGTGTTCCGCAAGTTCGT		
Multiplex 2	X-F	AACCTTAGAGGCTATTTAAGTTGCTGAT	376	[27]
	X-R	TGAGAGTCAATTTTATCTCATGTTTATAGC		
	L/M-F	GGATGAAAATATCAGCATCTGAAG	785	[27]
	L/M-R	CTGCAGGGGCGATTCTTTAGG		
	N-F	GTCTAACGAGCTTACCGAAG	559	[27]
	N-R	GTTTCAACTCTGCCAAGTTC		
	Multiplex 3	FIA-F	CCATGCTGGTCTAGAGAAGGTG	462
FIA-R		GTATATCCTTACTGGCTTCCGCAG		
FIB-F		GGAGTTCTGACACACGATTTCTG	702	[27]
FIB-R		CTCCCGTCTCAGGGCATT		
W-F		CCTAAGAACAACAAGCCCCCG	242	[27]
	W-R	GGTGCGCGGCATAGAACCCT		
	Multiplex 4	Y-F	AATTCAAACAACACTGTGCAGCCTG	765
Y-R		GCGAGAATGGACGATTACAAAACCTT		
P-F		CTATGGCCCTGCAAACGCGCCAGAAA	534	[27]
P-R		TCACGCGCCAGGGCGCAGCC		
FIC-F		GTGAACTGGCAGATGAGGAAGG	262	[27]
	FIC-R	TTCTCCTCGTCGCCAAACTAGAT		
	Multiplex 5	A/C-F	GAGAACCAAAGACAAGACCTGGA	465
A/C-R		ACGACAAACCTGAATTGCCTCCTT		
T-F		TTGGCCTGTTGTGCCTAAACCAT	750	[27]
T-R		CGTTGATTACACTTAGCTTTGGAC		
FIIS-F		CTGTCGTAAGCTGATGGC	270	[27]
	FIIS-R	CTTGCCACAACTTCAGC		
	Simplex 1	FrepB-F	TGATCGTTTAAGGAATTTTG	270
FrepB-R		GAAGATCAGTCACACCATCC		
Simplex 2	K-F	GCGGTCCGAAAGCCAGAAAAC	160	[27]
	K-R	TCTTTCACGAGCCCGCAAA		
Simplex 3	B/O-F	GCGGTCCGAAAGCCAGAAAAC	159	[27]
	B/O-R	TCTGCGTCCGCAAGTTCGA		
MLST	<i>fimA</i> -F	TCAGGGGAGAAAACAGAAAATAAT	760	[28]
	<i>fimA</i> -R	TCCCGATAGCCTCTTCC		
	<i>manB</i> -F	CCGGCACCCGAAGAGA	893	[28]
	<i>manB</i> -R	CGCCGCCATCCGGTC		
	<i>mdh</i> -F	ATGAAAGTCGAGTCTCGGCGTCTGGCGG	849	[28]
	<i>mdh</i> -R	ATATCTTYYTTCAGCGTATCCAGCAT		

PBRT, polymerase chain reaction-based replicon typing; MLST, multi-locus sequence typing.

in the GenBank database, of which the accession numbers are as follows: *mcr-1*, MW314264; *mcr-3*, MW314265; *mcr-5*, MW314266.

Carbapenem-resistance encoding genes were detected in all bacterial isolates using multiplex PCR with the specific primers [25], including *bla<sub>IMP</sub>*, *bla<sub>VIM</sub>*, *bla<sub>OXA</sub>*, *bla<sub>NDM-1</sub>*, and *bla<sub>KPC</sub>*.

### Conjugation experiments and plasmid replicon typing

A biparental mating experiment was performed in all *mcr*-positive *E. coli* and *Salmonella* isolates. All *mcr*-carrying *E. coli* (n = 57) were used as donors and spontaneous rifampicin-resistant *Salmonella* Enteritidis SE12 (SE12<sup>rif</sup>, MIC = 256 µg/mL) was used as the recipient [26]. The transconjugants were selected on LB agar containing rifampicin (32 µg/mL)



with colistin (2 µg/mL) and confirmed to be *Salmonella* on brilliant green agar and xylose lysine deoxycholate agar (Difco, MD, USA) containing 2 µg/mL colistin. For all *mcr*-positive *Salmonella* isolates (n=11), the spontaneous rifampicin-resistant *E. coli* K12 strain MG1655 (MG1655rif<sup>r</sup>) was used as the recipient [26]. The transconjugants were selected on LB agar containing rifampicin (32 µg/mL) with colistin (2 µg/mL) and confirmed to be *E. coli* on MacConkey agar and Eosin Methylene Blue agar (Difco, MD) containing 2 µg/mL colistin. The presence of the corresponding genes in all transconjugants was confirmed by PCR using specific primers as described above.

The plasmid Inc groups were determined in all the isolates containing the conjugative plasmid by a PCR-based replicon typing (PBRT) [27]. Eighteen plasmid Inc groups were detected using five multiplex PCR (i.e., HI1/HI2/I1-Iγ; X/L-M/N; FIA/FIB/W; Y/P/FIC and A-C/T/FIIA) and three simplex PCR (i.e., F, K, and B/O).

### Salmonella phylogenetic analysis

All *mcr*-positive *Salmonella* (n = 11) were characterized for the phylogenetic groups by multi-locus sequence typing (MLST). Three housekeeping genes, *fimA*, *manB*, and *mdh* were PCR amplified as described previously [28]. All PCR products were purified and submitted for nucleotide sequencing. The obtained sequences were assembled and proofread using the DNASTAR program [29] and aligned using the MUSCLE options in MEGA-X software [30]. A phylogenetic tree with 1000 bootstrap replicates was constructed based on the concatenated alignment of *fimA*, *manB*, and *mdh* sequences using the Neighbor-Joining (NJ) method [28]. The Maximum Composite Likelihood method was used to calculate the evolutionary distances [30]. All positions containing gaps and missing data were removed for each sequence pair. The final data set contained 18,129 positions. Evolutionary analyses were conducted using MEGA X [30].

### Statistical analysis

Statistical analysis was performed using the SPSS version 22.0 (IBM Corporation) program. A chi-squared test was used to compare the AMR phenotype and genotype. The statistical significance ( $p < 0.05$ ) of the differences between the AMR phenotypic and genotypic percentage was determined.

## RESULTS

### Phenotypic resistance to colistin and meropenem (n=1,619)

**Table 4** lists the colistin resistance rates of *E. coli* and *Salmonella* from Thailand, Lao PDR, and Cambodia. The prevalence of colistin-resistant *E. coli* was higher than that of *Salmonella* in all countries ( $p < 0.05$ ). Colistin resistance was commonly observed in either *E. coli* (17.8%;

**Table 4.** Colistin resistance rates of *E. coli* and *Salmonella* from pigs, pig carcasses, and pork in Thailand, Lao PDR, and Cambodia (n = 1,619)

Bacteria	Resistance rate*												Grand total
	Thailand				Lao PDR				Cambodia				
	Pig	Carcass	Pork	Total	Pig	Carcass	Pork	Total	Pig	Carcass	Pork	Total	
<i>E. coli</i>	9.4% (17/180)	5.4% (10/185)	7.7% (11/143)	7.5% <sup>†§</sup> (38/508)	7.8% (9/115)	6.1% (8/132)	1.9% (2/104)	5.4% <sup>†§</sup> (19/351)	22% (18/82)	9.1% (1/11)	11.1% (4/36)	17.8% <sup>†  </sup> (23/129)	8% (80/988)
<i>Salmonella</i>	3.4% (2/58)	0 (0)	0.6% (1/155)	1.1% <sup>†§</sup> (3/276)	0 (0)	1.2% (1/83)	0 (0)	0.5% <sup>†§</sup> (1/218)	4.5% (1/22)	0 (0)	1.5% (1/67)	1.7% <sup>†  </sup> (2/118)	1% (6/631)

\*In parenthesis, Number of colistin-resistant isolates/Number of bacterial isolates from each source; <sup>†</sup>+values bearing a different superscript in the same column are significantly different ( $p < 0.5$ ); <sup>§</sup>||values bearing a different superscript in the same row are significantly different ( $p < 0.5$ ).

23/129) or *Salmonella* (1.7%; 2/118) from Cambodia ( $p < 0.05$ ). When considering the sample sources, colistin-resistant *E. coli* were most common in pigs in Thailand (9.4%; 17/180), Lao PDR (7.8%; 9/115), and Cambodia (22%; 18/82). Colistin-resistant *Salmonella* was identified most frequently in pigs in Thailand (3.4%; 2/58) and Cambodia (4.5%; 1/22) (**Table 4**). None of the isolates in this collection was meropenem resistant.

### Distribution of *mcr* genes (n = 1,619)

Of the *E. coli* isolates (n = 988), 57 isolates (5.8%; 57/988) carried at least one *mcr* gene, including the isolates from Thailand (3.5%; 18/508), Lao PDR (6%; 21/351), and Cambodia (14%; 18/129) (**Table 5**). The percentage of *mcr*-carrying *E. coli* isolates in Cambodia was significantly higher than in Thailand and Lao PDR ( $p < 0.05$ ). The *mcr* genes were most common among the pig isolates (8.8%; 33/377), followed by the pig carcasses (4.3%; 14/328), and pork (3.5%; 10/283) isolates and corresponded to the colistin resistance phenotype. The percentage of *mcr*-carrying *E. coli* from pigs was significantly higher than pork ( $p < 0.05$ ). The *mcr-1* gene was predominant (56.1%; 32/57), followed by *mcr-3* (38.6%; 22/57), *mcr-1/mcr-3* (3.5%; 2/57), and *mcr-5* (1.8%; 1/57). The *mcr* genes were identified more commonly among *E. coli* (5.8%; 57/988) than the *Salmonella* (1.7%; 11/631) isolates ( $p < 0.05$ ). No *mcr-2* and *mcr-4* genes were detected.

**Table 5.** Presence and transfer of *mcr* genes in *E. coli* and *Salmonella* (n = 1,619)

Bacteria	Country	<i>mcr</i> -gene	Sample source	MIC ( $\mu\text{g/mL}$ )	No. of isolates (%)	Transferability*
<i>E. coli</i> (n = 988)	Thailand (n = 508)	<i>mcr-1</i>	Pig	8	4	-
			Pig carcass	0.5, 8	3	+ (1)
			Pork	4	1	+ (1)
		<i>mcr-3</i>	Pig	4-8	6	-
			Pig carcass	0.5	1	-
			Pork	8	1	+ (1)
		<i>mcr-1/ mcr-3</i>	Pig	1	2	-
		Subtotal			18 (4%) <sup>†</sup>	
		Lao PDR (n = 351)	<i>mcr-1</i>	Pig	8	3
	Pig carcass			4-8	6	-
	Pork			4-8	2	-
	<i>mcr-3</i>		Pig	4-8	5	+ (1)
			Pig carcass	2-8	3	-
	Pork		0.5-4	2	-	
	Subtotal				21 (6%) <sup>†</sup>	
	Cambodia (n = 129)	<i>mcr-1</i>	Pig	8-16	10	-
			Pig carcass	8	1	-
			Pork	4-8	2	-
		<i>mcr-3</i>	Pig	4	3	-
			Pork	0.5	1	-
		<i>mcr-5</i>	Pork	2	1	-
Subtotal			18 (14%) <sup>‡</sup>			
Grand total				57 (6%) <sup>§</sup>		
<i>Salmonella</i> (n = 631)	Thailand (n = 276)	<i>mcr-1</i>	Pig	1	1	-
			Pig carcass	1-8	3	+ (2)
		<i>mcr-3</i>	Pork	1-2	2	-
	Subtotal			6 (2.2%)		
	Cambodia (n = 118)	<i>mcr-3</i>	Pig	0.5-4	4	+ (1)
			Pork	4	1	-
	Subtotal			5 (4.2%)		
Grand total				11 (2.8%) <sup>  </sup>		

MIC, minimum inhibitory concentration.

<sup>†</sup>The number indicates the colistin-resistant isolate that can transfer *mcr*; <sup>‡</sup>Values for *E. coli* from different countries bearing a different superscript in the same column are significantly different ( $p \leq 0.05$ ); <sup>§</sup>Values for *E. coli* and *Salmonella* bearing a different superscript in the same column are statistical different ( $p \leq 0.05$ ).



**Table 6.** Characteristics of *E. coli* and *Salmonella* transconjugant (n = 7)

Bacteria	Country	Sample	MIC (µg/mL)	Bacterial donor		Bacterial recipient	
				<i>mcr</i> gene	Inc group	<i>mcr</i> gene	Inc group
<i>E. coli</i>	Thailand	Carcass	8	<i>mcr-1</i>	Incl, IncFIB, IncFrepB	<i>mcr-1</i>	Incl
<i>E. coli</i>	Thailand	Pork	4	<i>mcr-1</i>	IncFrepB	<i>mcr-1</i>	IncFrepB
<i>E. coli</i>	Thailand	Pork	8	<i>mcr-3</i>	IncY, IncFrepB	<i>mcr-3</i>	IncFrepB
<i>E. coli</i>	Lao PDR	Pig	8	<i>mcr-3</i>	IncFrepB	<i>mcr-3</i>	IncFrepB
<i>Salmonella</i>	Thailand	Pig	8	<i>mcr-3</i>	Incl, IncFIB	<i>mcr-3</i>	Incl
<i>Salmonella</i>	Thailand	Pig	8	<i>mcr-3</i>	Incl, IncFIB	<i>mcr-3</i>	Incl
<i>Salmonella</i>	Cambodia	Pig	4	<i>mcr-3</i>	IncFrepB	<i>mcr-3</i>	IncFrepB

MIC, minimum inhibitory concentration.

The co-occurrence of *mcr-1/mcr-3* was detected in two colistin-susceptible isolates (colistin MIC = 1 µg/mL) from pigs in Thailand. To the best of the authors' knowledge, this is the first report of *mcr-5* in pork isolates in Cambodia (colistin MIC=2 µg/mL).

In *Salmonella* (n = 631), 11 isolates (1.7%; 11/631) were positive to at least one *mcr* gene tested, including six Thai (2.2%; 6/276) and five Cambodian (4.2%; 5/118) isolates. The *mcr* genes were more common in the pig isolates than the pork isolates ( $p < 0.05$ ). The *mcr-3* gene (90.9%; 10/11) was most common, followed by *mcr-1* (9.1%; 1/11). None was positive to *mcr-2*, *mcr-4*, or *mcr-5*.

None of the *E. coli* and *Salmonella* isolates were positive to the carbapenem-resistance genes tested.

### Horizontal transfer and plasmid replicon typing of *mcr* gene carrying isolates


Two *E. coli* isolates from pig carcass and pork in Thailand were *mcr-1* transferred conjugally to *Salmonella* recipients. The *mcr-3* gene in two *E. coli* isolates (one from Thai pork and the others from a Laos pig) and three *Salmonella* (two from Thai pigs and one from Cambodian pig) were transferred horizontally. All the isolates were resistant to colistin (colistin MIC = 4–8 µg/mL) and harbored the IncF plasmid. The *mcr*-conjugative plasmids were in the IncF and Incl groups (Table 6).

### Evolutionary relationship among *mcr*-carrying *Salmonella*

The NJ phylogenetic tree showed that *mcr*-carrying *Salmonella* isolates (n = 11) could be classified into two clusters, Clusters A and B (Table 7). The percentage similarity within each cluster ranged from 94% to 97%, and the average branch lengths of the *Salmonella* isolates within the same cluster ranged from 0.00 to 0.01 (Fig. 1). The phylogenetic cluster A (n = 7) was predominant, including four isolates from Thailand [*S. Typhimurium* from pigs (n = 2), *S. Kedougou* from pork (n = 1), and *S. Rissen* from a pig (n = 1)] and three isolates from Cambodia [*S. Paratyphi* from pigs (n = 2) and *S. Kentucky* from a pig (n = 1)]. Cluster B

**Table 7.** Cluster and characteristics of selected *Salmonella* in multi-locus sequence typing (n = 11)

Cluster	<i>Salmonella</i>	Country	Source	<i>mcr</i> gene
	Serovars (n = 11)			
A	Paratyphi (n = 2)	Cambodia	Pig	<i>mcr 3</i>
	Kentucky (n = 1)	Cambodia	Pig	<i>mcr 1</i>
	Typhimurium (n = 2)	Thailand	Pig	<i>mcr 3</i>
	Kedougou (n = 1)	Thailand	Pork	<i>mcr 3</i>
	Rissen (n = 1)	Thailand	Pig	<i>mcr 1</i>
B	Typhimurium (n = 1)	Cambodia	Pig	<i>mcr 3</i>
	Typhimurium (n = 1)	Cambodia	Pork	<i>mcr 3</i>
	Huettwillen (n = 1)	Thailand	Pig	<i>mcr 3</i>
	Rissen (n = 1)	Thailand	Pork	<i>mcr 3</i>



**Fig. 1.** Phylogenetic tree of the *Salmonella* isolates (n = 11). Phylogenetic analysis was based on the alignment of the three multi-locus sequence typing genes (*manB*, *mdh*, and *fimA*) sequenced. The sequence was aligned by the MUSCLE options in MEGA-X, and a phylogenetic tree was constructed using the neighbor-joining method. The tree was drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree.

contained four *Salmonella* isolates, including three isolates from Thailand [*S. Typhimurium* from a pig (n = 1), *S. Huettwillen* from a pig (n = 1), and *S. Rissen* from pork (n = 1)], and one isolate from Cambodia [*S. Typhimurium* from pork (n = 1)].

## DISCUSSION

One of the major findings of this study was the presence of colistin-resistant *E. coli* and *Salmonella* harboring *mcr* in pigs, pig carcasses, and pork, indicating the circulation of colistin-resistant bacteria and genes in the pig production chain in the region. A previous study reported that flattening pigs entering slaughterhouses acted as vehicles for colistin-resistant bacteria distribution to the slaughterhouse environment [31]. This agrees with the observation of a high percentage of colistin-resistant *E. coli* in pigs in this study. Pig carcasses might be contaminated with resistant bacteria during the slaughtering process that is transferred to the pork in the retail meat markets [32]. The resistant bacteria may spread to workers at pig farms or slaughterhouses and butchers at the retail markets by direct contact and the food chain. These results show that the circulation of colistin-resistant bacteria in the pig and pork production chain is a potential route for introducing resistant bacteria to humans and the environment [32].

For Thailand, the colistin-resistant *E. coli* (7.5%) in this study was more common than a previous study reporting the absence of colistin-resistant *E. coli* in pig farms in other provinces of Northeastern Thailand [11]. Although the extent of antimicrobial use is largely unclear, the results may reflect the increase in colistin consumption in pig production in the area [31]. This discrepancy might be associated with diverse sources and different geographical origins of the samples as well. Furthermore, colistin resistance could be the result of co-selection caused by other antimicrobials. A previous study in *E. coli* showed that host cationic antimicrobial peptides (e.g., LL-37 and lysozyme) could promote cross-resistance to polymyxins, possibly explaining the spread of colistin resistance in the limited colistin use [33]. In contrast, the percentage of colistin resistance in Cambodia (17.8%) was

significantly higher than that in Thailand (7.5%) and Lao PDR (5.4%), but was slightly lower than a previous study in Cambodia [34]. This may be because of the higher use of colistin and other antimicrobial drugs in the country, which agrees with a previous study showing that the greater use and longer application of antimicrobials in pig feed on farms may increase the AMR prevalence in Cambodia [34].

The phenotypic colistin resistance and *mcr* in *E. coli* are more common than that in *Salmonella* ( $p < 0.05$ ). A previous study reported that the resistance rates to antimicrobials in different classes in *E. coli* were frequently higher than those in *Salmonella* [35]. *E. coli* are commensals existing in the intestinal tract of all animals and have a longer contact duration to antimicrobials administered to animals than *Salmonella*. Therefore, they have greater effectiveness in the development of resistance to antimicrobials than *Salmonella* [36]. In addition, AMR development in *E. coli* is affected more by the antimicrobials used than in *Salmonella* under the same condition [36].

The *mcr-1* and *mcr-3* genes were predominant *mcr* variants in this study. The *mcr-1* gene was most common among colistin-resistant *E. coli* (59.6%), which agrees with a previous study in pigs in China [31]. This gene has been detected in *E. coli* and *Salmonella* from humans and livestock in most parts of the world, including Asia (e.g., China) [32], Europe (e.g., Spain) [9], America (e.g., Argentina) [6] and South Africa (e.g., Algeria) [37]. The *mcr-1* gene was located on a conjugative plasmid, supporting its wide presence in *E. coli* and *Salmonella*. On the other hand, it was previously reported that *mcr-1* is stable and can be transferred without selective pressure from colistin [33]. The *mcr-3* gene was found in the *E. coli* and *Salmonella* isolates from pigs, pig carcasses, and pork in Thailand and Cambodia but was predominant among colistin-resistant *Salmonella* (10/11). Previous studies reported *mcr-3* in *Salmonella* in patients in Canada [38] and Denmark [7] who traveled to Thailand and *E. coli* and *Klebsiella pneumoniae* in patients in Thailand [39]. Plasmids carrying *mcr-3* were previously observed in *E. coli* from pigs in China and Malaysia, of which its sequence was similar to that in *K. pneumoniae* patient isolates in Thailand [40]. These data indicate the worldwide spread of *mcr-3* in humans and animals. An *E. coli* isolate from pork in Cambodia carried *mcr-5*. The *mcr-5* gene was previously detected in *E. coli* from pigs and poultry in China and Spain [9], suggesting the wide distribution of *mcr-5*. To the best of the authors' knowledge, this is the first report of *mcr-5* in *E. coli* from pork in Cambodia.

The coexistence of *mcr-1* and *mcr-3* was observed in *E. coli* from pigs in Thailand, which concurs with previous studies in Taiwan and China [41]. Almost all *E. coli* and *Salmonella* isolates carrying individual *mcr-1* or *mcr-3* were resistant to colistin (MIC $\geq$ 2  $\mu$ g/mL), suggesting that the presence of *mcr* in these isolates corresponded well to the colistin resistance phenotype. In contrast, the *E. coli* isolates carrying *mcr-1/mcr-3* were susceptible to colistin (MIC = 1  $\mu$ g/mL), which is in agreement with the study in China [41], supporting the premise that the coexistence of *mcr* in a single *E. coli* isolate does not generate a cumulative effect on colistin susceptibility [41]. These results indicate that the contribution of *mcr* to the colistin resistance phenotype varies and that the *mcr* genes should be screened in either colistin-susceptible or resistant isolates.

In this study, transconjugants ( $n = 7$ ) harbored the *mcr* gene on the IncF ( $n = 4$ ) or IncI ( $n = 3$ ) plasmids, which is in agreement with a previous study demonstrating IncF, IncI, IncHI2, and IncX4 plasmids carrying the *mcr-1* and *mcr-3* [42]. The IncF plasmid is associated with several

resistance genes worldwide. Overall, these observations highlight the potential distribution of *mcr* genes globally.

The prevalence of *mcr* genes in *E. coli* and *Salmonella* from Cambodia was significantly higher than in Thailand and Lao PDR, which agrees with the colistin resistance phenotype observed and suggests that the colistin resistance phenotype was being driven by the presence of the *mcr* genes [1]. Such higher prevalence may be associated with the use of colistin and antimicrobial drugs in other classes on-farm, resulting in the preservation and transfer of the genes [11].

Infection with *mcr*-carrying *Salmonella* may be difficult to treat and result in treatment failure. Knowledge of the genetic relatedness of *mcr*-harboring *Salmonella* will increase the understanding of their transmission mode [28]. In this study, all *mcr*-positive *Salmonella* were categorized into two clusters by MLST. Each cluster contained the pig and pork isolates from Thailand and Cambodia. The close similarity within the same cluster suggests the clonal expansion of *mcr*-positive *Salmonella* between the two countries. The *mcr*-positive *Salmonella* in Thailand and Cambodia were categorized into two clusters, possibly due to the multiple and independent acquisitions of *mcr* in bacteria in the same geographical area. This is similar to a study in China where *mcr-1* was harbored by multiple clones of *E. coli* isolates (e.g., ST10, ST101, and ST410) from hospital sewage water [43]. *E. coli* ST624 with *mcr-1* isolates from a patient in South Africa was identified in *E. coli* from poultry in Spain and Japan [44]. Overall, these observations suggest that *mcr-1* has circulated among different bacterial clones within and across countries, which may occur by diverse independent acquisitions of *mcr*-carrying plasmids [43].

CRE has spread globally and was previously reported in patients in the study areas [2]. In contrast, all *E. coli* isolates in this study were susceptible to meropenem and were not positive to any carbapenem resistance gene tested. Carbapenems are not approved for use in livestock in any country, resulting in minimized selective pressure conditions [3]. Simultaneously, carbapenem use is uncommon in livestock production in the region, due mainly to their high cost.

In summary, colistin resistance was detected in *E. coli* and *Salmonella* isolated from pigs, pig carcasses, and pork in Thailand, Lao PDR, and Cambodia border areas. The *mcr-1*, *mcr-3*, and *mcr-5* genes were found. Moreover, *mcr-1* and *mcr-3* co-occurred in the same *E. coli* isolates from pigs. The *mcr-1* and *mcr-3* genes were transferred horizontally, while resistance to meropenem was absent. Monitoring and surveillance of the resistance to last-line antibiotics at the phenotypic and genotypic levels should be encouraged as a collaboration between countries. Further study to characterize the genetic structure of *mcr*-carrying plasmids is currently underway.

## ACKNOWLEDGEMENTS

The authors wish to thank Dr. Tanittha Chatsuwana, department of microbiology, Chulalongkorn University, for providing control strains for *bla*<sub>KPC</sub>, *bla*<sub>NDM-1</sub>, *bla*<sub>OXA-1</sub>, *bla*<sub>IMP</sub>, and *bla*<sub>VIM</sub>.

## REFERENCES

1. Peng Z, Li X, Hu Z, Li Z, Lv Y, Lei M, et al. Characteristics of carbapenem-resistant and colistin-resistant *Escherichia coli* co-producing NDM-1 and MCR-1 from pig farms in China. *Microorganisms*. 2019;7(11):482. [PUBMED](#) | [CROSSREF](#)
2. Atterby C, Osbjørk K, Tepper V, Rajala E, Hernandez J, Seng S, et al. Carriage of carbapenemase- and extended-spectrum cephalosporinase-producing *Escherichia coli* and *Klebsiella pneumoniae* in humans and livestock in rural Cambodia; gender and age differences and detection of bla<sub>OXA-48</sub> in humans. *Zoonoses Public Health*. 2019;66(6):603-617. [PUBMED](#) | [CROSSREF](#)
3. Long H, Feng Y, Ma K, Liu L, McNally A, Zong Z. The co-transfer of plasmid-borne colistin-resistant genes *mcr-1* and *mcr-3.5*, the carbapenemase gene bla<sub>NDM-5</sub> and the 16S methylase gene *rmtB* from *Escherichia coli*. *Sci Rep*. 2019;9(1):696. [PUBMED](#) | [CROSSREF](#)
4. Liu L, Feng Y, Zhang X, McNally A, Zong Z. New variant of *mcr-3* in an extensively drug-resistant *Escherichia coli* clinical isolate carrying *mcr-1* and bla<sub>NDM-5</sub>. *Antimicrob Agents Chemother*. 2017;61(12):e01757-17. [PUBMED](#) | [CROSSREF](#)
5. Wang Z, Fu Y, Schwarz S, Yin W, Walsh TR, Zhou Y, et al. Genetic environment of colistin resistance genes *mcr-1* and *mcr-3* in *Escherichia coli* from one pig farm in China. *Vet Microbiol*. 2019;230:56-61. [PUBMED](#) | [CROSSREF](#)
6. Dominguez JE, Redondo LM, Figueroa Espinosa RA, Cejas D, Gutkind GO, Chacana PA, et al. Simultaneous carriage of *mcr-1* and other antimicrobial resistance determinants in *Escherichia coli* from poultry. *Front Microbiol*. 2018;9:1679. [PUBMED](#) | [CROSSREF](#)
7. Litrup E, Kiil K, Hammerum AM, Roer L, Nielsen EM, Torpdahl M. Plasmid-borne colistin resistance gene *mcr-3* in *Salmonella* isolates from human infections, Denmark, 2009-17. *Euro Surveill*. 2017;22(31):30587. [PUBMED](#) | [CROSSREF](#)
8. Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis*. 2016;16(2):161-168. [PUBMED](#) | [CROSSREF](#)
9. García-Meniño I, Díaz-Jiménez D, García V, de Toro M, Flament-Simon SC, Blanco J, et al. Genomic characterization of Prevalent *mcr-1*, *mcr-4*, and *mcr-5* *Escherichia coli* within swine enteric colibacillosis in Spain. *Front Microbiol*. 2019;10:2469. [PUBMED](#) | [CROSSREF](#)
10. European Food Safety Authority/European Centre for Disease Prevention and Control. The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2017. *EFSA J* 2019;17(2):e05598. [PUBMED](#)
11. Olaitan AO, Diene SM, Kempf M, Berrazeg M, Bakour S, Gupta SK, et al. Worldwide emergence of colistin resistance in *Klebsiella pneumoniae* from healthy humans and patients in Lao PDR, Thailand, Israel, Nigeria and France owing to inactivation of the PhoP/PhoQ regulator mgrB: an epidemiological and molecular study. *Int J Antimicrob Agents*. 2014;44(6):500-507. [PUBMED](#) | [CROSSREF](#)
12. Pungpian C, Sinwat N, Angkittitrakul S, Prathan R, Chuanchuen R. Presence and transfer of antimicrobial resistance determinants in *Escherichia coli* in pigs, pork, and humans in Thailand and Lao PDR border provinces. *Microb Drug Resist*. 2021;27(4):571-584. [PUBMED](#) | [CROSSREF](#)
13. Trongjit S, Angkittitrakul S, Tuttle RE, Pongseree J, Padungtod P, Chuanchuen R. Prevalence and antimicrobial resistance in *Salmonella enterica* isolated from broiler chickens, pigs and meat products in Thailand-Cambodia border provinces. *Microbiol Immunol*. 2017;61(1):23-33. [PUBMED](#) | [CROSSREF](#)
14. Sinwat N, Angkittitrakul S, Coulson KF, Pilapil FM, Meunsene D, Chuanchuen R. High prevalence and molecular characteristics of multidrug-resistant *Salmonella* in pigs, pork and humans in Thailand and Laos provinces. *J Med Microbiol*. 2016;65(10):1182-1193. [PUBMED](#) | [CROSSREF](#)
15. Trongjit S, Angkittitrakul S, Chuanchuen R. Occurrence and molecular characteristics of antimicrobial resistance of *Escherichia coli* from broilers, pigs and meat products in Thailand and Cambodia provinces. *Microbiol Immunol*. 2016;60(9):575-585. [PUBMED](#) | [CROSSREF](#)

16. FAO. *Monitoring and Surveillance of Antimicrobial Resistance in Bacteria from Healthy Food Animals Intended for Consumption*. Bangkok: Food and Agriculture Organization of the United Nations; 2019, 9-26.
17. U.S. Food and Drug Administration. Enumeration of *Escherichia coli* and the coliform bacteria [Internet]. Silver Spring: U.S. Food and Drug Administration; <https://www.fda.gov/food/laboratory-methods-food/bam-chapter-4-enumeration-escherichia-coli-and-coliform-bacteria>. Updated 2017.
18. ISO. *Microbiology of Food and Animal Feeding Stuffs – Horizontal Method for the Detection of Salmonella Spp: ISO6579-Fourth Edition*. Geneva: International Organization for Standardization; 2002.
19. Popoff M, LeMinor L. *Antigenic Formulas of the Salmonella Serovars*. Paris: Institut Pasteur; 1992.
20. CLSI. *Performance Standards for Antimicrobial Disk and Dilution Susceptibility Test for Bacteria Isolated from Animals 3rd VET01-S3*. Wayne: Clinical and Laboratory Standards Institute; 2015.
21. Dafopoulou K, Zarkotou O, Dimitroulia E, Hadjichristodoulou C, Gennimata V, Pournaras S, et al. Comparative evaluation of colistin susceptibility testing methods among carbapenem-nonsusceptible *Klebsiella pneumoniae* and *Acinetobacter baumannii* clinical isolates. *Antimicrob Agents Chemother*. 2015;59(8):4625-4630.  
[PUBMED](#) | [CROSSREF](#)
22. EUCAST. *Breakpoint Tables for Interpretation of MICs and Zone Diameters, Version 7.1, Valid from 2017-03-10*. Basel: European Committee on Antimicrobial Susceptibility Testing; 2017.
23. Lévesque C, Piché L, Larose C, Roy PH. PCR mapping of integrons reveals several novel combinations of resistance genes. *Antimicrob Agents Chemother*. 1995;39(1):185-191.  
[PUBMED](#) | [CROSSREF](#)
24. Rebelo AR, Bortolaia V, Kjeldgaard JS, Pedersen SK, Leekitcharoenphon P, Hansen IM, et al. Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* for surveillance purposes. *Euro Surveill*. 2018;23(6):  
[PUBMED](#) | [CROSSREF](#)
25. Poirel L, Walsh TR, Cuvillier V, Nordmann P. Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis*. 2011;70(1):119-123.  
[PUBMED](#) | [CROSSREF](#)
26. Khemtong S, Chuanchuen R. Class 1 integrons and *Salmonella* genomic island 1 among *Salmonella enterica* isolated from poultry and swine. *Microb Drug Resist*. 2008;14(1):65-70.  
[PUBMED](#) | [CROSSREF](#)
27. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods*. 2005;63(3):219-228.  
[PUBMED](#) | [CROSSREF](#)
28. Sukhnanand S, Alcaine S, Warnick LD, Su WL, Hof J, Craver MP, et al. DNA sequence-based subtyping and evolutionary analysis of selected *Salmonella enterica* serotypes. *J Clin Microbiol*. 2005;43(8):3688-3698.  
[PUBMED](#) | [CROSSREF](#)
29. Burland TG. DNASTAR's Lasergene sequence analysis software. *Methods Mol Biol*. 2000;132:71-91.  
[PUBMED](#)
30. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol*. 2018;35(6):1547-1549.  
[PUBMED](#) | [CROSSREF](#)
31. Zhang X, Zhang B, Guo Y, Wang J, Zhao P, Liu J, et al. Colistin resistance prevalence in *Escherichia coli* from domestic animals in intensive breeding farms of Jiangsu Province. *Int J Food Microbiol*. 2019;291:87-90.  
[PUBMED](#) | [CROSSREF](#)
32. Huang X, Yu L, Chen X, Zhi C, Yao X, Liu Y, et al. High prevalence of colistin resistance and *mcr-1* Gene in *Escherichia coli* isolated from food animals in China. *Front Microbiol*. 2017;8:562.  
[PUBMED](#) | [CROSSREF](#)
33. Xu F, Zeng X, Hinenoya A, Lin J. MCR-1 confers cross-resistance to bacitracin, a widely used in-feed antibiotic. *MSphere*. 2018;3(5):e00411-18.  
[PUBMED](#) | [CROSSREF](#)
34. Strom G, Halje M, Karlsson D, Jiwakanon J, Pringle M, Fernstrom LL, et al. Antimicrobial use and antimicrobial susceptibility in *Escherichia coli* on small- and medium-scale pig farms in north-eastern Thailand. *Antimicrob Resist Infect Control*. 2017;6:75.  
[PUBMED](#) | [CROSSREF](#)
35. Langata LM, Maingi JM, Musonye HA, Kiiru J, Nyamache AK. Antimicrobial resistance genes in *Salmonella* and *Escherichia coli* isolates from chicken droppings in Nairobi, Kenya. *BMC Res Notes*. 2019;12(1):22.  
[PUBMED](#) | [CROSSREF](#)



36. Varga C, Rajić A, McFall ME, Reid-Smith RJ, Deckert AE, Pearl DL, et al. Comparison of antimicrobial resistance in generic *Escherichia coli* and *Salmonella* spp. cultured from identical fecal samples in finishing swine. *Can J Vet Res.* 2008;72(2):181-187.  
[PUBMED](#)
37. Touati M, Hadjadj L, Berrazeg M, Baron SA, Rolain JM. Emergence of *Escherichia coli* harbouring *mcr-1* and *mcr-3* genes in North West Algerian farmlands. *J Glob Antimicrob Resist.* 2020;21:132-137.  
[PUBMED](#) | [CROSSREF](#)
38. Mulvey MR, Bharat A, Boyd DA, Irwin RJ, Wylie J. Characterization of a colistin-resistant *Salmonella enterica* 4,[5],12:i:- harbouring *mcr-3.2* on a variant IncHI-2 plasmid identified in Canada. *J Med Microbiol.* 2018;67(12):1673-1675.  
[PUBMED](#) | [CROSSREF](#)
39. Wise MG, Estabrook MA, Sahm DE, Stone GG, Kazmierczak KM. Prevalence of *mcr*-type genes among colistin-resistant *Enterobacteriaceae* collected in 2014-2016 as part of the INFORM global surveillance program. *PLoS One.* 2018;13(4):e0195281.  
[PUBMED](#) | [CROSSREF](#)
40. Yin W, Li H, Shen Y, Liu Z, Wang S, Shen Z, et al. Novel Plasmid-Mediated Colistin Resistance Gene *mcr-3* in *Escherichia coli*. *MBio.* 2017;8(3):e00543-17.  
[PUBMED](#) | [CROSSREF](#)
41. Liu L, Feng Y, Zhang X, McNally A, Zong Z. New variant of *mcr-3* in an extensively drug-resistant *Escherichia coli* clinical isolate carrying *mcr-1* and *bla<sub>KDM-5</sub>*. *Antimicrob Agents Chemother.* 2017;61(12):e01757-17.  
[PUBMED](#) | [CROSSREF](#)
42. Khine NO, Lugsomya K, Kaewgun B, Honhanrob L, Pairojrit P, Jermprasert S, et al. Multidrug resistance and virulence factors of *Escherichia coli* harboring plasmid-mediated colistin resistance: *mcr-1* and *mcr-3* genes in contracted pig farms in Thailand. *Front Vet Sci.* 2020;7:582899.  
[PUBMED](#) | [CROSSREF](#)
43. Jin L, Wang R, Wang X, Wang Q, Zhang Y, Yin Y, et al. Emergence of *mcr-1* and carbapenemase genes in hospital sewage water in Beijing, China. *J Antimicrob Chemother.* 2018;73(1):84-87.  
[PUBMED](#) | [CROSSREF](#)
44. Poirel L, Kieffer N, Brink A, Coetze J, Jayol A, Nordmann P. Genetic features of MCR-1-producing colistin-resistant *Escherichia coli* isolates in South Africa. *Antimicrob Agents Chemother.* 2016;60(7):4394-4397.  
[PUBMED](#) | [CROSSREF](#)