



## OUTER APPROXIMATION METHOD FOR ZEROS OF SUM OF MONOTONE OPERATORS AND FIXED POINT PROBLEMS IN BANACH SPACES

Hammad Anuoluwapo Abass<sup>1</sup>, Akindele Adebayo Mebawondu<sup>2</sup>,  
Ojen Kumar Narain<sup>3</sup> and Jong Kyu Kim<sup>4</sup>

<sup>1</sup>School of Mathematics, Statistics and Computer Science

University of KwaZulu-Natal, Durban, South Africa,

DSI-NRF Centre of Excellence in Mathematical and Statistical Sciences (CoE-MaSS)

Johannesburg, South Africa

e-mail: [hammedabass548@gmail.com](mailto:hammedabass548@gmail.com)

<sup>2</sup>School of Mathematics, Statistics and Computer Science

University of KwaZulu-Natal, Durban, South Africa,

DSI-NRF Centre of Excellence in Mathematical and Statistical Sciences (CoE-MaSS)

Johannesburg, South Africa,

Department of Computer Science and Mathematics, Mountain Top University

Prayer City, Ogun State, Nigeria

e-mail: [dele@aims.ac.za](mailto:dele@aims.ac.za)

<sup>3</sup>School of Mathematics, Statistics and Computer Science

University of KwaZulu-Natal, Durban, South Africa

e-mail: [naraino@ukzn.ac.za](mailto:naraino@ukzn.ac.za)

<sup>4</sup>Department of Mathematics Education, Kyungnam University

Changwon, Gyeongnam, 51767, Korea

e-mail: [jongkyuk@kyungnam.ac.kr](mailto:jongkyuk@kyungnam.ac.kr)

**Abstract.** In this paper, we investigate a hybrid algorithm for finding zeros of the sum of maximal monotone operators and Lipschitz continuous monotone operators which is also a common fixed point problem for finite family of relatively quasi-nonexpansive mappings and split feasibility problem in uniformly convex real Banach spaces which are also uniformly smooth. The iterative algorithm employed in this paper is design in such a way that it does not require prior knowledge of operator norm. We prove a strong convergence result for approximating the solutions of the aforementioned problems and give applications of our main result to minimization problem and convexly constrained linear inverse problem.

---

<sup>0</sup>Received August 27, 2020. Revised January 26, 2021. Accepted February 19, 2021.

<sup>0</sup>2010 Mathematics Subject Classification: 47H06, 47H09, 47J05, 47J25.

<sup>0</sup>Keywords: Maximal monotone operators, relatively quasi-nonexpansive mapping, hybrid iterative scheme, split feasibility problem, fixed point problem.

<sup>0</sup>Corresponding author: H. A. Abass([hammedabass548@gmail.com](mailto:hammedabass548@gmail.com)).

## 1. INTRODUCTION

The monotone inclusion problem is to find an element  $x \in H$  such that

$$0 \in B(x),$$

where  $B : H \rightarrow 2^H$  is a multi-valued operator and  $H$  is a real Hilbert space. This problem is very important in many areas such as convex optimization and monotone variational inequality problems. It is worth mentioning that every monotone operator on Hilbert spaces can be regularized into single-valued, nonexpansive, Lipschitz continuous monotone operator by means of Yosida approximation notion. The inclusion problem can also be defined in terms of sum of two monotone operators  $M$  and  $B$ , where one of these operators is  $\alpha$ -inverse strongly monotone which is  $\frac{1}{\alpha}$ -Lipschitz continuous.

Let  $E$  be a real Banach space with  $\|\cdot\|$ , dual space  $E^*$  and  $\langle f, x \rangle$  the value of  $f \in E^*$  at  $x \in E$ . Let  $B : E \rightarrow 2^{E^*}$  be a maximal monotone operator and  $M : E \rightarrow E^*$  be a Lipschitz continuous monotone operator.

In this paper, we consider the following inclusion problem: find  $x \in E$  such that

$$0 \in (M + B)x. \tag{1.1}$$

We denote by  $(M + B)^{-1}(0)$  the solution set of (1.1).

Based on a series of studies in the past years, the splitting method has been known to be a popular method for solving (1.1). The splitting methods for linear equations were introduced by Peaceman and Rashford [27]. Extensions to nonlinear equations in Hilbert spaces were carried out by Lions and Mercier [22]. Since then, many authors have considered approximating solutions of variational inclusion (1.1) using this method, (see [2, 3, 12, 18, 20, 21, 31, 36] and the references contained in).

Recently, Zhang and Jiang [42] proved the following strong convergence theorem for approximating solutions for a common zero point of the sum of two monotone operators which is also a fixed point of a family of countable quasi-nonexpansive mapping in the framework of Hilbert spaces as follows:

**Theorem 1.1.** ([42]) *Let  $C$  be a nonempty, closed and convex subset of a real Hilbert space  $H$ ,  $A : C \rightarrow H$  be an  $\alpha$ -inverse strongly monotone operator and  $B$  be a maximal monotone operator on  $H$  such that  $\text{Dom}(B)$  is included in  $C$ . Let  $\{S_n\} : C \rightarrow C$  be a family of countable quasi-nonexpansive mappings which are uniformly closed. Assume that*

$$\Gamma := F(S_n) \cap (A + B)^{-1}(0) \neq \emptyset.$$

Let  $\{r_n\}$  be a positive real number sequence and  $\{\alpha_n\}$  be a real number sequence in  $(0,1)$ . Let  $\{x_n\}$  be a sequence of  $C$  generated by

$$\begin{cases} x_1 \in C_1 = C, \text{ chosen arbitrarily;} \\ z_n = J_{r_n}(x_n - r_n Ax_n); \\ y_n = \alpha_n z_n + (1 - \alpha_n) S_n z_n; \\ C_{n+1} = \{z \in C_n : \|z_n - z\| \leq \|y_n - z\| \leq \|x_n - z\|\}; \\ x_{n+1} = P_{C_{n+1}} x_1, \quad n \geq 1; \end{cases}$$

where  $J_{r_n} = (I + r_n B)^{-1}$ ,  $\liminf_{n \rightarrow \infty} r_n > 0$ ,  $r_n \leq 2\alpha$  and  $\limsup_{n \rightarrow \infty} \alpha_n < 1$ . Then the sequence  $\{x_n\}$  converges strongly to  $q = P_{\Gamma} x_0$ .

Very recently, Shehu [31] considered splitting method for finding zeros of the sum of maximal monotone operator and Lipschitz continuous monotone operator in Banach spaces. He proved weak and strong convergence results and give some applications of his main result.

The split feasibility problem (SFP) introduced by Censor and Elfving [10] is to find an element

$$x^* \in C \text{ such that } Ax^* \in Q, \tag{1.2}$$

where  $C$  and  $Q$  are nonempty, closed and convex subsets of real Banach spaces  $E_1$  and  $E_2$  respectively, and  $A : E_1 \rightarrow E_2$  is a bounded linear operator. The SFP arises from phase retrievals and in medical image reconstruction to mention a few. For more details on SFP, we refer readers to (see [1, 2, 3, 11, 19, 23, 25, 26, 37] and other references therein).

In 2018, Ma et al. [23] introduced an iterative algorithm to solve the SFP (1.2) and fixed point problem of quasi- $\phi$ -nonexpansive mappings in Banach spaces. They proved a strong convergence result to a common solution of the aforementioned problems and apply their result to convexly constrained inverse problem and split null point problem.

**Remark 1.2.**

- (1) We observe that the inclusion problem considered in [42] is quite different from the one in (1.1) in the sense that one of the operators is a Lipschitz continuous monotone operator.
- (2) The iterative algorithm employed in this article does not require prior knowledge of operator norm as the ones employed in [23] requires prior knowledge of operator norm which gives difficulties in computation.
- (3) We extend the result of [42] from Hilbert spaces to a more general Banach spaces.
- (4) The split feasibility problem considered in this paper finds its applications in signal processing, image reconstruction and medical care.

Motivated by the works of Shehu [31], Zhang and Jiang [42] and Ma et al. [23], we introduced a shrinking iterative algorithm for finding zeros of the sum of maximal monotone operators and Lipschitz continuous monotone operators, which is also a common fixed point of a finite family of relatively quasi-nonexpansive mappings and split feasibility problem in Banach spaces. We prove a strong convergence result for approximating solutions of the aforementioned problems and give applications of our main result to minimization problem and convexly constrained linear inverse problem. The result present in this paper extends the result of Ma et al. [23], Zhang and Jiang [42] and other related results in literature.

## 2. PRELIMINARIES

We give some definitions and important results which will be useful in establishing our main results. In the sequel, we denote strong and weak convergence by " $\rightarrow$ " and " $\rightharpoonup$ ", respectively.

Throughout this paper, we assume  $C$  to be a nonempty, closed and convex subset of a real Banach space with norm  $\|\cdot\|$ ,  $J : E \rightarrow 2^{E^*}$  be the normalized duality mapping defined by

$$J(x) = \{x^* \in E^* : \langle x, x^* \rangle = \|x\|^2 = \|x^*\|^2, \forall x \in E\}.$$

Consider the Lyapunov functional  $\phi : E \times E \rightarrow [0, \infty)$  defined [4, 5] by

$$\phi(x, y) = \|x\|^2 - 2\langle x, Jy \rangle + \|y\|^2, \forall x, y \in E.$$

Alber [4] introduced a generalized projection operator  $\Pi_C : E \rightarrow C$  which is an analogue of the metric projection defined as follows:

$$\Pi_C(x) = \operatorname{argmin}_{y \in C} \phi(y, x), x \in E.$$

That is,  $\Pi_C(x) = \bar{x}$ , where  $\bar{x}$  is the unique solution to the minimization problem  $\phi(\bar{x}, x) = \inf_{y \in C} \phi(y, x)$ . In real Hilbert space, we observe that  $\Pi_C(x) \equiv P_C(x)$  and  $\phi(x, y) = \|x - y\|^2$ . It is obvious from the definition of the functional  $\phi$  that

$$(\|x\| - \|y\|)^2 \leq \phi(x, y) \leq (\|x\| + \|y\|)^2. \quad (2.1)$$

Apart from inequality (2.1), the Lyapunov functional  $\phi$  also satisfy the following inequalities:

- (A<sub>1</sub>)  $\phi(x, y) = \phi(x, z) + \phi(z, y) + 2\langle x - z, Jz - Jy \rangle;$
- (A<sub>2</sub>)  $2\langle x - y, Jz - Jw \rangle = \phi(x, w) + \phi(y, z) - \phi(x, z) - \phi(y, w);$
- (A<sub>3</sub>)  $\phi(x, y) \leq \|x\| \|Jx - Jy\| + \|y\| \|x - y\|.$

**Note:** If  $E$  is a reflexive, strictly convex, and smooth Banach space, then for  $x, y \in E$ ,  $\phi(x, y) = 0$  if and only if  $x = y$ , see [35].

We are also concerned with the functional  $V : E \times E^* \rightarrow \mathbb{R}$  which is defined by

$$V(x, x^*) = \|x\|^2 - 2\langle x, x^* \rangle + \|x^*\|^2 \tag{2.2}$$

for all  $x \in E$  and  $x^* \in E^*$ . Observe that,  $V(x, x^*) = \phi(x, J^{-1}x^*)$ , if  $E$  is a reflexive, strictly convex and smooth Banach space and

$$V(x, x^*) \leq V(x, x^* + y^*) - 2\langle J^{-1}x^* - x, y^* \rangle \tag{2.3}$$

for all  $x \in E$  and all  $x^*, y^* \in E^*$ , see [30].

Let  $C$  be a closed and convex subset of  $E$  and  $T : C \rightarrow C$  be a mapping. Then,  $x \in C$  is called a fixed point of  $T$ , if  $x = Tx$ . We denote the set of fixed points of  $T$  by  $F(T)$ . A point  $p \in C$  is called an asymptotic fixed point of  $T$ , if  $C$  contains a sequence  $\{x_n\}$  such that  $x_n \rightarrow p$  and  $\|x_n - Tx_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . We denote by  $\widehat{F(T)}$  the set of asymptotic fixed points of  $T$ . A mapping  $T : C \rightarrow C$  is said to be relatively nonexpansive (see [24]) if the following conditions are satisfied:

- (L1)  $F(T) \neq \emptyset$ ;
- (L2)  $\phi(p, Tx) \leq \phi(p, x), \forall x \in C, p \in F(T)$ ;
- (L3)  $F(T) = \widehat{Fix(T)}$ .

If  $T$  satisfies (L1) and (L2), then  $T$  is said to be relatively quasi-nonexpansive. It is easy to see that the class of relative quasi-nonexpansive mappings contains the class of relatively nonexpansive mappings. Many authors have considered the relative quasi-nonexpansive mappings, (see [33, 38]).

**Definition 2.1.** Let  $X \subset E$  be a nonempty subset. Then a mapping  $A : X \rightarrow E^*$  is called

- (i)  $\gamma$ -strongly monotone with modulus  $\gamma > 0$  on  $X$  if

$$\langle Ax - Ay, x - y \rangle \geq \gamma \|x - y\|^2, \forall x, y \in X;$$

- (ii) monotone on  $X$  if

$$\langle Ax - Ay, x - y \rangle \geq 0, \forall x, y \in X;$$

- (iii) Lipschitz continuous on  $X$  if there exists a constant  $L > 0$  such that

$$\|Ax - Ay\| \leq L \|x - y\|, \forall x, y \in X.$$

Below is an example of a monotone operator in quantum mechanics.

**Example 2.2.** ([31]) Let the operator

$$Au := -b^2 \Delta u + (f(x) + c)u(x) + u(x) \int_{\mathbb{R}^3} \frac{u^2(y)}{|x - y|} dy,$$

where  $\Delta := \sum_{i=1}^3 \frac{\partial^2}{\partial x_i^2}$  is the Laplacian in  $\mathbb{R}^3$ ,  $b$  and  $c$  are constants,  $f(x) = f_0(x) + f_1(x)$ , where  $f_0(x) \in L^\infty(\mathbb{R}^3)$  and  $f_1(x) \in L^2(\mathbb{R}^3)$ . Let  $A := L + B$ ,

where the operator  $L$  which is the schrödinger operator is the linear part of  $A$  and  $B$  defined by the last term. It is known that  $B$  is a monotone operator on  $L^2(\mathbb{R}^3)$ , (see p.23 of [6]) which also implies that  $A : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)$  is also a monotone operator.

**Definition 2.3.** A multi-valued operator  $B : E \rightarrow 2^{E^*}$  with domain  $Dom(B) = \{x \in E : Bx \neq 0\}$  and the range  $R(B) = \{Bx : x \in D(B)\}$  is said to be monotone if for  $x, y \in D(B)$ ,  $a \in Bx, b \in By$ , the following inequality holds:

$$\langle x - y, a - b \rangle \geq 0.$$

A monotone operator  $B$  is said to be maximal if its graph  $Gra(B) = \{(x, y) : y \in Bx\}$  is not properly contained in the graph of any other monotone operator.

If  $E$  is a strictly convex, reflexive and smooth Banach space and  $B : E \rightarrow 2^{E^*}$  is a maximal monotone operator. Then, for any positive real number  $\lambda$ , we can define a nonexpansive single-valued operator  $J_\lambda^B : E \rightarrow E$  by

$$J_\lambda^B(x) := (J + \lambda B)^{-1}J(x), \quad x \in E.$$

This operator is called the resolvent of  $B$  for  $\lambda > 0$ . It is well known that  $B^{-1}(0) = F(J_\lambda^B)$  for all  $\lambda > 0$  and  $B^{-1}(0)$  is a closed and convex subset of  $E$ .

For a real Banach space  $E$ , the modulus of convexity of  $E$  is the function  $\delta_E : [0, 2] \rightarrow [0, 1]$  defined as

$$\delta_E(\epsilon) = \inf \left\{ 1 - \frac{1}{2} \|x + y\| : \|x\| = \|y\| = 1, \|x - y\| \geq \epsilon \right\}. \quad (2.4)$$

Recall that  $E$  is said to be uniformly convex if  $\delta_E(\epsilon) > 0$  for any  $\epsilon \in (0, 2]$ .  $E$  is said to be strictly convex if  $\frac{\|x + y\|}{2} < 1$  for all  $x, y \in E$  with  $\|x\| = \|y\| = 1$  and  $x \neq y$ . Also,  $E$  is  $p$ -uniformly convex if there exists a constant  $c_p > 0$  such that  $\delta_E(\epsilon) > c_p \epsilon^p$  for any  $\epsilon \in (0, 2]$ .

The modulus of smoothness of  $E$  is the function  $\rho_E : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  defined by

$$\rho_E(t) = \sup \left\{ \frac{1}{2} (\|x + ty\| - \|x - ty\|) - 1 : \|x\| = \|y\| = 1 \right\}. \quad (2.5)$$

$E$  is said to be uniformly smooth if  $\lim_{t \rightarrow 0} \frac{\rho_E(t)}{t} = 0$ . Let  $1 < q \leq 2$ . Then  $E$  is  $q$ -uniformly smooth if there exists  $c_q > 0$  such that  $\rho_E(t) \leq c_q t^q$  for  $t > 0$ . It is known that  $E$  is  $p$ -uniformly convex if and only if  $E^*$  is  $q$ -uniformly smooth, where  $p^{-1} + q^{-1} = 1$ . It is also known that every  $q$ -uniformly smooth Banach space is uniformly smooth. It is also widely known that if  $E$  is uniformly smooth, then the duality mapping  $J$  is norm-to-norm continuous on each bounded subset of  $E$ .

The following are some important and useful properties of duality mapping  $J$ , for further details see [35]:

- For every  $x \in E$ ,  $Jx$  is nonempty, closed, convex and bounded subset of  $E^*$ .
- If  $E$  is smooth or  $E^*$  is strictly convex, then  $J$  is single-valued. Also, If  $E$  is reflexive, then  $J$  is onto.
- If  $E$  is strictly convex, then  $J$  is strictly monotone, that is

$$\langle x - y, Jx - Jy \rangle > 0, x \neq y \quad \forall x, y \in E.$$

- If  $E$  is smooth, strictly convex and reflexive and  $J^* : E^* \rightarrow 2^E$  is the normalized duality mapping on  $E^*$ , then  $J^{-1} = J^*$ ,  $JJ^* = I_{E^*}$  and  $J^*J = I_E$ , where  $I_E$  and  $I_{E^*}$  are the identity mappings on  $E$  and  $E^*$  respectively.
- If  $E$  is uniformly convex and uniformly smooth, then  $J$  is uniformly norm-to-norm continuous on bounded subsets of  $E$  and  $J^* = J^{-1}$  is also uniformly norm-to-norm continuous on bounded subsets of  $E^*$ .

We now state the following results which will be useful to prove our main result.

**Lemma 2.4.** ([8]) *Let  $\frac{1}{p} + \frac{1}{q} = 1$ , for  $p, q > 1$ . Then, the space  $E$  is  $q$ -uniformly smooth if and only if its dual space  $E^*$  is  $p$ -uniformly convex.*

**Lemma 2.5.** ([39]) *Let  $E$  be a 2-uniformly smooth Banach space with the best smoothness constant  $k > 0$ . Then, the following inequality holds:*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, Jx \rangle + 2\|ky\|^2, \quad \forall x, y \in E.$$

**Lemma 2.6.** ([39]) *Given a number  $r > 0$ , a real Banach space  $E$  is uniformly convex if and only if there exists a continuous strictly increasing function  $g : [0, \infty) \rightarrow [0, \infty)$  with  $g(0) = 0$  such that*

$$\|\lambda x + (1 - \lambda)y\|^2 \leq \lambda\|x\|^2 + (1 - \lambda)\|y\|^2 - \lambda(1 - \lambda)g(\|x - y\|);$$

for all  $x, y \in E$  with  $\|x\| \leq r$  and  $\|y\| \leq r$  and  $\lambda \in [0, 1]$ .

**Lemma 2.7.** ([5]) *Let  $E$  be a smooth, strictly convex and reflexive Banach space and  $C$  be a nonempty closed convex subset of  $E$ . Then, the following conclusions hold:*

- (i)  $\phi(x, \Pi_C y) + \phi(\Pi_C y, y) \leq \phi(x, y), \quad \forall x \in C, y \in E.$
- (ii) *If  $x \in E$  and  $z \in C$ , then  $z = \Pi_C x$  if and only if*

$$\langle z - y, Jx - Jz \rangle \geq 0, \quad \forall y \in C.$$

- (iii) *For  $x, y \in E$ ,  $\phi(x, y) = 0$  if and only if  $x = y$ .*

**Lemma 2.8.** ([17]) *Let  $E$  be a uniformly convex and smooth Banach space, and let  $\{x_n\}, \{y_n\}$  be two sequences of  $E$ . If  $\phi(x_n, y_n) \rightarrow 0$  and either of  $\{x_n\}$  or  $\{y_n\}$  is bounded. Then,  $\|x_n - y_n\| \rightarrow 0$ .*

**Lemma 2.9.** ([7]) *Let  $E$  be a real uniformly convex, smooth Banach space. Then, the following identities hold:*

- (i)  $\phi(x, y) = \phi(x, z) + \phi(z, y) + 2\langle x - z, Jz - Jy \rangle, \forall x, y \in E;$
- (ii)  $\phi(x, y) + \phi(y, x) = 2\langle x - y, Jx - Jy \rangle, \forall x, y \in E.$

**Lemma 2.10.** ([35]) *Let  $E$  be a smooth, strictly convex, and reflexive Banach space. Let  $C$  be a nonempty, closed and convex subset of  $E$  and let  $x_1 \in E$  and  $z \in C$ . Then, the following conclusions hold:*

- (i)  $z = P_C x_1,$
- (ii)  $\langle z - y, J(x_1 - z) \rangle \geq 0, \forall y \in C.$

**Lemma 2.11.** ([9]) *Let  $B : E \rightarrow 2^{E^*}$  be a maximal monotone operator and  $M : E \rightarrow E^*$  be a Lipschitz continuous monotone operator. Then the operator  $M + B$  is a maximal monotone operator.*

**Lemma 2.12.** ([31]) *Let  $B : E \rightarrow 2^{E^*}$  be a maximal monotone operator and  $M : E \rightarrow E^*$  be an operator. Define an operator*

$$T_\lambda x := J_\lambda^B \circ J^{-1}(J - \lambda M), \quad x \in E, \quad \lambda > 0.$$

*Then  $F(T_\lambda) = (M + B)^{-1}(0)$ .*

### 3. MAIN RESULT

We suppose that  $E$  is  $p$ -uniformly convex and uniformly smooth, which implies that its dual space  $E^*$  is  $q$ -uniformly smooth and uniformly convex. Throughout this section, we assume that  $E_1$  is a 2-uniformly convex real Banach space which is also 2-uniformly smooth and  $E_2$  is a smooth, strictly convex and reflective Banach space,  $E_1^*$  is a 2-uniformly smooth real Banach space which is also uniformly convex. Furthermore, we suppose that  $J_1$  and  $J_2$  represent the normalized duality mapping of  $E_1$  and  $E_2$  respectively and  $J_1 = (J_1^*)^{-1}$ , where  $J_1^*$  is the normalized duality mapping of  $E_1^*$ .

**Theorem 3.1.** *Let  $E_1$  be 2-uniformly convex and 2-uniformly smooth real Banach space with the best smoothness constant  $0 < k \leq \frac{1}{\sqrt{2}}$ ,  $E_2$  be a smooth, strictly convex and reflective Banach space. Let  $\{S_i\}_{i=1}^N : E_1 \rightarrow E_1$  be a finite family of closed relatively quasi-nonexpansive mapping,  $A : E_1 \rightarrow E_2$  be a bounded linear operator with adjoint  $A^*$  and  $Q$  be a nonempty, closed and convex subset of  $E_2$ . Suppose that  $B : E_1 \rightarrow 2^{E_1^*}$  is a maximal monotone operator and  $M : E_1 \rightarrow E_1^*$  is monotone and  $L$ -Lipschitz continuous. Assume that*

$$\Gamma := \{\bar{x} \in \bigcap_{i=1}^N F(S_i) \cap (B + M)^{-1}(0) : A\bar{x} \in Q\} \neq \emptyset.$$



Let  $x_1 \in E_1$ ,  $C_1 = E_1$ , and  $\{x_n\}$  be a sequence generated by

$$\begin{cases} w_n = J_1^{-1}(J_1x_n + \gamma_n A^* J_2(P_Q - I)Ax_n); \\ y_n = J_{\lambda_n}^B \circ J_1^{-1}(J_1w_n - \lambda_n M w_n); \\ u_n = J_1^{-1}(J_1y_n - \lambda_n(My_n - Mx_n)); \\ t_n = J_1^{-1}[(1 - \alpha_n)J_1u_n + \alpha_n J_1S_i u_n]; \\ C_{n+1} = \{v \in C_n : \phi(v, t_n) \leq \phi(v, x_n)\}; \\ x_{n+1} = \Pi_{C_{n+1}}x_1; \quad n \geq 1; \end{cases} \tag{3.1}$$

where  $P_Q$  is the metric projection of  $E_2$  onto  $Q$  and  $\Pi_{C_{n+1}}$  is the generalized projection of  $E_1$  onto  $C_{n+1}$ . Suppose  $\{\alpha_n\}_{n=1}^\infty$  is a sequence in  $(0, 1)$  such that

$$\liminf_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0,$$

and the step size  $\gamma_n$  is chosen in such a way that for  $\varepsilon > 0$ ,

$$\gamma_n \in \left( \varepsilon, \frac{\|(P_Q - I)Ax_n\|^2}{\|A^*k^2J_2(P_Q - I)Ax_n\|^2} - \varepsilon \right),$$

for all  $P_Q Ax_n \neq Ax_n$ ,  $\gamma_n = \gamma$  otherwise ( $\gamma$  being any nonnegative real number) with  $\{\lambda_n\}_{n=1}^\infty$  satisfying the following condition:

$$0 < d \leq \lambda_n \leq e < \frac{1}{\sqrt{2\mu\rho L}},$$

where  $\mu$  is the 2-uniform convexity constant of  $E_1$ ,  $\rho$  is the 2-uniform smoothness constant of  $E_1^*$ , and  $L$  is the Lipschitz constant of  $M$ . Then,  $\{x_n\}$  converges strongly to a point  $\bar{x} = \Pi_\Gamma x_1$ .

*Proof.* We divide our proof into several steps:

**Step 1:** We prove using Theorem 3.1 that  $C_n$  is closed and convex for each  $n \geq 1$ . We obtain from Theorem 3.1 that  $C_1 = E_1$ , therefore  $C_1$  is closed and convex. Now assume that  $C_n$  is closed and convex, then we have

$$\phi(v, t_n) \leq \phi(v, x_n).$$

It means that

$$\|v\|^2 - 2\langle v, J_1t_n \rangle + \|t_n\|^2 \leq \|v\|^2 - 2\langle v, J_1x_n \rangle + \|x_n\|^2.$$

Hence we obtain that

$$2\langle v, J_1x_n - J_1t_n \rangle \leq \|x_n\|^2 - \|t_n\|^2. \tag{3.2}$$

We have from (3.2) that  $C_{n+1}$  is closed and convex subset of  $E_1$ . Therefore,  $\Pi_{C_{n+1}}$  is well-defined.

**Step 2:** We show that  $\Gamma \subseteq C_n$  for all  $n \geq 1$ . Let  $x^* \in \Gamma \subseteq C_n$ , for some  $n \geq 1$ . Then we have from (3.1) and Lemma 2.6 that

$$\begin{aligned}
\phi(x^*, t_n) &= \phi(x^*, J_1^{-1}((1 - \alpha_n)J_1u_n + \alpha_nJ_1S_iu_n)) \\
&= \|x^*\|^2 - 2\langle x^*, (1 - \alpha_n)J_1u_n + \alpha_nJ_1S_iu_n \rangle \\
&\quad + \|(1 - \alpha_n)J_1u_n + \alpha_nJ_1S_iu_n\|^2 \\
&\leq \|x^*\|^2 - 2(1 - \alpha_n)\langle x^*, J_1u_n \rangle - 2\alpha_n\langle x^*, J_1S_iu_n \rangle + (1 - \alpha_n)\|u_n\|^2 \\
&\quad + \alpha_n\|S_iu_n\|^2 - \alpha_n(1 - \alpha_n)g(\|J_1u_n - J_1S_iu_n\|) \\
&= (1 - \alpha_n)\phi(x^*, u_n) + \alpha_n\phi(x^*, S_iu_n) \\
&\quad - \alpha_n(1 - \alpha_n)g(\|J_1u_n - J_1S_iu_n\|) \\
&\leq (1 - \alpha_n)\phi(x^*, u_n) + \alpha_n\phi(x^*, u_n) \\
&\quad - \alpha_n(1 - \alpha_n)g(\|J_1u_n - J_1S_iu_n\|) \\
&= \phi(x^*, u_n) - \alpha_n(1 - \alpha_n)g(\|J_1u_n - J_1S_iu_n\|) \\
&\leq \phi(x^*, u_n) \\
&= \phi(x^*, J_1^{-1}(J_1y_n - \lambda_n(My_n - Mw_n))) \\
&= \|x^*\|^2 - 2\langle x^*, (J_1y_n - \lambda_n(My_n - Mw_n)) \rangle \\
&\quad + \|J_1^{-1}(J_1y_n - \lambda_n(My_n - Mw_n))\|^2 \\
&= \|x^*\|^2 - 2\langle x^*, J_1y_n - \lambda_n(My_n - Mw_n) \rangle \\
&\quad + \|(J_1y_n - \lambda_n(My_n - Mw_n))\|^2 \\
&= \|x^*\|^2 - 2\langle x^*, J_1y_n \rangle + 2\lambda_n\langle x^*, My_n - Mw_n \rangle \\
&\quad + \|J_1y_n - \lambda_n(My_n - Mw_n)\|^2. \tag{3.3}
\end{aligned}$$

But from Lemma 2.5, we have that

$$\begin{aligned}
\|J_1y_n - \lambda_n(My_n - Mw_n)\|^2 &\leq \|J_1y_n\|^2 - 2\lambda_n\langle My_n - Mw_n, y_n \rangle \\
&\quad + 2k^2\|\lambda_n(My_n - Mw_n)\|^2. \tag{3.4}
\end{aligned}$$

On substituting (3.4) into (3.3), we obtain

$$\begin{aligned}
\phi(x^*, u_n) &\leq \|x^*\|^2 - 2\langle x^*, J_1y_n \rangle + 2\lambda_n\langle x^*, My_n - Mw_n \rangle + \|J_1y_n\|^2 \\
&\quad - 2\lambda_n\langle My_n - Mw_n, y_n \rangle + 2k^2\|\lambda_n(My_n - Mw_n)\|^2 \\
&= \|x^*\|^2 - 2\lambda_n\langle My_n - Mw_n, y_n - x^* \rangle - 2\langle x^*, J_1y_n \rangle \\
&\quad + 2k^2\|\lambda_n(My_n - Mw_n)\|^2 + \|y_n\|^2 \\
&= \phi(x^*, y_n) - 2\lambda_n\langle My_n - Mw_n, y_n - x^* \rangle \\
&\quad + 2k^2\|\lambda_n(My_n - Mw_n)\|^2. \tag{3.5}
\end{aligned}$$

Observe from Lemma 2.9 (i) that

$$\begin{aligned}\phi(x^*, y_n) &= \phi(x^*, w_n) + \phi(w_n, y_n) + 2\langle x^* - w_n, J_1 w_n - J_1 y_n \rangle \\ &= \phi(x^*, w_n) + \phi(w_n, y_n) + 2\langle w_n - x^*, J_1 y_n - J_1 w_n \rangle.\end{aligned}\quad (3.6)$$

Also, using Lemma 2.9 (ii), we get

$$\phi(w_n, y_n) = 2\langle y_n - w_n, J_1 y_n - J_1 w_n \rangle - \phi(y_n, w_n). \quad (3.7)$$

On substituting (3.6) and (3.7) into (3.5), we obtain that

$$\begin{aligned}\phi(x^*, u_n) &= \phi(x^*, w_n) + \phi(w_n, y_n) + 2\langle w_n - x^*, J_1 y_n - J_1 w_n \rangle \\ &\quad - 2\lambda_n \langle My_n - Mw_n, y_n - x^* \rangle + 2k^2 \|\lambda_n(My_n - Mw_n)\|^2 \\ &= \phi(x^*, w_n) + \phi(w_n, y_n) - 2\langle y_n - w_n, J_1 y_n - J_1 w_n \rangle \\ &\quad + 2\langle y_n - x^*, J_1 y_n - J_1 w_n \rangle - 2\lambda_n \langle My_n - Mw_n, y_n - x^* \rangle \\ &\quad + 2k^2 \|\lambda_n(My_n - Mw_n)\|^2 \\ &\leq \phi(x^*, w_n) - \phi(y_n, w_n) + 2\langle y_n - x^*, J_1 y_n - J_1 w_n \rangle \\ &\quad - 2\lambda_n \langle My_n - Mw_n, y_n - x^* \rangle + 2k^2 \|\lambda_n(My_n - Mw_n)\|^2 \\ &= \phi(x^*, w_n) - \phi(y_n, w_n) + 2k^2 \|\lambda_n(My_n - Mw_n)\|^2 \\ &\quad - 2\langle J_1 w_n - J_1 y_n - \lambda_n(Mw_n - My_n), y_n - x^* \rangle.\end{aligned}\quad (3.8)$$

Using (3.1), it is clear that

$$J_1 w_n - \lambda_n M w_n \in (J_1 + \lambda_n B) y_n.$$

Also, using the fact that  $B$  is a maximal monotone, there exists  $r_n \in B y_n$  such that

$$J_1 w_n - \lambda_n M w_n = J_1 y_n + \lambda_n r_n.$$

Hence

$$r_n = \frac{1}{\lambda_n} (J_1 w_n - J_1 y_n - \lambda_n M w_n). \quad (3.9)$$

Since  $M + B$  is maximal monotone and  $M y_n + r_n \in (M + B) y_n$ , we obtain

$$\langle M y_n + r_n, y_n - x^* \rangle \geq 0. \quad (3.10)$$

On substituting (3.9) into (3.10), we have

$$\langle J_1 w_n - J_1 y_n - \lambda_n (M w_n - M y_n), y_n - x^* \rangle \geq 0. \quad (3.11)$$

Using (3.11) in (3.8), we obtain that

$$\begin{aligned}\phi(x^*, u_n) &\leq \phi(x^*, w_n) - \phi(y_n, w_n) + 2k^2 \|\lambda_n(My_n - Mw_n)\|^2 \\ &\leq \phi(x^*, w_n) - \phi(y_n, w_n) + 2k^2 \lambda_n^2 L^2 \mu \phi(y_n, w_n) \\ &\leq \phi(x^*, w_n) - (1 - 2k^2 \lambda_n^2 L^2 \mu) \phi(y_n, w_n).\end{aligned}\quad (3.12)$$

By applying the condition on  $\lambda_n$ , (3.6), we have that

$$\begin{aligned}
\phi(x^*, u_n) &\leq \phi(x^*, y_n) \\
&\leq \phi(x^*, w_n) \\
&= \phi(x^*, J_1^{-1}(J_1 x_n + \gamma_n A^* J_2(P_Q - I)Ax_n)) \\
&= \|x^*\|^2 - 2\langle x^*, J_1 x_n + \gamma_n A^* J_2(P_Q - I)Ax_n \rangle \\
&\quad + \|J_1 x_n + \gamma_n A^* J_2(P_Q - I)Ax_n\|^2 \\
&\leq \|x^*\|^2 - 2\langle x^*, J_1 x_n \rangle - 2\langle x^*, \gamma_n A^* J_2(P_Q - I)Ax_n \rangle \\
&\quad + \|x_n\|^2 + 2\gamma_n \langle Ax_n, J_2(P_Q - I)Ax_n \rangle + 2\|k\gamma_n A^* J_2(P_Q - I)Ax_n\|^2 \\
&\leq \phi(x^*, x_n) - 2\gamma_n \langle Ax^* - Ax_n, J_2(P_Q - I)Ax_n \rangle \\
&\quad + 2k^2 \gamma_n^2 \|A^* J_2(P_Q - I)Ax_n\|^2. \tag{3.13}
\end{aligned}$$

But, from Lemma 2.10, we have that

$$\begin{aligned}
&\langle Ax^* - Ax_n, J_2(P_Q - I)Ax_n \rangle \\
&= \langle Ax^* - P_Q Ax_n + P_Q Ax_n - Ax_n, J_2(P_Q - I)Ax_n \rangle \\
&= \langle Ax^* - P_Q Ax_n, J_2(P_Q - I)Ax_n \rangle \\
&\quad + \langle (P_Q - I)Ax_n, J_2(P_Q - I)Ax_n \rangle \\
&= \langle Ax^* - P_Q Ax_n, J_2(P_Q - I)Ax_n \rangle + \|(P_Q - I)Ax_n\|^2 \\
&\geq \|(P_Q - I)Ax_n\|^2. \tag{3.14}
\end{aligned}$$

On substituting (3.14) into (3.13), we obtain that

$$\begin{aligned}
&\phi(x^*, w_n) \\
&\leq \phi(x^*, x_n) - 2\gamma_n \|(P_Q - I)Ax_n\|^2 + 2k^2 \gamma_n^2 \|A^* J_2(P_Q - I)Ax_n\|^2 \\
&\leq \phi(x^*, x_n) - 2\gamma_n [\|(P_Q - I)Ax_n\|^2 - k^2 \gamma_n \|A^* J_2(P_Q - I)Ax_n\|^2]. \tag{3.15}
\end{aligned}$$

By applying the condition on  $\gamma_n$  in Theorem 3.1 we have that

$$\phi(x^*, u_n) \leq \phi(x^*, w_n) \leq \phi(x^*, x_n), \tag{3.16}$$

and hence

$$\phi(x^*, t_n) \leq \phi(x^*, x_n). \tag{3.17}$$

We therefore conclude that  $x^* \in C_{n+1}$ . This implies that  $\Gamma \subseteq C_n$  for all  $n \geq 1$ . Hence, (3.1) is well-defined.

**Step 3:** We show that  $\{x_n\}$  is a Cauchy sequence. Let  $x^* \in \Gamma$ , by using the definition of  $C_n$ , we have that  $x_n = \Pi_{C_n} x_1$  for all  $n \geq 1$ . It follows from

Lemma 2.7, we have that

$$\begin{aligned} \phi(x_n, x_1) &= \phi(\Pi_{C_n} x_1, x_1) \\ &\leq \phi(x^*, x_1) - \phi(x^*, \Pi_{C_n} x_1) \\ &\leq \phi(x^*, x_1), \quad \forall n \geq 1. \end{aligned}$$

This implies that  $\{\phi(x_n, x_1)\}$  is bounded. More so, since  $x_n = \Pi_{C_n} x_1$  and  $x_{n+1} = \Pi_{C_{n+1}} x_1 \in C_{n+1} \subseteq C_n$ , we have that

$$\phi(x_n, x_1) \leq \phi(x_{n+1}, x_1), \quad \forall n \geq 1. \tag{3.18}$$

Therefore,  $\{\phi(x_n, x_1)\}$  is non-decreasing. So, the limit also exists. From Lemma 2.7, we obtain that

$$\begin{aligned} \phi(x_{n+1}, x_n) &= \phi(x_{n+1}, \Pi_{C_n} x_1) \\ &\leq \phi(x_{n+1}, x_1) - \phi(\Pi_{C_n} x_1, x_1) \\ &= \phi(x_{n+1}, x_1) - \phi(x_n, x_1), \end{aligned} \tag{3.19}$$

thus, we have that

$$\lim_{n \rightarrow \infty} \phi(x_{n+1}, x_n) = 0. \tag{3.20}$$

Applying Lemma 2.8, we obtain that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \tag{3.21}$$

Suppose  $x_n = \Pi_{C_n} x_1 \subseteq C_m$ , for some positive integers  $m, n$  with  $m \leq n$ , then applying Lemma 2.7 and using the same approach as in (3.19), we obtain that

$$\begin{aligned} \phi(x_m, x_n) &= \phi(x_m, \Pi_{C_n} x_1) \\ &\leq \phi(x_m, x_1) - \phi(\Pi_{C_n} x_1, x_1) \\ &= \phi(x_m, x_1) - \phi(x_n, x_1). \end{aligned} \tag{3.22}$$

Since  $\lim_{n \rightarrow \infty} \phi(x_n, x_1)$  exists, it follows from (3.22) and Lemma 2.8 that  $\lim_{n \rightarrow \infty} \|x_n - x_m\| = 0$ . Hence, we conclude that  $\{x_n\}$  is a Cauchy sequence.

**Step 4:** Let  $\{x_n\}$  be a sequence generated by (3.1). Then we have the followings.

- (i)  $\lim_{n \rightarrow \infty} \|(P_Q - I)Ax_n\| = 0$ .
- (ii)  $\lim_{n \rightarrow \infty} \|S_i u_n - u_n\| = 0$ .
- (iii)  $\lim_{n \rightarrow \infty} \|y_n - w_n\| = 0$ .

Since  $x_{n+1} = \Pi_{C_{n+1}} x_1 \in C_{n+1} \subseteq C_n$ , by the definition of  $C_{n+1}$ , (3.18) and (3.20), we have that

$$\phi(x_{n+1}, t_n) \leq \phi(x_{n+1}, x_n) \rightarrow 0, \quad (n \rightarrow \infty). \tag{3.23}$$

It follows from Lemma 2.8 that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - t_n\| = 0. \quad (3.24)$$

Also, from (3.21) and (3.24), we have that

$$\lim_{n \rightarrow \infty} \|t_n - x_n\| = 0. \quad (3.25)$$

From (3.3), (3.12) and (3.15), we have that

$$\begin{aligned} \phi(x^*, t_n) &\leq \phi(x^*, x_n) - \alpha_n(1 - \alpha_n)g(\|J_1 u_n - S_i u_n\|) \\ &\quad - 2\gamma_n [ \|(P_Q - I)Ax_n\|^2 - k^2\gamma_n \|A^* J_2(P_Q - I)Ax_n\|^2 ] \\ &\quad - (1 - 2k^2\lambda_n^2 L^2 \mu)\phi(y_n, w_n). \end{aligned} \quad (3.26)$$

It then follows that

$$\begin{aligned} &\alpha_n(1 - \alpha_n)g(\|J_1 u_n - S_i u_n\|) \\ &\leq \phi(x^*, x_n) - \phi(x^*, t_n) \\ &= \|x^*\|^2 - 2\langle x^*, J_1 x_n \rangle + \|x_n\|^2 - \|x^*\|^2 + 2\langle x^*, J_1 t_n \rangle - \|t_n\|^2 \\ &= 2\langle x^*, J_1 t_n - J_1 x_n \rangle + \|x_n\|^2 - \|t_n\|^2 \\ &\leq 2\|x^*\| \|J_1 t_n - J_1 x_n\| + \|x_n - t_n\| (\|x_n\| + \|t_n\|). \end{aligned} \quad (3.27)$$

Since  $E_1$  is 2-uniformly convex and uniformly smooth Banach space,  $J_1$  is uniformly continuous from norm-to-norm. Then, we obtain from (3.25) that

$$\lim_{n \rightarrow \infty} \|J_1 t_n - J_1 x_n\| = 0. \quad (3.28)$$

By applying the condition  $\liminf_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0$  and (3.28) in (3.26), we obtain that

$$\lim_{n \rightarrow \infty} g(\|J_1 u_n - J_1 S_i u_n\|) = 0. \quad (3.29)$$

Using the property of  $g$  in Lemma 2.6, we have that

$$\lim_{n \rightarrow \infty} \|J_1 u_n - J_1 S_i u_n\| = 0. \quad (3.30)$$

Since  $J_1^{-1}$  is uniformly norm-to-norm continuous on bounded sets, we have

$$\lim_{n \rightarrow \infty} \|u_n - S_i u_n\| = 0. \quad (3.31)$$

Also, from (3.26) and following the same approach in (3.27), we have that

$$\phi(y_n, w_n) = 0. \quad (3.32)$$

Applying Lemma 2.8 in (3.32), we have

$$\lim_{n \rightarrow \infty} \|y_n - w_n\| = 0. \quad (3.33)$$

From (3.26), condition on  $\gamma_n$  in (3.1) and following the approach in (3.27), we get that

$$\lim_{n \rightarrow \infty} \|(P_Q - I)Ax_n\| = 0. \tag{3.34}$$

From (3.1), we have

$$J_1(t_n) - J_1(u_n) = \alpha_n(J_1S_i u_n - J_1(u_n)),$$

it implies that

$$\|J_1 t_n - J_1 u_n\| = \alpha_n \|J_1 S_i u_n - J_1 u_n\|.$$

Thus, from (3.30), we have that

$$\lim_{n \rightarrow \infty} \|J_1 t_n - J_1 u_n\| = 0. \tag{3.35}$$

Since  $E_1$  is 2-uniformly convex and uniformly smooth real Banach space and  $J_1^{-1}$  is uniformly norm-to-norm weakly continuous on bounded subset of  $E_1^*$ , we obtain that

$$\lim_{n \rightarrow \infty} \|t_n - u_n\| = 0. \tag{3.36}$$

From (3.24) and (3.36), we get that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - u_n\| = 0. \tag{3.37}$$

Also, from (3.25) and (3.36), we obtain that

$$\lim_{n \rightarrow \infty} \|u_n - x_n\| = 0. \tag{3.38}$$

More so, from (3.37) and (3.38), we get

$$\|u_{n+1} - u_n\| \leq \|u_{n+1} - x_{n+1}\| + \|x_{n+1} - u_n\| \rightarrow 0 \text{ as } n \rightarrow \infty. \tag{3.39}$$

We also have that

$$\begin{aligned} \|u_n - S_{i+l}u_n\| &\leq \|u_n - u_{n+l}\| + \|u_{n+l} - S_{i+l}u_{n+l}\| \\ &\quad + \|S_{i+l}u_{n+l} - S_{i+l}u_n\|, \end{aligned}$$

for all  $l = 1, 2, \dots, N$ . Using the assumption of  $S_l$ , we know that  $S_l$  is uniformly continuous. It then follows from (3.31) and (3.39) that

$$\lim_{n \rightarrow \infty} \|u_n - S_{i+l}u_n\| = 0, \forall l = 1, 2, \dots, N. \tag{3.40}$$

Thus, we have

$$\lim_{n \rightarrow \infty} \|u_n - S_l u_n\| = 0, \forall l = 1, 2, \dots, N. \tag{3.41}$$

Since  $S_l$  is closed for each  $l = 1, 2, \dots, N$  and  $\{x_n\} \rightharpoonup \bar{x}$ , we have that

$$\bar{x} \in \bigcap_{i=1}^N F(S_i).$$

**Step 5:** We show that  $\bar{x} \in (M + B)^{-1}(0)$ . Since  $\{x_n\}$  is bounded, there exists a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$  and  $\bar{x} \in E_1$  such that  $x_{n_k} \rightharpoonup \bar{x}$ . Suppose  $(v, u) \in \text{Gra}(M + B)$ . This implies that

$$J_1 u - Mv \in Bv.$$

More so, we obtain from (3.1) that

$$y_{n_k} = (J_1 + \lambda_{n_k} B)^{-1} J_1 \circ J_1^{-1} (J_1 w_{n_k} - \lambda_{n_k} M w_{n_k}),$$

which implies

$$(J_1 - \lambda_{n_k} M) w_{n_k} \in (J_1 + \lambda_{n_k} B) y_{n_k},$$

and thus

$$\frac{1}{\lambda_{n_k}} (J_1 w_{n_k} - J_1 y_{n_k} - \lambda_{n_k} M w_{n_k}) \in B y_{n_k}.$$

Using the fact that  $B$  is maximal monotone, we obtain

$$\langle v - y_{n_k}, J_1 u - Mv - \frac{1}{\lambda_{n_k}} (J_1 w_{n_k} - J_1 y_{n_k} - \lambda_{n_k} M w_{n_k}) \rangle \geq 0.$$

Hence, we have

$$\begin{aligned} \langle v - y_{n_k}, J_1 u \rangle &\geq \langle v - y_{n_k}, Mv + \frac{1}{\lambda_{n_k}} (J_1 w_{n_k} - J_1 y_{n_k} - \lambda_{n_k} M w_{n_k}) \rangle \\ &= \langle v - y_{n_k}, Mv - M w_{n_k} \rangle \\ &\quad + \langle v - y_{n_k}, \frac{1}{\lambda_{n_k}} (J_1 w_{n_k} - J_1 y_{n_k}) \rangle \\ &= \langle v - y_{n_k}, Mv - M y_{n_k} \rangle \\ &\quad + \langle v - y_{n_k}, M y_{n_k} - M w_{n_k} \rangle \\ &\quad + \langle v - y_{n_k}, \frac{1}{\lambda_{n_k}} (J_1 w_{n_k} - J_1 y_{n_k}) \rangle \\ &\geq \langle v - y_{n_k}, M y_{n_k} - M w_{n_k} \rangle \\ &\quad + \langle v - y_{n_k}, \frac{1}{\lambda_{n_k}} (J_1 w_{n_k} - J_1 y_{n_k}) \rangle. \end{aligned}$$

Applying (3.33) and using the fact that  $M$  is Lipschitz continuous, we obtain that

$$\lim_{n \rightarrow \infty} \|M y_{n_k} - M w_{n_k}\| = 0.$$

More so, we obtain that  $\langle v - \bar{x}, J_1 u \rangle \geq 0$ . By the maximal monotonicity of  $M + B$ , we obtain that  $0 \in (M + B)\bar{x}$ . Therefore, we conclude that

$$\bar{x} \in (M + B)^{-1}(0).$$



**Step 6:** We show that  $A\bar{x} \in Q$ . Using Lemma 2.7, we have that

$$\begin{aligned} \|(I - P_Q)A\bar{x}\|^2 &= \langle J_2(A\bar{x} - P_Q(A\bar{x})), A\bar{x} - P_Q(A\bar{x}) \rangle \\ &= \langle J_2(A\bar{x} - P_Q(A\bar{x})), A\bar{x} - Ax_n + Ax_n \\ &\quad - P_Q(Ax_n) + P_Q(Ax_n) - P_Q(A\bar{x}) \rangle \\ &= \langle J_2(A\bar{x} - P_Q(A\bar{x})), A\bar{x} - Ax_n \rangle \\ &\quad + \langle J_2(A\bar{x} - P_Q(A\bar{x})), Ax_n - P_Q(Ax_n) \rangle \\ &\quad + \langle J_2(A\bar{x} - P_Q(A\bar{x})), P_Q(Ax_n) - P_Q(A\bar{x}) \rangle \\ &\leq \langle J_2(A\bar{x} - P_Q(A\bar{x})), A\bar{x} - Ax_n \rangle \\ &\quad + \langle J_2(A\bar{x} - P_Q(A\bar{x})), Ax_n - P_Q(Ax_n) \rangle. \end{aligned} \tag{3.42}$$

Using the fact that  $A$  is a bounded linear operator and (3.34), we have that

$$\lim_{n \rightarrow \infty} \|Ax_n - A\bar{x}\| = 0,$$

this implies that  $\|(I - P_Q)A\bar{x}\| = 0$ , and hence

$$A\bar{x} \in Q.$$

From the Step 5 and Step 6, we conclude that  $\bar{x} \in \Gamma$ .

**Step 7:** We prove that  $\{x_n\}$  converges strongly to  $\bar{x}$ . Let  $\bar{x} = \Pi_\Gamma x_1$  and  $\bar{x} \in \Gamma$ . Then, from  $x_n = \Pi_{C_n} x_1$  and  $\bar{x} \in \Gamma \subseteq C_n$ , we have

$$\phi(x_n, x_1) \leq \phi(\bar{x}, x_1), \tag{3.43}$$

which implies that

$$\begin{aligned} \phi(\bar{x}, x_1) &\leq \liminf_{n \rightarrow \infty} \phi(x_n, x_1) \\ &\leq \phi(\bar{x}, x_1). \end{aligned} \tag{3.44}$$

From the definition of  $\bar{x} = \Pi_\Gamma x_1$ , we have that  $x^* = \bar{x}$ . Hence

$$\liminf_{n \rightarrow \infty} x_n = \bar{x} = \Pi_C x_1.$$

We therefore conclude that  $\{x_n\}$  converges strongly to  $\bar{x} \in \Gamma$ , where  $\bar{x} = \Pi_\Gamma x_1$ . This completes the proof.  $\square$

**Corollary 3.2.** *Suppose that  $E_1$ , and  $C$  be as defined in Theorem 3.1 and  $S : E_1 \rightarrow E_1$  be a nonexpansive mapping. Suppose that  $B : E_1 \rightarrow 2^{E_1^*}$  is a maximal monotone operator and  $M : E_1 \rightarrow E_1^*$  is monotone and  $L$ -Lipschitz continuous. Assume that*

$$\Gamma := \{\bar{x} \in C : \bar{x} \in F(S) \cap (B + M)^{-1}(0)\} \neq \emptyset.$$

*Let  $x_1 \in E_1$  and  $C = E_1$ , and  $\{x_n\}$  be a sequence generated by*

$$\begin{cases} y_n = J_{\lambda_n}^B \circ J_1^{-1}(J_1x_n - \lambda_n Mx_n); \\ u_n = J_1^{-1}(J_1y_n - \lambda_n(My_n - Mx_n)); \\ t_n = J_1^{-1}[(1 - \alpha_n)J_1u_n + \alpha_n J_1Su_n]; \\ C_{n+1} = \{v \in C_n : \phi(v, t_n) \leq \phi(v, x_n)\}; \\ x_{n+1} = \Pi_{C_{n+1}}x_1; \quad n \geq 1; \end{cases} \tag{3.45}$$

where  $\Pi_{C_{n+1}}$  is the generalized projection of  $E_1$  onto  $C_{n+1}$ . Suppose  $\{\alpha_n\}_{n=1}^\infty$  is a sequence in  $(0, 1)$  such that  $\liminf_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0$ , with  $\{\lambda_n\}_{n=1}^\infty$  satisfying the following condition:

$$0 < d \leq \lambda_n \leq e < \frac{1}{\sqrt{2\mu\rho L}},$$

where  $\mu$  is the 2-uniform convexity constant of  $E_1$ ,  $\rho$  is the 2-uniform smoothness constant of  $E_1^*$ , and  $L$  is the Lipschitz constant of  $M$ . Then the sequence  $\{x_n\}$  converges strongly to a point  $\bar{x} = \Pi_{\Gamma}x_1$ .

**Corollary 3.3.** *Let  $C$  and  $Q$  be nonempty, closed and convex subsets of real Hilbert spaces  $H_1$  and  $H_2$ , respectively. Let  $M : H_1 \rightarrow H_1$  be an  $\alpha$ -inverse strongly monotone operator with  $\alpha > 0$  and  $B : H_1 \rightarrow 2^{H_1}$  be a maximal monotone operator on  $H_1$  such that  $\text{Dom}(B)$  is included in  $H_1$ . Let  $\{S_n\} : H_1 \rightarrow H_1$  be a family of countable quasi-nonexpansive mappings which are uniformly closed, and  $A : H_1 \rightarrow H_2$  be a bounded linear operator with its adjoint  $A^*$ . Assume that*

$$\Gamma := \{\bar{x} \in C : \bar{x} \in F(S_n) \cap (M + B)^{-1}(0) \text{ and } A\bar{x} \in Q\} \neq \emptyset.$$

Let  $\{r_n\}$  be a positive real number sequence and  $\{\alpha_n\}$  be a real number sequence in  $[0, 1)$ . Let  $\{x_n\}$  be a sequence in  $C$  generated by

$$\begin{cases} x_1 \in C_1 = C, \text{ chosen arbitrarily}; \\ w_n = x_n + \gamma_n A^*(P_Q - I)Ax_n; \\ z_n = J_{r_n}(w_n - r_n Aw_n); \\ y_n = \alpha_n z_n + (1 - \alpha_n)S_n z_n; \\ C_{n+1} = \{z \in C_n : \|z_n - z\| \leq \|y_n - z\| \leq \|x_n - z\|\}; \\ x_{n+1} = P_{C_{n+1}}x_1, \quad n \geq 1; \end{cases}$$

where  $P_Q$  is the metric projection on  $H_2$ ,  $J_{r_n} = (I + r_n B)^{-1}$ ,  $\liminf_{n \rightarrow \infty} r_n > 0$ ,  $r_n \leq 2\alpha$ ,  $\limsup_{n \rightarrow \infty} \alpha_n < 1$ . and and the step size  $\gamma_n$  is chosen in such a way that for  $\varepsilon > 0$ ,

$$\gamma_n \in \left( \varepsilon, \frac{\|(P_Q - I)Ax_n\|^2}{\|A^*(P_Q - I)Ax_n\|^2} - \varepsilon \right),$$

for all  $P_Q Ax_n \neq Ax_n, \gamma_n = \gamma$  otherwise ( $\gamma$  being any nonnegative real number). Suppose  $\{\alpha_n\}_{n=1}^\infty$  is a sequence in  $(0, 1)$  such that  $\liminf_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0$ . Then the sequence  $\{x_n\}$  converges strongly to  $q = P_\Gamma x_1$ .

**Remark 3.4.** We observe that Corollary 3.3 coincide with the main result of [42]. Just that a new problem (split feasibility problem) was added to their iterative algorithm.

#### 4. APPLICATIONS

**1. Convexly Constrained Linear Inverse Problem:** Consider the convexly constrained linear inverse problem (see [13]) which is defined by

$$\begin{cases} Ax = b, \\ x \in C, \end{cases} \tag{4.1}$$

where  $H_1$  and  $H_2$  are two real Hilbert spaces,  $C, Q$  are closed convex subset of  $H_1$  and  $H_2$  respectively,  $A : H_1 \rightarrow H_2$  is a bounded linear operator and  $b \in Q$ . We denote by  $\Omega$  the solution set of (4.1).

Landweber introduced the following iterative algorithm to approximate the solution of (4.1) (see [14]) as follows:

$$\begin{cases} x_1 \in C, \\ x_{n+1} = P_C(x_n - \gamma A^*(Ax_n - b)), \quad n \geq 1, \end{cases}$$

where  $A^*$  is the adjoint of  $A$ ,  $0 < \gamma < 2\alpha$  with  $\alpha = \frac{1}{\|A\|^2}$ , then  $\{x_n\}$  converges weakly to a solution of (4.1).

Now, we introduce an iterative algorithm to approximate (4.1) and prove the following strong convergence result.

**Theorem 4.1.** *Let  $H_1$  and  $H_2$  be two real Hilbert spaces. Let  $C, Q$  be a nonempty, closed and convex subsets of a real Hilbert space  $H_1$  and  $H_2$  respectively. Let  $M : H_1 \rightarrow H_1$  be an  $\alpha$ -inverse strongly monotone operator with  $\alpha > 0$  and  $B : H_1 \rightarrow 2^{H_1}$  be a maximal monotone operator on  $H_1$  such that  $\text{Dom}(B)$  is included in  $H_1$ . Let  $\{S_n\} : H_1 \rightarrow H_1$  be a family of countable quasi-nonexpansive mappings which are uniformly closed, and  $A : H_1 \rightarrow H_2$  be a bounded linear operator with its adjoint  $A^*$ . Assume that*

$$\Gamma := \{\bar{x} \in C : \bar{x} \in F(S_n) \cap (M + B)^{-1}(0) \cap \Omega\} \neq \emptyset.$$

*Let  $\{r_n\}$  be a positive real number sequence and  $\{\alpha_n\}$  be a real number sequence in  $(0, 1)$ . Let  $\{x_n\}$  be a sequence in  $C$  generated by*

$$\begin{cases} x_1 \in C_1 = C, \text{ chosen arbitrarily;} \\ w_n = x_n - \gamma A^*(Ax_n - b); \\ z_n = J_{r_n}(w_n - r_n M w_n); \\ y_n = \alpha_n z_n + (1 - \alpha_n) S_n z_n; \\ C_{n+1} = \{z \in C_n : \|z_n - z\| \leq \|y_n - z\| \leq \|x_n - z\|\}; \\ x_{n+1} = P_{C_{n+1}} x_1, \quad n \geq 1; \end{cases}$$

where  $J_{r_n} = (I + r_n B)^{-1}$ ,  $\liminf_{n \rightarrow \infty} r_n > 0$ ,  $r_n \leq 2\alpha$ ,  $\limsup_{n \rightarrow \infty} \alpha_n < 1$ . and  $\gamma$  is a positive constant satisfying  $0 < \gamma < \frac{1}{\|A\|^2}$ . Suppose  $\{\alpha_n\}_{n=1}^{\infty}$  is a sequence in  $(0, 1)$  such that  $\liminf_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0$ . Then the sequence  $\{x_n\}$  converges strongly to  $q = P_{\Gamma} x_1$ .

*Proof.* This is a consequence of Corollary 3.3 by taking  $P_Q(Ax_n) = b$ .  $\square$

## 2. Minimization Problem:

**Definition 4.2.** Let  $Q$  be a convex subset of a vector space  $X$  and  $f : Q \rightarrow \mathbb{R} \cup \{+\infty\}$  be a map. Then,

(i)  $f$  is called convex if for each  $\lambda \in [0, 1]$  and  $x, y \in Q$ , we have

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y);$$

(ii)  $f$  is called proper if there exists at least one  $x \in Q$  such that

$$f(x) \neq +\infty;$$

(iii)  $f$  is called lower semi-continuous at  $x_0 \in Q$  if

$$f(x_0) \leq \liminf_{x \rightarrow x_0} f(x).$$

Let  $E$  be a real Banach space, we consider the following minimization of composite objective function of the type:

$$\min_{x \in E} f(x) + g(x), \quad (4.2)$$

where  $f : E \rightarrow \mathbb{R} \cup \{+\infty\}$  is a proper, convex and lower semi-continuous function and  $g : E \rightarrow \mathbb{R}$  is a convex function. In this setting, we assume that  $g$  is the smooth part of the functionals, while  $f$  is assumed to be non-smooth.

Precisely, we assume that  $g$  is Gâteaux-differentiable with derivative  $\nabla g$  which is Lipschitz continuous with constant  $L$ . It is easy to see from Theorem 3.13 ([28]) that

$$\langle \nabla g(x) - \nabla g(y), x - y \rangle \geq \frac{1}{L} \|\nabla g(x) - \nabla g(y)\|^2, \quad \forall x, y \in E.$$

Hence,  $\nabla g$  is monotone and Lipschitz continuous. It can be seen that (4.2) is equivalent to finding  $x \in E$  such that

$$0 \in \partial f(x) + \nabla g(x). \tag{4.3}$$

Problem (4.3) is a special case of (1.1) with  $M := \nabla g$  and  $B = \partial f$ .

We denote by  $\Omega$  the solution set of (4.3). Also, for fixed  $r > 0$  and  $z \in E$ , it has been shown in [31] that the resolvent of  $\partial f$  which is denoted as  $J_r^{\partial f}$  is defined as

$$J_r^{\partial f}(z) = \arg \min_{y \in E} \left\{ f(y) + \frac{1}{2r} \|y\|^2 - \frac{1}{r} \langle y, Jz \rangle \right\}.$$

This can be re-written using (3.1) as

$$y_n = \arg \min_{y \in E} \left\{ f(y) + \frac{1}{2\lambda_n} \|y\|^2 - \frac{1}{\lambda_n} \langle y, Jw_n - \lambda_n \nabla g(w_n) \rangle \right\}.$$

**Theorem 4.3.** *Let  $E_1, E_2, A, A^*$ , and  $Q$  be as defined in Theorem 3.1 and suppose that*

$$\Gamma := \{ \bar{x} \in C : \bar{x} \in \cap_{i=1}^N F(S_i) \cap \Omega : A\bar{x} \in Q \} \neq \emptyset.$$

Let  $x_1 \in E_1$  and  $C_1 = E_1$ , and  $\{x_n\}$  be a sequence generated by

$$\begin{cases} w_n = J_1^{-1}(J_1 x_n + \gamma_n A^* J_2(P_Q - I)Ax_n); \\ y_n = \operatorname{argmin}_{y \in E} \left\{ f(y) + \frac{1}{2\lambda_n} \|y\|^2 - \frac{1}{\lambda_n} \langle y, J_1 w_n - \lambda_n \nabla g(w_n) \rangle \right\}; \\ u_n = J_1^{-1}(J_1 y_n - \lambda_n (\nabla g(y_n) - \nabla g(x_n))); \\ t_n = J_1^{-1}[(1 - \alpha_n)J_1 u_n + \alpha_n J_1 S_i u_n]; \\ C_{n+1} = \{v \in C_n : \phi(v, t_n) \leq \phi(v, x_n)\}; \\ x_{n+1} = \Pi_{C_{n+1}} x_1; \quad n \geq 1; \end{cases} \tag{4.4}$$

where  $P_Q$  is the metric projection of  $E_2$  onto  $Q$  and  $\Pi_{C_{n+1}}$  is the generalized projection of  $E_1$  onto  $C_{n+1}$ . Suppose  $\{\alpha_n\}_{n=1}^\infty$  is a sequence in  $(0, 1)$  such that  $\liminf_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0$ , and the step size  $\gamma_n$  is chosen in such a way that for  $\varepsilon > 0$ ,

$$\gamma_n \in \left( \varepsilon, \frac{\|(P_Q - I)Ax_n\|^2}{\|A^*k^2 J_2(P_Q - I)Ax_n\|^2} - \varepsilon \right),$$

for all  $P_Q Ax_n \neq Ax_n$ ,  $\gamma_n = \gamma$  otherwise ( $\gamma$  being any nonnegative real number) with  $\{\lambda_n\}_{n=1}^\infty$  satisfying the following condition:

$$0 < d \leq \lambda_n \leq e < \frac{1}{\sqrt{2\mu\rho L}},$$

where  $\mu$  is the 2-uniform convexity constant of  $E_1$ ,  $\rho$  is the 2-uniform smoothness constant of  $E_1^*$ , and  $L$  is the Lipschitz constant of  $\nabla g$ . Then,  $\{x_n\}$  converges strongly to a point  $\bar{x} = \Pi_\Gamma x_1$ .

**Acknowledgments:** The first and second authors acknowledge with thanks the bursary and financial support from Department of Science and Technology and National Research Foundation, Republic of South Africa, Centre of Excellence in Mathematical and Statistical Sciences (DST-NRF COE-MaSS) Post Doctoral Bursary. Opinions expressed and conclusions arrived are those of the authors and are not necessarily to be attributed to the CoE-MaSS. And the fourth author was supported by the Basic Science Research Program through the National Research Foundation(NRF) Grant funded by Ministry of Education of the republic of Korea (2018R1D1A1B07045427).

## REFERENCES

- [1] H.A. Abass, F.U. Ogbuisi and O.T. Mewomo, *Common solution of split equilibrium problem with no prior knowledge of operator norm*, UPB Sci. Bull., Series A, **80** (2018), 175–190.
- [2] H.A. Abass, C. Izuchukwu, O.T. Mewomon and Q.L. Dong, *Strong convergence of an inertial forward-backward splitting method for accretive operators in real Banach space*, Fixed Point Theory, **21** (2020), 397–412.
- [3] H.A. Abass, K.O. Aremu, L.O. Jolaoso and O.T. Mewomo, *An inertial forward-backward splitting method for approximating solutions of certain optimization problem*, J. Nonlinear Funct. Anal., **2020** (2020), Article ID 6.
- [4] Y.I. Alber, *Metric and generalized projection operators in Banach spaces: properties and applications* in: Kartsatos, A.G (Ed). Theory and Applications of Nonlinear Operators and Accretive and Monotone Type. Lecture Notes Pure Appl. Math., **178**, Dekker, New York (1996), 15–50.
- [5] Y.I. Alber and S. Reich, *An iterative method for solving a class of nonlinear operator equations in Banach spaces*, PanAmer. Math. J., **4** (1994), 39–54.
- [6] Y. Alber and L. Ryazantseva, *Nonlinear ill-posed problems of monotone type*, Springer, Dordrecht (2006). xiv+410 pp. ISBN:978-1-4020-4395-6, 1-4020-4395-3.
- [7] K. Aoyama and F. Koshaka, *Strongly relatively nonexpansive sequences generated by firmly nonexpansive-like mappings*, Fixed Point Theory Appl., **95** (2014), pp.13.
- [8] K. Avetisyan, O. Djordjevic and M. Pavlovic, *Littlewood-Paley inequalities in uniformly convex and uniformly smooth Banach spaces*, J. Math. Anal. Appl., **336** (2007), 31–43.
- [9] V. Barbu, *Nonlinear semigroups and differential equations in Banach spaces*, Editura Academiei, R.S.R, Bucharest, English transl. Noordhof, Leyden, 1976.
- [10] Y. Censor and T. Elfving, *A multiprojection algorithm using Bregman projection in a product space*, Numer. Algor., **8** (1994), 221–239.
- [11] Y. Censor, T. Elfving, N. Kopf and T. Bortfield, *The multiple-sets split feasibility problem and its applications for inverse problems*, Inverse Prob., **21** (2005), 2071–2084.
- [12] Q.L. Dong, D. Jiang, P. Cholmjiak and Y. Shehu, *A strong convergence result involving an inertial forward-backward splitting algorithm for monotone inclusions*, J. Fixed Theory Appl., **19** (2017), 3097–3118.
- [13] B. Eicke, *Iteration methods for convexly constrained ill-posed problems in Hilbert space*, Numer. Funct. Anal. Optim., **13** (1992), 413–429.
- [14] H.W. Engl, M. Hanke and A. Neubauer, *Regularization of inverse problems*, Kluwer Academic Publishers Group, Dordrecht, 1996.

- [15] J.N. Ezeora, H.A. Abass and C. Izuchukwu, *Strong convergence of an inertial-type algorithm to a common solution of minimization and fixed point problems*, *Mathematik Vesnik*, **71** (2019), 338–350.
- [16] Z. Jouymandi and F. Moradlou, *Retraction algorithms for solving variational inequalities, pseudomonotone equilibrium problems and fixed point problems in Banach spaces*, *Numer. Algor.*, **78** (2018), 1153–1182.
- [17] S. Kamimura and W. Takahashi, *Strong convergence of a proximal-type algorithm in Banach space*, *SIAM J. Optim.*, **13** (2002), 938–945.
- [18] J.K. Kim, A.H. Dar and Salahuddin, *Existence theorems for the generalized relaxed pseudomonotone variational inequalities*, *Nonlinear Funct. Anal. and Appl.*, **25**(1) (2020), 25–34. doi.org/10.22771/nfaa.2020.25.01.03
- [19] J.K. Kim, T.M. Tuyen and M.T. Ngoc Ha, *Two projection methods for solving the split common fixed point problem with multiple output sets in Hilbert spaces*, *Numer. Funct. Anal. Optimiz.*, **42**(8) (2021), 973–988, <https://doi.org/10.1080/01630563.2021.1933528>
- [20] J.K. Kim and T.M. Tuyen, *A parallel iterative method for a finite family of Bregman strongly nonexpansive mappings in reflexive Banach spaces*, *J. Korean Math. Soc.*, **57**(3) (2020), 617–640. <https://doi.org/10.4134/JKMS.j190268>
- [21] J.K. Kim and T.M. Tuyen, *Parallel iterative method for solving the split common null point problem in Banach spaces*, *J. Nonlinear and Convex Anal.*, **20**(10) (2019), 2075–2093.
- [22] P.L. Lions and B. Mercier, *Splitting algorithms for the sum of two nonlinear operators*, *SIAM J. Numer. Anal.*, **16** (1979), 964–979.
- [23] Z. Ma, L. Wang and S.S. Chang, *On the split feasibility problem and fixed point problem of quasi- $\phi$ -nonexpansive in Banach spaces*, *Numer. Algor.*, **80**(4) (2019), 1203–1218. <https://doi.org/10.1007/s11075-018-0523-1>.
- [24] S. Matsushita and W. Takahashi, *A strong convergence theorem for relatively nonexpansive mappings in Banach spaces*, *J. Approx. Theory*. **134** (2005), 257–266.
- [25] A.A. Mebawondu, *Proximal Point Algorithms for Finding Common Fixed Points of a Finite Family of Nonexpansive Multivalued Mappings in Real Hilbert Spaces*, *Khayyam Journal of Mathematics*, **5**(2), 113–123.
- [26] A. Moudafi and B.S. Thakur, *Solving proximal split feasibility problems without prior knowledge of operator norms*, *Optim. Letter*, **8** (2014), 2099–2110.
- [27] D.H. Peaceman and H.H. Rashford, *The numerical solution of parabolic and elliptic differential equations*, *J. Soc. Ind. Appl. Math.*, **3** (1995), 267–275.
- [28] J. Peypouquet, *Convex optimization in Normed spaces. Theory, Methods and Examples*. With a foreword by H. Attouch. Springer briefs in optimization. Springer, Cham., 2015. xiv+124 pp. ISBN: 978-3-319-13709-4, 978-3-319-13710-0.
- [29] X. Qin, Y.J. Cho and S.M. Kang, *Convergence theorems of common elements for equilibrium problem and fixed point problems in Banach spaces*, *J. Comput. Appl. Math.*, **225** (2009), 20–30.
- [30] R.T. Rockafellar, *Monotone operators and the proximal point algorithm*, *SIAM J. Control Optim.*, **14** (1977), 877–808.
- [31] Y. Shehu, *Convergence results of forward-backward algorithms for sum of monotone operators in Banach spaces*, *Results Math.*, **74** (2019), 138.
- [32] Y. Shehu, *Iterative approximation for zeros of sum of accretive operators in Banach spaces*, *J. Funct. Spaces*, (2015), Article ID 5973468, 9 pages.
- [33] Y. Shehu, *Iterative approximation method for finite family of relatively quasi-nonexpansive mapping and systems of equilibrium problem*, *J. Glob. Optim.*, (2014), DOI.10.1007/s10898-010-9619-4.

- [34] Y. Shehu and O.S. Iyiola, *Convergence analysis for the proximal split feasibility using an inertial extrapolation term method*, J. Fixed Point Theory Appl., **19** (2017), 2483–2510.
- [35] W. Takahashi, *Nonlinear Functional Analysis*, Fixed Theory Appl., Yokohama-Publishers, 2000.
- [36] Ng.T.T. Thuy, P.T.T. Hoai and Ng.T.T. Hoa, *Explicit iterative methods for maximal monotone operators in Hilbert spaces*, Nonlinear Funct. Anal. Appl., **25**(4) (2020), 753–767. doi.org/10.22771/nfaa.2020.25.04.09.
- [37] P.T. Vuong, J.J. Stroduot and V.H. Nguyen, *A gradient projection method for solving split equality and split feasibility problems in Hilbert space*, Optimization, **64** (2015), 2321–2341.
- [38] K. Wattanawitoon and P. Kuman, *Strong convergence theorems by a new hybrid projection algorithm for fixed point problem and equilibrium problems of two relatively quasi-nonexpansive mappings*, Nonlinear Anal. Hybrid Syst., **3** (2009), 11–20.
- [39] H.K. Xu, *Inequalities in Banach spaces with applications*, Nonlinear Anal. Theory Methods Appl., **16** (1991), 1127–1138.
- [40] J.C. Yao, *Variational inequalities with generalized monotone operators*, Math. Oper. Res., **19** (1994), 691–705.
- [41] H. Zhang and L. Ceng, *Projection splitting methods for sums of maximal monotone operators with applications*, J. Math. Anal. Appl., **406** (2013), 323–334.
- [42] J. Zhang and N. Jiang, *Hybrid algorithm for common solution of monotone inclusion problem and fixed point problem and applications to variational inequalities*, Springer Plus, **5**: 803 (2016). <https://doi.org/10.1186/s40064-016-2389-9>