

Original Research



Reproducibility of a food frequency questionnaire: Korea Nurses' Health Study

Sihan Song ¹, Bohye Kim ², Yanghee Pang ², Oksoo Kim ², Jung Eun Lee ^{1,3*}

¹Department of Food and Nutrition, College of Human Ecology, Seoul National University, Seoul 08826, Korea

²College of Nursing, Ewha Womans University, Seoul 03760, Korea

³Research Institute of Human Ecology, Seoul National University, Seoul 08826, Korea



Received: Oct 27, 2020

Revised: Mar 26, 2021

Accepted: Jul 5, 2021

Published online: Jul 12, 2021

*Corresponding Author:


Jung Eun Lee


Department of Food and Nutrition, College of Human Ecology, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Korea.
Tel. 82-2-880-6834
Fax. 82-2-884-0305
Email. jungelee@snu.ac.kr


©2022 The Korean Nutrition Society and the Korean Society of Community Nutrition
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.


ORCID iDs

Sihan Song 
<https://orcid.org/0000-0003-1634-8004>

Bohye Kim 
<https://orcid.org/0000-0001-5419-3467>

Yanghee Pang 
<https://orcid.org/0000-0003-3768-156X>

Oksoo Kim 
<https://orcid.org/0000-0001-9071-6093>

Jung Eun Lee 
<https://orcid.org/0000-0003-1141-878X>

Funding

This work was supported by the Research Program funded by the Korea National Institute of Health (2013-E63006-00, 2013-E63006-01, 2019-ER710-100, 2019-ER710-102).

<https://e-nrp.org>

ABSTRACT

BACKGROUND/OBJECTIVES: This study aimed to examine the reproducibility of food frequency questionnaires (FFQs) designed for young female nurses in the Korea Nurses' Health Study.

SUBJECTS/METHODS: The reproducibility of web-based, self-administered FFQs was evaluated among 243 Korean female nurses. The first FFQ (FFQ1) was administered from March 2014 to February 2019 and the second FFQ (FFQ2) from November 2019, with a mean interval of 2.8 years between the FFQs (range, 9 months–5.6 years). Pearson and Spearman correlation coefficients (r values) and quartile agreements between FFQ1 and FFQ2 were calculated for intakes of energy, nutrients, and foods.

RESULTS: Pearson correlation coefficients ranged from 0.41 to 0.55 (median r = 0.51) for energy and raw nutrients and from 0.16 to 0.46 (median r = 0.36) for energy-adjusted nutrients. Spearman correlation coefficients ranged from 0.25 to 0.72 (median r = 0.41) for food items. The percentages of women who were classified into the same or adjacent quartile were 77% to 84% (median = 82%) for raw nutrients and 69% to 86% (median = 78%) for foods.

CONCLUSIONS: The results indicated that the web-based FFQ used in the Korea Nurses' Health Study has acceptable reproducibility.

Keywords: Food frequency questionnaire; reproducibility; Korea Nurses' Health Study

INTRODUCTION

Non-communicable diseases have emerged as a new global agenda for women's health [1,2]. A growing body of evidence from the long-term cohort studies has contributed to understand and improve the health of women [3-6]. Diet is one of the most important determinants of health throughout the life and across generations [7,8]. To ensure the research findings regarding the association between diet and disease outcome, an accurate assessment of dietary exposure is essential [9]. In the Nurses' Health Study (NHS), a large cohort study of female nurses in the United States (US), various dietary factors assessed by validated questionnaires have been identified as determinants of chronic diseases in women [10]. For example, the NHS found that high consumption of red meat or sugar-sweetened beverages and low consumption of dietary fiber or fruits and vegetables were associated with increased risk of coronary heart disease [11].

Conflict of Interest

The authors declare no potential conflicts of interests.

Author Contributions

Conceptualization: Lee JE; Formal analysis: Song S; Funding acquisition: Kim O; Investigation: Kim B, Pang Y, Kim O; Methodology: Song S, Lee JE; Supervision: Kim O, Lee JE; Writing - original draft: Song S; Writing - review & editing: Kim B, Pang Y, Kim O, Lee JE.

The Korea Nurses' Health Study (KNHS), which began in 2013, is a prospective study of young female nurses in Korea [12]. The purpose of the KNHS is to investigate the effects of occupational, environmental, and lifestyle factors on women's health. This long-term follow-up study of women of reproductive age is expected to contribute to framing of evidence-based policies and guidelines for women's health. In the KNHS, dietary information was assessed using the web-based food frequency questionnaire (FFQ) [12]. Given that FFQs are designed to evaluate the long-term dietary intake of the study population, the capacity of the FFQ to measure the participants' diet over an extended period, despite the within-person variation, should be evaluated. The FFQ is typically validated by comparing the reference methods, and its reproducibility is generally assessed by administering the same FFQ twice to the same group of participants. The FFQ used in the KNHS was initially developed for adults aged 19–64 years in the Korean National Health and Nutrition Examination Survey (KNHANES) [13]. It has been reported that the FFQ used in this study has acceptable reproducibility and modest validity [14]; significant associations between dietary factors, as assessed by this FFQ, and disease-related outcomes [15–19] suggest that this FFQ appropriately captures diet-disease associations.

However, the performance of the FFQ in the KNHS has not been addressed. Therefore, the present study aimed to evaluate the reproducibility of the web-based FFQ used in the KNHS in women aged 20–45 years in Korea.

SUBJECTS AND METHODS

Participants and study design

The KNHS is an ongoing web-based prospective cohort study of female registered nurses aged 20–45 years that began in 2013 in Korea [12]. The protocol and questionnaires of the KNHS are based on the NHS3, an ongoing web-based prospective study in the US and Canada that began in 2010 [20]. In the first phase of the KNHS (March 2013–December 2015), survey modules 1–4 and pregnancy modules were open to participants. The initial baseline survey (module 1) was conducted from July 2013 to November 2014, and a total of 20,613 women were enrolled. Participants who responded to the module 1 survey were invited through a text message to continue with subsequent online surveys. The second phase of the study was conducted from June 2016 to March 2019, in which participants provided answers to survey modules 5–7. The third phase began in March 2019 with module 8, and module 9 in the third phase was administered beginning October 2020. Some questions such as job status, disease history, and reproductive factors have been repeated in the modules. The KNHS has been described in detail elsewhere [12].

In the baseline survey module 1, participants were asked about demographic characteristics, weight and height, disease history, reproductive factors, employment, occupational exposure, and lifestyle factors, including alcohol consumption and smoking status. In the subsequent survey module 2, usual dietary intake during the past year was assessed using the web-based FFQ. For the current analysis, we used dietary data collected from May 2014 to February 2019. Among the 15,355 participants who agreed to participate in module 2, a total of 14,485 participants completed the FFQ. We aimed to recruit 300 participants in November 2019 to examine the reproducibility of the FFQ. In epidemiological studies, a sample size of about 100 to 200 participants is reasonable to evaluate the performance of the FFQ [9].

Among those who had completed the first FFQ (FFQ1), we excluded women who were pregnant at the time of dietary assessment ($n = 684$), had been diagnosed with a chronic disease, including type 2 diabetes, hypertension, cardiovascular diseases, or cancer ($n = 743$), or had a short interval (< 9 months) from the date of FFQ1 completion to November 2019 ($n = 3$). We further excluded 875 participants who reported an implausible energy intake (< 500 kcal/day, > 95 th percentile of energy intake). Consequently, 12,218 participants were eligible for re-contact, and 900 individuals were selected as potential participants for this study as follows. We initially included potential participants who had completed the FFQ1 within 9 months to less than 1 year ($n = 151$) and selected the rest of the participants using stratified random sampling ($n = 749$). In other words, we oversampled participants who had recently completed the FFQ1 to minimize the effect of dietary change over time. Participants who had completed the FFQ1 ≥ 1 year prior were stratified into 3 groups by time since the completion of the FFQ1 (1 to < 2.5 years, 2.5 to < 4 years, and ≥ 4 years) and were randomly selected in a ratio of 3:2:1 using PROC SURVEYSELECT in SAS (SAS Institute Inc., Cary, NC, USA).

The questionnaire comprised the second FFQ (FFQ2) and questions on reproductive factors (pregnancy, childbirth, and breastfeeding), history of chronic diseases (type 2 diabetes, hypertension, cardiovascular diseases, and cancer), and occupational status (current job status and shift work information). The invitation to participate in the survey was sent to potential participants via text message that included a link to the questionnaire. We sent a reminder message to non-responders 3 days after sending the web-based questionnaire. The online survey was closed when the number of respondents reached 300. A coffee coupon was given to participants as an incentive. Among participants enrolled from November 5 to November 20, 2019 ($n = 300$), we excluded participants who were pregnant, had given birth, or had breastfed an infant in the last year ($n = 45$), had been diagnosed with a chronic disease ($n = 15$), or reported an implausible energy intake ($n = 4$). Consequently, a total of 243 women were included in this analysis. The study was approved by the Institutional Review Board of the Ewha Womans University (ewha-201904-0012-05). All participants provided the informed consent after understanding the purpose of the study.

The FFQ

The semi-quantitative FFQ was developed to assess the usual dietary intake over the past year in Korean adults [13]. The validity of the FFQ among 126 Korean adults have been described in detail elsewhere [14]. In brief, the validity of the FFQ was assessed by comparing energy as well as intake of thirteen nutrients between the FFQ findings and 4 seasonal 3-day dietary records (DR). The correlation coefficients between the first FFQ and 12-day DR ranged from 0.27 to 0.45 (mean $r = 0.38$) for energy and nutrients and from 0.15 to 0.64 (mean $r = 0.39$) for energy-adjusted and de-attenuated nutrients [14]. The FFQ includes 112 food items commonly consumed by Korean adults. Participants were asked to select their average frequency of consumption over the past year by selecting one of nine frequency categories (never or seldom, once per month, 2–3 times per month, once per week, 2–4 times per week, 5–6 times per week, once per day, twice per day, or three times per day). Participants were also asked to select their usual portion size as small, medium, or large. In the KNHS, figures for the portion sizes of food items were provided to aid the estimation of portion sizes. To calculate the daily intake, the selected frequency category for each food item was converted to a daily frequency of consumption. Daily intakes of energy, nutrients, and foods were calculated by multiplying the frequency of consumption by the nutrient content of the selected portion size. The nutrient database for the FFQ in the KNHANES was primarily based on the food composition table from the Korean National

Academy of Agricultural Science and that provided by the Korea Disease Control and Prevention Agency [21].

Statistical analysis

We compared the baseline characteristics of 300 participants who were administered FFQ2 with those of all study participants of the KNHS ($n = 20,613$). Descriptive statistics were presented as mean and SD for continuous variables, and as number and proportion (%) for categorical variables. Among the 243 subjects who were included in the reproducibility analysis, group means and SDs were separately calculated from FFQ1 and FFQ2 for daily intake of energy, % energy from macronutrients (protein, fat, and carbohydrate), and twenty-two nutrients. We used a paired t-test to examine the differences in energy and nutrient intakes between the FFQ1 and FFQ2.

To assess the reproducibility, we estimated the relative ranking of participants by the test–retest responses. We calculated their Pearson's correlation coefficients for energy and nutrients and Spearman's correlation coefficients for food items. We also calculated their energy-adjusted nutrient intake using a residual method to estimate energy-adjusted Pearson's correlation coefficient [9]. To improve the normality of data, energy, % energy from macronutrients, and nutrient intake values were log-transformed. We also calculated the agreement in quartile classification of the dietary intake between FFQ1 and FFQ2. The percentages of participants classified into the same quartile, the same or adjacent quartile, and the opposite quartile were calculated. A few food items were not divided into quartiles because of skewed distribution and therefore we did not compute % quartile agreement. Because a longer time interval between FFQs decreases the reproducibility due to true changes in diet as well as variations in response [9], we also evaluated the mean daily intakes and reproducibility of FFQs administered in a relatively short interval. A time interval of < 3 years was chosen to perform analysis with the sample size of more than 100 ($n = 139$).

All statistical analyses were performed using SAS statistical software version 9.4 (SAS Institute Inc.). All hypothesis tests were evaluated using 2-tailed tests of significance at P -value < 0.05.

RESULTS

The baseline characteristics of the current reproducibility study respondents did not differ from those of all cohort participants in terms of age, body mass index, alcohol consumption, smoking status, education levels, or marital status (**Table 1**). Among the 243 women included in this study, the mean time interval between FFQ1 and FFQ2 was 2.76 years and ranged from 9.24 months to 5.64 years. In addition, the mean time interval between FFQs was 1.74 years among 139 women who completed FFQ2 within the 3 years since FFQ1 administration.

Mean daily intakes, correlation coefficients, and quartile agreement between FFQs are shown in **Table 2**. Mean daily intakes were lower in FFQ2 than in FFQ1, except for % energy from protein and fat, carotene, and vitamin C intakes. Pearson's correlation coefficient between FFQ1 and FFQ2 was 0.46 for energy intake. The correlation coefficients between FFQ1 and FFQ2 ranged from 0.41 for carbohydrate to 0.55 for protein or iron (median $r = 0.51$) for nutrient intake. Adjustment for energy intake attenuated the correlation coefficients, except for the correlation coefficient for carbohydrate. The correlations ranged from 0.16 for

carotene to 0.46 for carbohydrate (median $r = 0.36$) for energy-adjusted nutrient intake. The correlation coefficients for % energy from macronutrients were similar to those of energy-adjusted macronutrients using the residual methods. The median percentages of the same

Table 1. Characteristics of the respondents of the reproducibility study and the cohort participants

Characteristics ¹⁾	Reproducibility study respondents (n = 300)	KNHS participants (n = 20,613)
Age (yr)	29.63 ± 5.86	29.38 ± 5.92
BMI ²⁾ (kg/m ²)	20.90 ± 2.54	20.95 ± 2.72
Alcohol consumption (cups/day)	0.36 ± 0.73	0.32 ± 0.69
Smoking status		
Never	289 (96.33)	19,938 (96.74)
Ever	11 (3.67)	671 (3.26)
Education levels		
3-year college	142 (47.33)	9,771 (47.40)
4-year university	139 (46.33)	9,316 (45.20)
Master's or higher	19 (6.33)	1,525 (7.40)
Marital status		
Unmarried	187 (62.33)	13,548 (65.73)
Married	110 (36.67)	6,965 (33.79)
Others (cohabiting/divorced/separated/widowed)	3 (1.00)	98 (0.48)

Values are presented as mean ± SD or number (%).

KNHS, Korea Nurses' Health Study; BMI, body mass index.

¹⁾Number may not total to 300 or 20,613 for each characteristic due to missing values. Data are missing for BMI (n = 4 among reproducibility study respondents; n = 93 among overall participants) and alcohol consumption (n = 2 among reproducibility study respondents; n = 31 among overall participants).

²⁾BMI was calculated as weight divided by height squared (kg/m²).

Table 2. Mean daily intakes and reproducibility of energy and nutrients between FFQ1 and FFQ2 (n = 243)

Daily intake	Mean ± SD ¹⁾		Pearson's correlation ²⁾		Quartile agreement		
	FFQ1	FFQ2	Crude r	Energy-adjusted r	Same	Same or adjacent	Opposite
Energy (kcal)	2,253.10 ± 1,029.69	1,845.85 ± 923.29	0.46	-	41	79	4
% Energy from protein	14.19 ± 2.62	14.46 ± 2.70*	-	0.40	37	77	7
% Energy from fat	22.25 ± 5.99	23.54 ± 5.85	-	0.36	35	75	6
% Energy from carbohydrate	60.81 ± 8.66	59.42 ± 8.33	-	0.46	39	80	5
Protein (g)	81.66 ± 47.34	66.53 ± 37.44	0.55	0.40	42	84	4
Fat (g)	57.35 ± 36.09	47.41 ± 26.25	0.52	0.36	43	82	4
SFA (g)	17.45 ± 11.36	14.26 ± 8.07	0.51	0.42	40	82	4
MUFA (g)	18.35 ± 12.23	15.11 ± 8.67	0.53	0.39	41	83	4
PUFA (g)	13.54 ± 8.00	11.64 ± 6.55	0.51	0.34	42	79	4
n-3 FA (g)	1.51 ± 0.94	1.30 ± 0.77	0.53	0.32	44	79	4
n-6 FA (g)	12.19 ± 7.22	10.49 ± 5.88	0.51	0.35	43	79	4
Cholesterol (mg)	341.96 ± 220.83	303.55 ± 200.46	0.51	0.31	43	84	2
Carbohydrate (g)	337.63 ± 145.37	275.85 ± 143.91	0.41	0.46	39	77	4
Dietary fiber (g)	20.45 ± 12.27	16.93 ± 11.09	0.53	0.35	41	81	4
Calcium (mg)	520.32 ± 295.29	423.77 ± 250.21	0.50	0.40	44	82	4
Phosphorus (mg)	1,121.37 ± 573.78	926.19 ± 501.21	0.53	0.39	40	81	4
Iron (mg)	14.63 ± 7.87	12.13 ± 7.13	0.55	0.39	42	83	5
Sodium (mg)	3,836.82 ± 2,445.90	3,079.85 ± 1,986.50	0.50	0.25	47	84	4
Potassium (mg)	2,908.51 ± 1,611.46	2,474.58 ± 1,522.71	0.53	0.38	40	83	3
Vitamin A (µg RE)	636.42 ± 385.50	562.48 ± 390.81	0.47	0.21	47	79	4
Retinol (µg)	121.48 ± 77.70	102.42 ± 63.71	0.50	0.42	38	83	3
Carotene (µg)	2,870.83 ± 1,936.34	2,597.78 ± 2,021.09*	0.43	0.16	43	79	2
Thiamin (mg)	2.03 ± 1.04	1.67 ± 0.88	0.50	0.38	39	80	5
Riboflavin (mg)	1.60 ± 0.88	1.36 ± 0.75	0.53	0.36	40	84	2
Niacin (mg)	15.70 ± 9.12	12.64 ± 7.39	0.51	0.34	47	82	6
Vitamin C (mg)	106.09 ± 112.61	92.98 ± 89.11*	0.49	0.36	39	78	3

FFQ1, the first food frequency questionnaire; FFQ2, the second food frequency questionnaire; SFA, saturated fatty acids; MUFA, monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; FA, fatty acids.

¹⁾Non-significant differences in mean intakes at $\alpha = 0.05$ between FFQs were noted as “**”.

²⁾All correlation coefficients were statistically significant at $\alpha = 0.05$.

quartile, the same or adjacent quartile, and the opposite quartile classification for energy and nutrients were 42%, 82%, and 4%, respectively.

The same sets of analyses for energy and nutrients were conducted among women who completed FFQ2 during the period from 9 months to less than 3 years since they had completed FFQ1 (**Table 3**). Mean daily intakes were generally lower in FFQ2 than in FFQ1; however, there were no statistically significant differences between the 2 FFQs in terms of fat (fat subtypes and cholesterol), vitamin A, retinol, carotene, and vitamin C intakes. The reproducibility was slightly increased among women who had a shorter time interval. Pearson's correlation coefficient between FFQ1 and FFQ2 for energy intake was 0.49. For nutrient intake, correlation coefficients between FFQ1 and FFQ2 ranged from 0.42 for carbohydrate to 0.62 for potassium (median $r = 0.57$) for raw nutrients and from 0.11 for carotene to 0.49 for potassium or retinol (median $r = 0.39$) for energy-adjusted nutrients. The median percentages of the same quartile, the same or adjacent quartile, and the opposite quartile classification for energy and nutrients were 45%, 82%, and 4%, respectively.

Spearman's correlation coefficients and quartile agreement of 40 food items according to the time interval between FFQs are shown in **Table 4**. The correlation coefficients ranged from 0.25 for cooked rice to 0.72 for beer (median $r = 0.41$) among the 243 women and from 0.24 for kimchi stew/stir-fried kimchi to 0.74 for beer (median $r = 0.45$) among the 139 women who completed FFQ2 within < 3 years since FFQ1. The median percentages of the same quartile, the same or adjacent quartile, and the opposite quartile classification were

Table 3. Mean daily intakes and reproducibility of energy and nutrients between FFQ1 and FFQ2, < 3 years apart (n = 139)

Daily intake	Mean \pm SD ¹⁾		Pearson's correlation ²⁾		Quartile agreement		
	FFQ1	FFQ2	Crude r	Energy-adjusted r	Same	Same or adjacent	Opposite
Energy (kcal)	2,110.76 \pm 984.91	1,859.44 \pm 951.83	0.49	-	45	80	6
% Energy from protein	14.05 \pm 2.65	14.58 \pm 2.83	-	0.43	38	79	5
% Energy from fat	22.34 \pm 6.31	24.22 \pm 6.31	-	0.39	40	76	5
% Energy from carbohydrate	60.73 \pm 8.94	58.76 \pm 8.35	-	0.44	37	78	6
Protein (g)	75.14 \pm 44.08	67.69 \pm 40.43	0.58	0.44	45	83	6
Fat (g)	53.75 \pm 35.81	48.79 \pm 27.91*	0.57	0.38	45	80	4
SFA (g)	16.37 \pm 11.36	14.70 \pm 8.56*	0.58	0.47	42	82	3
MUFA (g)	17.23 \pm 12.33	15.60 \pm 9.26*	0.58	0.44	47	83	3
PUFA (g)	12.78 \pm 7.94	11.92 \pm 6.89*	0.53	0.30	45	81	5
n-3 FA (g)	1.42 \pm 0.93	1.32 \pm 0.82*	0.56	0.26	45	81	3
n-6 FA (g)	11.52 \pm 7.19	10.76 \pm 6.19*	0.52	0.31	42	82	5
Cholesterol (mg)	325.66 \pm 220.12	314.29 \pm 217.52*	0.56	0.37	47	82	2
Carbohydrate (g)	316.72 \pm 139.68	275.33 \pm 145.29	0.42	0.43	42	78	6
Dietary fiber (g)	18.52 \pm 10.93	17.16 \pm 11.30	0.59	0.38	47	86	3
Calcium (mg)	479.81 \pm 273.13	432.94 \pm 250.77	0.56	0.39	46	83	4
Phosphorus (mg)	1,040.00 \pm 534.00	942.82 \pm 523.86	0.57	0.43	43	81	4
Iron (mg)	13.46 \pm 7.24	12.47 \pm 7.59	0.60	0.38	44	86	4
Sodium (mg)	3,460.70 \pm 2,174.75	3,134.49 \pm 2,071.77	0.54	0.16 [†]	44	84	3
Potassium (mg)	2,674.20 \pm 1,482.59	2,491.17 \pm 1,529.11	0.62	0.49	49	86	4
Vitamin A (μ g RE)	595.63 \pm 361.76	577.54 \pm 400.01*	0.53	0.17	45	82	4
Retinol (μ g)	116.85 \pm 78.58	105.51 \pm 67.61*	0.59	0.49	40	83	2
Carotene (μ g)	2,651.50 \pm 1,757.31	2,666.50 \pm 2,065.52*	0.45	0.11 [†]	46	81	4
Thiamin (mg)	1.90 \pm 1.00	1.70 \pm 0.92	0.55	0.45	42	82	6
Riboflavin (mg)	1.53 \pm 0.87	1.40 \pm 0.77	0.57	0.39	41	86	2
Niacin (mg)	14.47 \pm 8.71	12.88 \pm 8.00	0.56	0.43	47	86	5
Vitamin C (mg)	98.21 \pm 101.76	88.99 \pm 81.50*	0.57	0.39	40	83	2

FFQ1, the first food frequency questionnaire; FFQ2, the second food frequency questionnaire; SFA, saturated fatty acids; MUFA, monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; FA, fatty acids.

¹⁾Non-significant differences in mean intakes at $\alpha = 0.05$ between FFQs were noted as “**”.

²⁾Correlation coefficients were statistically significant at $\alpha = 0.05$, unless otherwise noted as “†”.

Table 4. Reproducibility of selected food items between the first and second food frequency questionnaires

Food item	All (n = 243)				< 3 years apart (n = 139)			
	Spearman's correlation	Quartile agreement			Spearman's correlation	Quartile agreement		
		Same	Same or adjacent	Opposite		Same	Same or adjacent	Opposite
Cooked rice	0.25	35	69	8	0.33	42	76	6
Cooked rice with other grains and legumes	0.51	43	80	6	0.52	45	82	6
Instant noodles, instant cup noodles	0.45	42	82	4	0.47	42	84	5
Loaf bread	0.45	43	79	7	0.47	49	81	9
Butter, margarine	0.36	-	-	-	0.46	-	-	-
Jam	0.49	-	-	-	0.55	-	-	-
Sweet red-beans buns, steamed sweet red-bean buns, cream buns	0.41	-	-	-	0.47	-	-	-
Pizza	0.36	33	76	7	0.46	37	79	6
Hamburger, sandwich	0.35	37	74	8	0.47	42	79	6
Plain steamed rice-cake, steamed rice-cake with red-bean, cubed rice-cake with soybean powder, plain cubed rice-cake	0.49	-	-	-	0.53	-	-	-
Cereal	0.36	-	-	-	0.41	-	-	-
Sea mustard soup	0.44	41	83	2	0.39	40	81	2
Bean paste soup	0.33	40	71	5	0.34	42	74	4
Kimchi stew, stir-fried kimchi	0.33	39	77	8	0.24	42	74	9
Bean curd, bean curd boiled in soy sauce, pan-fried bean curd	0.43	37	77	6	0.44	40	78	4
Soybean boiled in soy sauce	0.41	-	-	-	0.37	-	-	-
Fried egg, fried egg roll	0.41	39	79	3	0.44	41	81	4
Grilled pork belly	0.50	40	81	4	0.60	41	83	2
Stir-fried beef	0.45	37	77	3	0.52	35	80	5
Ham	0.43	39	78	8	0.48	40	78	8
Fried chicken	0.53	44	86	2	0.53	47	83	1
Anchovy, stir-fried anchovy	0.41	45	78	6	0.38	39	77	6
Bean sprout (seasoned, soup), seasoned mung bean sprout	0.40	33	78	7	0.45	42	81	5
Cucumber (seasoned, raw)	0.35	38	77	6	0.45	40	81	4
Vegetables salad	0.39	37	77	7	0.38	40	77	6
Boiled broccoli, boiled cabbage	0.46	45	73	7	0.54	50	78	7
Korean cabbage kimchi	0.46	42	79	4	0.51	48	84	8
Pickled vegetable (pepper, garlic, sesame leaf, onion, radish), pickled cucumber	0.37	46	74	10	0.36	40	72	10
Grilled laver, raw laver, seasoned laver	0.34	40	72	7	0.34	35	76	6
Stir-fried potatoes, potatoes boiled with soy sauce	0.45	39	81	5	0.45	42	76	5
Milk (normal)	0.31	-	-	-	0.26	-	-	-
Soybean milk	0.38	-	-	-	0.36	-	-	-
Tomato, cherry tomato	0.36	-	-	-	0.42	-	-	-
Apple	0.38	42	70	7	0.41	41	76	7
Banana	0.34	35	79	7	0.41	36	78	5
Coffee	0.54	46	83	5	0.54	46	83	5
Soft drink (cola, soda, fruit juice soda)	0.51	44	76	5	0.51	47	80	4
Snack	0.47	41	79	4	0.42	43	79	5
Cookie, cracker	0.36	33	80	7	0.40	42	77	9
Beer	0.72	58	86	2	0.74	60	87	2

Correlation coefficients were statistically significant at $\alpha = 0.05$, unless otherwise noted as “**”.

40%, 78%, and 6%, respectively, among the 243 women and 42%, 79%, and 5%, respectively, among the 139 women. The correlation coefficients and quartile agreements of all food items of the FFQs are listed in **Supplementary Table 1**. Spearman's correlation coefficients of food intake between FFQ1 and FFQ2 ranged from 0.18 for persimmon/dried persimmon to 0.72 for beer (median $r = 0.39$) among the 243 women and from 0.10 for strawberry to 0.74 for beer (median $r = 0.41$) among 139 women with a time interval of < 3 years since FFQ1. The median of the same and opposite classification was 40% and 6% among the 243 women and 42% and 6% among the 139 women.

DISCUSSION

This study aimed to assess the reproducibility of the web-based FFQ used in the KNHS among 243 female nurses. A 112-item FFQ was administered twice at a mean interval of 2.76 years, ranging from 9.24 months to 5.64 years. The mean daily intakes of energy and nutrients were lower from FFQ2 than from FFQ1. The median of reproducibility correlation coefficients was 0.51 (range, 0.41–0.55) for energy and raw nutrients and 0.41 (ranged, 0.25–0.72) for food items. The median percentages of participants who were classified into the same quartile and the same or adjacent quartile were 42% and 82%, respectively, for energy and nutrients. For food items, the median percentages of participants who were classified into the same quartile and the same or adjacent quartile were 40% and 78%, respectively.

The following reasons may have caused the differences in mean intakes between FFQ1 and FFQ2: aging, changes in dietary intakes, or random variation. However, although the comparison of mean intakes is a simple approach for evaluating the performance of FFQs, it does not provide information on the ability of the FFQ to rank individuals [9]. Correlations between the repeated measures of FFQs could provide useful information for distinguishing among individuals by dietary intakes. We found a moderate correlation between nutrient intakes from FFQ1 and those from FFQ2. In our study, energy adjustment slightly attenuated the correlation coefficients. The attenuation of the correlation coefficients for the reproducibility of the FFQ has also been observed in some previous studies in Korea [22–24], Japan [25–27], and Western countries [28–30]. Although the underlying reasons remain unclear, adjustment for energy intake could add unexplained variation.

We included women who had a time interval between FFQs that ranged from 9.24 months to 5.64 years. Because FFQ reproducibility has been often evaluated by assessing diet using the same questionnaires 1-year apart, we limited the analysis to women who completed FFQ2 within the 3 years following completion of FFQ1. Reproducibility was slightly higher when we limited the analysis to women with a shorter interval from FFQ1 completion than those with a longer interval. A few studies examined temporal changes in the reproducibility of the FFQ and observed a decline in correlation coefficients over time [31,32]. It may be natural that we observed slightly higher correlations because diet can change over time. The mean time interval between FFQs was 2.76 years among all participants in our study, comparable to those of several previous studies reporting various time intervals between FFQs, ranging from 9 months to 5.7 years [3,14,29,33–35].

Our results are comparable with those of a previous study that examined the reproducibility of the same FFQ [14]. In the previous validity and reproducibility study of the FFQ, FFQs were administered twice at 9 months apart to 126 Korean men and women aged 20–65 years. The median correlation coefficients between the 2 FFQs were 0.55 (range, 0.53 for phosphorus to 0.62 for fat) for energy and thirteen nutrients and 0.61 (range, 0.33 for rice to 0.87 for alcoholic beverages) for eleven food groups. The median proportions of participants whose dietary intakes were classified into the same or adjacent quartile was 83% for energy and nutrients, and 87% for food groups. The median of correlations for energy-adjusted nutrients and food groups was 0.55 for both nutrients and food groups [14].

Similar reproducibility results have been reported in several previous Korean studies. The validity and reproducibility of the FFQ were assessed among middle-aged men and women [22,23,36] and young women [24] in Korea. The correlation coefficients for the

reproducibility of the FFQ generally ranged from 0.3 to 0.6 for energy and nutrients [22-24,36], and energy adjustment attenuated the correlation coefficients in most studies [22-24]. For example, in the Korean diet and cancer research, 115 men and 173 women aged 30–66 years completed 2 FFQs administered approximately 9 months apart [23]. Among the 173 women, the correlation coefficients between 2 FFQs ranged from 0.42 for vitamin C to 0.59 for fat (median $r = 0.54$) for energy and fifteen nutrients and from 0.24 for niacin to 0.57 for calcium (median $r = 0.49$) for energy-adjusted nutrients. For eleven food groups, the median (range) correlation coefficient was 0.56 (0.35 for fruit to 0.72 for alcoholic beverages) for unadjusted values and 0.50 (0.36 for fruit to 0.72 for alcoholic beverages) for energy-adjusted values among women [23].

Our results are also consistent with findings of studies in other populations. The correlation coefficients for the reproducibility of the FFQ generally ranged from 0.4 to 0.7 for energy and nutrients [26,27,33-35,37-49] and ranged from 0.3 to 0.8 for foods [26,27,33,35,38,41,46,47,50-52] among female adults. Most previous studies used a self-administered FFQ to assess dietary intake over the preceding year and re-administered the questionnaire at approximately 1-year interval. For the reproducibility of food intake, correlations were the highest for alcoholic beverages in several studies [26,27,33,35,51]. In the US Nurses' Health Study, 1,497 female nurses self-administered 2 FFQs 9 months apart, and the mean (range) of correlation coefficients was 0.57 (0.40 for trans-fatty acids to 0.71 for vitamin E) for energy-adjusted nutrients and 0.56 (0.34 for ready-made pie to 0.76 for tea) for foods [38]. In the Shanghai Women's Health Study, 2 FFQs were administered approximately 2 years apart to 191 Chinese women by interviewers [46]. In this study, the mean (range) of correlation coefficients was 0.51 (0.30 for carotene to 0.59 for retinol) for energy and nutrients and 0.49 (0.37 for soy foods to 0.66 for fruits) for food groups. The mean (range) percentage of the same quartile classification was 36% (32% to 39%) for energy and nutrients and 41% (35–51%) for food groups.

Web-based surveys have several advantages over traditional survey methods, such as access to individuals at distant locations, ability to collect real-time data, low administration cost, and ease of follow-up [53,54]. Because of the advances in technologies and increased internet access, online surveys have become common research tools. Dietary assessment methods are also integrated with novel technologies in current epidemiological research [55]. FFQs are suitable for a web-based survey because the respondent's burden is modest and they can be self-administered. In addition, data processing of the FFQ is inexpensive and can be easily computerized [9]. Paper and web-based FFQ have shown comparable performance in several studies [56-59]. Reproducibility of the web-based FFQs has shown moderate-to-high correlation coefficients among adults [60-64] and adolescents [65-67]. Most studies used web-based FFQs that assess dietary intake over the preceding month, and FFQs were administered twice at one-month interval [60-62,64-67].

The strength of this study is a large sample size and extensive evaluation of various nutrients and foods for the reproducibility of the FFQ. The limitation of our study is a relatively long-time interval between FFQs compared to the previous studies. However, when we examined the reproducibility among those with a shorter time interval between FFQs, we found acceptable reproducibility. Our results were comparable to those from previous studies that examined the reproducibility of the FFQ with intervals of 9 months or 1 year. Further research, including that on the association of diet and disease, is needed to support the validity of the questionnaire.

In conclusion, the FFQ used in the KNHS study has acceptable reproducibility. The estimates of reproducibility in our study are similar to a study that examined the reproducibility of the same FFQ in Korean men and women, and studies that evaluated the reproducibility of FFQs among women. This study may be a useful resource for testing future KNHS hypotheses about dietary roles in women's health and wellness.

SUPPLEMENTARY MATERIAL

Supplementary Table 1

Reproducibility of food items between the first and second food frequency questionnaires

[Click here to view](#)

REFERENCES

1. Peters SA, Woodward M, Jha V, Kennedy S, Norton R. Women's health: a new global agenda. *BMJ Glob Health* 2016;1:e000080.
[CROSSREF](#)
2. World Health Organization. *World Health Statistics 2019: Monitoring Health for the SDGs, Sustainable Development Goals*. Geneva: World Health Organization; 2019.
3. Colditz GA, Philpott SE, Hankinson SE. The impact of the Nurses' Health Study on population health: prevention, translation, and control. *Am J Public Health* 2016;106:1540-5.
[PUBMED](#) | [CROSSREF](#)
4. Design of the Women's Health Initiative clinical trial and observational study. The Women's Health Initiative Study Group. *Control Clin Trials* 1998;19:61-109.
[PUBMED](#) | [CROSSREF](#)
5. Zheng W, Chow WH, Yang G, Jin F, Rothman N, Blair A, Li HL, Wen W, Ji BT, Li Q, et al. The Shanghai Women's Health Study: rationale, study design, and baseline characteristics. *Am J Epidemiol* 2005;162:1123-31.
[PUBMED](#) | [CROSSREF](#)
6. Folsom AR, Kushi LH, Anderson KE, Mink PJ, Olson JE, Hong CP, Sellers TA, Lazovich D, Prineas RJ. Associations of general and abdominal obesity with multiple health outcomes in older women: the Iowa Women's Health Study. *Arch Intern Med* 2000;160:2117-28.
[PUBMED](#) | [CROSSREF](#)
7. Darnton-Hill I, Nishida C, James WP. A life course approach to diet, nutrition and the prevention of chronic diseases. *Public Health Nutr* 2004;7:101-21.
[PUBMED](#) | [CROSSREF](#)
8. Barker DJ. Maternal nutrition, fetal nutrition, and disease in later life. *Nutrition* 1997;13:807-13.
[PUBMED](#) | [CROSSREF](#)
9. Willett W. *Nutritional Epidemiology*. New York (NY): Oxford University Press; 2012.
10. Hu FB, Satija A, Rimm EB, Spiegelman D, Sampson L, Rosner B, Camargo CA Jr, Stampfer M, Willett WC. Diet assessment methods in the Nurses' Health Studies and contribution to evidence-based nutritional policies and guidelines. *Am J Public Health* 2016;106:1567-72.
[PUBMED](#) | [CROSSREF](#)
11. Yu E, Rimm E, Qi L, Rexrode K, Albert CM, Sun Q, Willett WC, Hu FB, Manson JE. Diet, lifestyle, biomarkers, genetic factors, and risk of cardiovascular disease in the Nurses' Health Studies. *Am J Public Health* 2016;106:1616-23.
[PUBMED](#) | [CROSSREF](#)
12. Kim O, Ahn Y, Lee HY, Jang HJ, Kim S, Lee JE, Jung H, Cho E, Lim JY, Kim MJ, et al. The Korea Nurses' Health Study: a prospective cohort study. *J Womens Health (Larchmt)* 2017;26:892-9.
[PUBMED](#) | [CROSSREF](#)
13. Yun SH, Shim JS, Kweon S, Oh K. Development of a food frequency questionnaire for the Korea National Health and Nutrition Examination Survey: data from the fourth Korea National Health and Nutrition Examination Survey (KNHANES IV). *Korean J Nutr* 2013;46:186-96.
[CROSSREF](#)

14. Kim DW, Song S, Lee JE, Oh K, Shim J, Kweon S, Paik HY, Joung H. Reproducibility and validity of an FFQ developed for the Korea National Health and Nutrition Examination Survey (KNHANES). *Public Health Nutr* 2015;18:1369-77.
[PUBMED](#) | [CROSSREF](#)
15. Kim Y, Je Y. Moderate coffee consumption is inversely associated with the metabolic syndrome in the Korean adult population. *Br J Nutr* 2018;120:1279-87.
[PUBMED](#) | [CROSSREF](#)
16. Kim Y, Je Y. A modified Mediterranean diet score is inversely associated with metabolic syndrome in Korean adults. *Eur J Clin Nutr* 2018;72:1682-9.
[PUBMED](#) | [CROSSREF](#)
17. Shin S, Kim SA, Ha J, Lim K. Sugar-sweetened beverage consumption in relation to obesity and metabolic syndrome among Korean adults: a cross-sectional study from the 2012–2016 Korean national health and nutrition examination survey (KNHANES). *Nutrients* 2018;10:1467.
[CROSSREF](#)
18. Choi A, Ha K, Joung H, Song Y. Frequency of consumption of whole fruit, not fruit juice, is associated with reduced prevalence of obesity in Korean adults. *J Acad Nutr Diet* 2019;119:1842-1851.e2.
[PUBMED](#) | [CROSSREF](#)
19. Kim H, Lee K, Rebholz CM, Kim J. Association between unhealthy plant-based diets and the metabolic syndrome in adult men and women: a population-based study in South Korea. *Br J Nutr* 2021;125:1-14.
[PUBMED](#) | [CROSSREF](#)
20. Bao Y, Bertoin ML, Lenart EB, Stampfer MJ, Willett WC, Speizer FE, Chavarro JE. Origin, methods, and evolution of the three Nurses' Health Studies. *Am J Public Health* 2016;106:1573-81.
[PUBMED](#) | [CROSSREF](#)
21. Korea Disease Control and Prevention Agency. Nutrient Database for Food Frequency Questionnaire in the Korea National Health and Nutrition Examination Survey. Cheongju: Korea Disease Control and Prevention Agency; 2015.
22. Ahn Y, Kwon E, Shim JE, Park MK, Joo Y, Kimm K, Park C, Kim DH. Validation and reproducibility of food frequency questionnaire for Korean genome epidemiologic study. *Eur J Clin Nutr* 2007;61:1435-41.
[PUBMED](#) | [CROSSREF](#)
23. Park MK, Noh HY, Song NY, Paik HY, Park S, Joung H, Song WO, Kim J. Validity and reliability of a dish-based, semi-quantitative food frequency questionnaire for Korean diet and cancer research. *Asian Pac J Cancer Prev* 2012;13:545-52.
[PUBMED](#) | [CROSSREF](#)
24. Bae YJ, Choi HY, Sung MK, Kim MK, Choi MK. Validity and reproducibility of a food frequency questionnaire to assess dietary nutrients for prevention and management of metabolic syndrome in Korea. *Nutr Res Pract* 2010;4:121-7.
[PUBMED](#) | [CROSSREF](#)
25. Imaeda N, Fujiwara N, Tokudome Y, Ikeda M, Kuriki K, Nagaya T, Sato J, Goto C, Maki S, Tokudome S. Reproducibility of a semi-quantitative food frequency questionnaire in Japanese female dietitians. *J Epidemiol* 2002;12:45-53.
[PUBMED](#) | [CROSSREF](#)
26. Ishihara J, Sobue T, Yamamoto S, Yoshimi I, Sasaki S, Kobayashi M, Takahashi T, Itoi Y, Akabane M, Tsugane S, et al. Validity and reproducibility of a self-administered food frequency questionnaire in the JPHC Study Cohort II: study design, participant profile and results in comparison with Cohort I. *J Epidemiol* 2003;13:S134-47.
[PUBMED](#) | [CROSSREF](#)
27. Tsubono Y, Kobayashi M, Sasaki S, Tsugane S; JPHC. Validity and reproducibility of a self-administered food frequency questionnaire used in the baseline survey of the JPHC Study Cohort I. *J Epidemiol* 2003;13:S125-33.
[PUBMED](#) | [CROSSREF](#)
28. Franceschi S, Barbone F, Negri E, Decarli A, Ferraroni M, Filiberti R, Giacosa A, Gnagnarella P, Nanni O, Salvini S, et al. Reproducibility of an Italian food frequency questionnaire for cancer studies. Results for specific nutrients. *Ann Epidemiol* 1995;5:69-75.
[PUBMED](#) | [CROSSREF](#)
29. Riboli E, Toniolo P, Kaaks R, Shore RE, Casagrande C, Pasternack BS. Reproducibility of a food frequency questionnaire used in the New York University Women's Health Study: effect of self-selection by study subjects. *Eur J Clin Nutr* 1997;51:437-42.
[PUBMED](#) | [CROSSREF](#)

30. Bohlscheid-Thomas S, Hoting I, Boeing H, Wahrendorf J. Reproducibility and relative validity of energy and macronutrient intake of a food frequency questionnaire developed for the German part of the EPIC project. European Prospective Investigation into Cancer and Nutrition. *Int J Epidemiol* 1997;26 Suppl 1:S71-81.
[PUBMED](#) | [CROSSREF](#)
31. Goldbohm RA, van't Veer P, van den Brandt PA, van't Hof MA, Brants HA, Sturmans F, Hermus RJ. Reproducibility of a food frequency questionnaire and stability of dietary habits determined from five annually repeated measurements. *Eur J Clin Nutr* 1995;49:420-9.
[PUBMED](#)
32. Tsubono Y, Nishino Y, Fukao A, Hisamichi S, Tsugane S. Temporal change in the reproducibility of a self-administered food frequency questionnaire. *Am J Epidemiol* 1995;142:1231-5.
[PUBMED](#) | [CROSSREF](#)
33. Pisani P, Faggiano F, Krogh V, Palli D, Vineis P, Berrino F. Relative validity and reproducibility of a food frequency dietary questionnaire for use in the Italian EPIC centres. *Int J Epidemiol* 1997;26 Suppl 1:S152-60.
[PUBMED](#) | [CROSSREF](#)
34. Willett WC, Sampson L, Browne ML, Stampfer MJ, Rosner B, Hennekens CH, Speizer FE. The use of a self-administered questionnaire to assess diet four years in the past. *Am J Epidemiol* 1988;127:188-99.
[PUBMED](#) | [CROSSREF](#)
35. Nagel G, Zoller D, Ruf T, Rohrmann S, Linseisen J. Long-term reproducibility of a food-frequency questionnaire and dietary changes in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Heidelberg cohort. *Br J Nutr* 2007;98:194-200.
[PUBMED](#) | [CROSSREF](#)
36. Lee Y, Park K. Reproducibility and validity of a semi-quantitative FFQ for trace elements. *Br J Nutr* 2016;116:864-73.
[PUBMED](#) | [CROSSREF](#)
37. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH, Speizer FE. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol* 1985;122:51-65.
[PUBMED](#) | [CROSSREF](#)
38. Colditz GA, Willett WC, Stampfer MJ, Sampson L, Rosner B, Hennekens CH, Speizer FE. The influence of age, relative weight, smoking, and alcohol intake on the reproducibility of a dietary questionnaire. *Int J Epidemiol* 1987;16:392-8.
[PUBMED](#) | [CROSSREF](#)
39. Munger RG, Folsom AR, Kushi LH, Kaye SA, Sellers TA. Dietary assessment of older Iowa women with a food frequency questionnaire: nutrient intake, reproducibility, and comparison with 24-hour dietary recall interviews. *Am J Epidemiol* 1992;136:192-200.
[PUBMED](#) | [CROSSREF](#)
40. Martin-Moreno JM, Boyle P, Gorgojo L, Maisonneuve P, Fernandez-Rodriguez JC, Salvini S, Willett WC. Development and validation of a food frequency questionnaire in Spain. *Int J Epidemiol* 1993;22:512-9.
[PUBMED](#) | [CROSSREF](#)
41. Männistö S, Virtanen M, Mikkonen T, Pietinen P. Reproducibility and validity of a food frequency questionnaire in a case-control study on breast cancer. *J Clin Epidemiol* 1996;49:401-9.
[PUBMED](#) | [CROSSREF](#)
42. Xing X, Burr JA, Brasure JR, Neugut AI, Marshall JR. Reproducibility of nutrient intake in a food frequency questionnaire used in a general population. *Nutr Cancer* 1996;25:259-68.
[PUBMED](#) | [CROSSREF](#)
43. Katsouyanni K, Rimm EB, Gnardellis C, Trichopoulos D, Polychronopoulos E, Trichopoulou A. Reproducibility and relative validity of an extensive semi-quantitative food frequency questionnaire using dietary records and biochemical markers among Greek schoolteachers. *Int J Epidemiol* 1997;26 Suppl 1:S118-27.
[PUBMED](#) | [CROSSREF](#)
44. Ocké MC, Bueno-de-Mesquita HB, Pols MA, Smit HA, van Staveren WA, Kromhout D. The Dutch EPIC food frequency questionnaire. II. Relative validity and reproducibility for nutrients. *Int J Epidemiol* 1997;26 Suppl 1:S49-58.
[PUBMED](#) | [CROSSREF](#)
45. Romieu I, Parra S, Hernández JF, Madrigal H, Willett W, Hernández M. Questionnaire assessment of antioxidants and retinol intakes in Mexican women. *Arch Med Res* 1999;30:224-39.
[PUBMED](#) | [CROSSREF](#)
46. Shu XO, Yang G, Jin F, Liu D, Kushi L, Wen W, Gao YT, Zheng W. Validity and reproducibility of the food frequency questionnaire used in the Shanghai Women's Health Study. *Eur J Clin Nutr* 2004;58:17-23.
[PUBMED](#) | [CROSSREF](#)

47. Parr CL, Veierød MB, Laake P, Lund E, Hjartåker A. Test-retest reproducibility of a food frequency questionnaire (FFQ) and estimated effects on disease risk in the Norwegian Women and Cancer Study (NOWAC). *Nutr J* 2006;5:4.
[PUBMED](#) | [CROSSREF](#)
48. Horn-Ross PL, Lee VS, Collins CN, Stewart SL, Canchola AJ, Lee MM, Reynolds P, Clarke CA, Bernstein L, Stram DO. Dietary assessment in the California Teachers Study: reproducibility and validity. *Cancer Causes Control* 2008;19:595-603.
[PUBMED](#) | [CROSSREF](#)
49. Mirmiran P, Esfahani FH, Mehrabi Y, Hedayati M, Azizi F. Reliability and relative validity of an FFQ for nutrients in the Tehran lipid and glucose study. *Public Health Nutr* 2010;13:654-62.
[PUBMED](#) | [CROSSREF](#)
50. Xing X, Burr JA, Brasure JR, Neugut AI, Marshall JR. Reproducibility of food intake in a food frequency questionnaire used in a general population. *Nutr Cancer* 1995;24:85-95.
[PUBMED](#) | [CROSSREF](#)
51. Ocké MC, Bueno-de-Mesquita HB, Goddijn HE, Jansen A, Pols MA, van Staveren WA, Kromhout D. The Dutch EPIC food frequency questionnaire. I. Description of the questionnaire, and relative validity and reproducibility for food groups. *Int J Epidemiol* 1997;26 Suppl 1:S37-48.
[PUBMED](#) | [CROSSREF](#)
52. Esfahani FH, Asghari G, Mirmiran P, Azizi F. Reproducibility and relative validity of food group intake in a food frequency questionnaire developed for the Tehran Lipid and Glucose Study. *J Epidemiol* 2010;20:150-8.
[PUBMED](#) | [CROSSREF](#)
53. Evans JR, Mathur A. The value of online surveys. *Internet Res* 2005;15:195-219.
[CROSSREF](#)
54. Wright KB. Researching Internet-based populations: advantages and disadvantages of online survey research, online questionnaire authoring software packages, and web survey services. *J Comput Mediat Commun* 2005;10:JCMC1034.
[CROSSREF](#)
55. Naska A, Lagiou A, Lagiou P. Dietary assessment methods in epidemiological research: current state of the art and future prospects. *F1000 Res* 2017;6:926.
[PUBMED](#) | [CROSSREF](#)
56. Beasley JM, Davis A, Riley WT. Evaluation of a web-based, pictorial diet history questionnaire. *Public Health Nutr* 2009;12:651-9.
[PUBMED](#) | [CROSSREF](#)
57. González Carrascosa R, García Segovia P, Martínez Monzó J. Paper and pencil vs online self-administered food frequency questionnaire (FFQ) applied to university population: a pilot study. *Nutr Hosp* 2011;26:1378-84.
[PUBMED](#)
58. Lo Siou G, Csizmadi I, Boucher BA, Akawung AK, Whelan HK, Sharma M, Al Rajabi A, Vena JE, Kirkpatrick SI, Koushik A, et al. The comparative reliability and feasibility of the past-year Canadian diet history questionnaire II: comparison of the paper and web versions. *Nutrients* 2017;9:133.
[PUBMED](#) | [CROSSREF](#)
59. Zazpe I, Santiago S, De la Fuente-Arrillaga C, Nuñez-Córdoba JM, Bes-Rastrollo M, Martínez-González MA. Paper-based versus web-based versions of self-administered questionnaires, including food-frequency questionnaires: prospective cohort study. *JMIR Public Health Surveill* 2019;5:e11997.
[PUBMED](#) | [CROSSREF](#)
60. Engle A, Lynn LL, Koury K, Boyar AP. Reproducibility and comparability of a computerized, self-administered food frequency questionnaire. *Nutr Cancer* 1990;13:281-92.
[PUBMED](#) | [CROSSREF](#)
61. Labonté ME, Cyr A, Baril-Gravel L, Royer MM, Lamarche B. Validity and reproducibility of a web-based, self-administered food frequency questionnaire. *Eur J Clin Nutr* 2012;66:166-73.
[PUBMED](#) | [CROSSREF](#)
62. Fallaize R, Forster H, Macready AL, Walsh MC, Mathers JC, Brennan L, Gibney ER, Gibney MJ, Lovegrove JA. Online dietary intake estimation: reproducibility and validity of the Food4Me food frequency questionnaire against a 4-day weighed food record. *J Med Internet Res* 2014;16:e190.
[PUBMED](#) | [CROSSREF](#)
63. Kristal AR, Kolar AS, Fisher JL, Plascak JJ, Stumbo PJ, Weiss R, Paskett ED. Evaluation of web-based, self-administered, graphical food frequency questionnaire. *J Acad Nutr Diet* 2014;14:613-21.
[PUBMED](#) | [CROSSREF](#)

64. Marshall SJ, Livingstone KM, Celis-Morales C, Forster H, Fallaize R, O'Donovan CB, Woolhead C, Marsaux CF, Macready AL, Navas-Carretero S, et al. Reproducibility of the online Food4Me food-frequency questionnaire for estimating dietary intakes across Europe. *J Nutr* 2016;146:1068-75.
[PUBMED](#) | [CROSSREF](#)
65. Matthys C, Pynaert I, De Keyzer W, De Henauw S. Validity and reproducibility of an adolescent web-based food frequency questionnaire. *J Am Diet Assoc* 2007;107:605-10.
[PUBMED](#) | [CROSSREF](#)
66. Filippi AR, Amodio E, Napoli G, Breda J, Bianco A, Jemni M, Censi L, Mammina C, Tabacchi G. The web-based ASSO-food frequency questionnaire for adolescents: relative and absolute reproducibility assessment. *Nutr J* 2014;13:119.
[PUBMED](#) | [CROSSREF](#)
67. Overby NC, Johannesen E, Jensen G, Skjævesland AK, Haugen M. Test-retest reliability and validity of a web-based food-frequency questionnaire for adolescents aged 13–14 to be used in the Norwegian Mother and Child Cohort Study (MoBa). *Food Nutr Res* 2014;58.
[PUBMED](#) | [CROSSREF](#)