

Review



Association of added sugar intake with all-cause and cardiovascular disease mortality: a systematic review of cohort studies

SuJin Song ¹, Jae Eun Shim ^{2§}, and YoonJu Song ³

¹Department of Food and Nutrition, Hannam University, Daejeon 34054, Korea

²Department of Food and Nutrition, Daejeon University, Daejeon 34520, Korea

³Department of Food Science and Nutrition, The Catholic University of Korea, Bucheon 14662, Korea



Received: Nov 4, 2021

Revised: Jan 5, 2022

Accepted: Jan 10, 2022

Published online: Feb 17, 2022

§Corresponding Author:

Jae Eun Shim

Department of Food and Nutrition, Daejeon University, 62 Daehak-ro, Dong-gu, Daejeon 34520, Korea.

Tel. +82-42-280-2469

Fax. +82-42-280-2468

Email. jshim@dju.kr


©2022 The Korean Nutrition Society and the Korean Society of Community Nutrition

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.


ORCID iDs

SuJin Song 

<https://orcid.org/0000-0003-1871-4346>

Jae Eun Shim 

<https://orcid.org/0000-0001-8458-9112>

YoonJu Song 

<https://orcid.org/0000-0002-4764-5864>

Funding

This research was supported by Policy Research Program for (project No. 2018O415A13-00, 25193068200, 25203084501) from the Ministry of Health and Welfare in 2018–2020.

ABSTRACT

BACKGROUND/OBJECTIVES: The long-term health impacts of dietary sugar have begun to be reported. The committee on the 2020 Dietary Reference Intakes for Koreans for sugar intended to update the scientific evidence on the association of added sugar intake with health outcomes, including mortality. This study evaluated the evidence on the association of dietary sugar with the all-cause and cardiovascular disease mortality from cohort studies.

SUBJECTS/METHODS: The PubMed and ScienceDirect databases were searched until July 2021. The search terms were based on multiple combinations of keywords, such as added sugar, beverage, and mortality, included in the title or abstract. Two authors independently assessed the eligibility criteria of study selection and extracted the information from each selected article for this systematic review.

RESULTS: The literature search identified 276 articles. Seventeen cohort studies met the inclusion criteria and were included in this systematic review. This study extracted information on the cohort and participants, dietary assessment methods, exposure and its comparison groups, health outcomes, and risk measures. Five articles reported added sugar intake, and 15 articles reported the food sources of the added sugar in the association with mortality. Increased mortality was observed in higher added sugar intake and a lower intake of less than 5% of energy, while higher discrepancies between gender, age groups, and countries were noted. In addition, nutritious sugary foods and beverages did not increase mortality.

CONCLUSIONS: A ‘U’ or ‘J’ shaped relationship between the added sugar intake and mortality was estimated. The observed discrepancy indicated the need for more evidence to establish the dose-response relationship for Koreans.

Keywords: Dietary sugars; sugar-sweetened beverages; mortality

INTRODUCTION

Dietary sugar intake has been reported to link with several cardio-metabolic risk factors. Meta-analysis of prospective cohort studies and randomized clinical trials of the sugar intake reported a significant association with obesity and the morbidity of major non-communicable diseases (NCDs). Free sugar intake was positively correlated with the body weight gain in both adults and children, but there was no association with isoenergetic

Conflict of Interest

The authors declare no potential conflicts of interests.

Author Contributions

Conceptualization: Shim JE; Formal analysis: Song S, Shim JE, Song Y; Supervision: Shim JE; Writing - original draft: Song S, Shim JE, Song Y; Writing - review & editing: Song S, Shim JE.

replacement with other carbohydrate sources [1]. The results of meta-analysis of blood pressure and lipids also suggested a clear correlation with the sugar intake [2]. Many studies have been conducted on sugar-sweetened beverages (SSB), a major source of added sugar, and meta-analyses have also reported associations with obesity and major NCDs. The SSB intake was significantly associated with weight gain in both children and adults [3], and the group high in SSB intake increased the risk of type 2 diabetes and metabolic syndrome [4].

Dietary sugar intake in Korean adults is relatively low. According to the Korea National Health and Nutrition Examination Survey (KNHANES), the average daily intake of total sugar was 67.9 g in 2016, which decreased slightly to 61.7 g in 2019 [5]. However, SSB consumption increased, and the average energy contribution from SSB was 32 kcal/day in 1998 KNHANES, but it more than doubled to 82 kcal/day in the 2007–2009 KNHANES [6].

In studies of SSB in Korean adults, higher consumption of soft drinks (≥ 4 times per week) was associated with increased risk of metabolic syndrome only in women [7,8]. Other studies for Korean adults also reported that higher SSB consumption (\geq once per day) was associated with an increased risk of obesity in both men and women [9], and higher SSB consumption (median 3.5 servings per week) was associated with an increased incidence of hypertension [10].

On the other hand, a study of 2,599 Korean children and adolescents reported that the total dietary sugar intake was relatively low (12.5% in girls and 10.8% in boys), and even sugar intake from milk and fruits was inversely associated with pediatric overweight and obesity among girls [11]. Another study of Korean adults reported that high sugar intake ($> 20\%$ of energy) was associated with an increased risk of obesity and metabolic syndrome in men [12].

Although most studies reported a strong association of sugar intake or SSB consumption with an increased risk of cardio-metabolic risk factors, the long-term health impacts of dietary sugars are unclear. Several large cohort studies have reported a relationship between dietary sugar and mortality. A study of US adults using NHANES data (1988–1994) reported that those who consumed an added sugar intake of 25% of their energy intake or more showed 2.75 times higher cardiovascular disease (CVD) mortality than those who consumed less than 10% of energy [13]. On the other hand, a study of 4,000 Chinese elderly reported that the added sugar intake in the highest quintile was only 6.9% of energy. Those in the highest quintile had 25% lower CVD mortality than those in the lowest quintile.

In addition, a large prospective study of 353,751 US adults reported that the total fructose intake was weakly positively associated with the all-cause mortality in both men and women. In contrast, added sugar differed by the food source in that added sugar from solid foods was inversely associated with the all-cause mortality in both men and women, but added sugar from beverages was positively associated with the all-cause mortality in women [14]. This suggests that the health effects of dietary sugar should be determined from the levels of intake and its source.

Sugar intake is a modifiable dietary risk factor for NCDs. The World Health Organization (WHO) [15,16] set population intake goals to prevent NCDs, the leading causes of death worldwide. Recently, WHO developed evidence-based guidelines to provide recommendations on the intake of free sugars, focusing on the prevention and control of unhealthy weight gain and dental caries [1,17,18]. The first Dietary Reference Intakes for Koreans (KDRIs) for sugar was set in 2005, which was the acceptable macronutrient

distribution range for the total sugar as 10–20%. The committee on the 2015 KDRI examined the previous references for total sugar and recommended added sugar intake according to the free sugar guideline from WHO. The 2015 committee reviewed the scientific evidence on the association of sugar intake with the diet quality and NCDs morbidity more than obesity and dental caries. The 2020 KDRI committee for sugar updated the scientific evidence on the association of added sugar with NCDs morbidity and expanded the evidence into the association with mortality. This study provided a systematic review on the association of added sugar intake and SSB consumption with all-cause and CVD mortality to establish the 2020 KDRI for sugar.

SUBJECTS AND METHODS

Search strategy

Studies examining the associations of added sugar intake and its sources with mortality were identified by searching the PubMed and ScienceDirect published databases through July 2021. The following keywords were used for the search: (total sugar or added sugar or free sugar or dietary sugar or SSB or fruit juice or carbonated beverage or soft drink or sweet drink or soda appeared in the title or abstract) and (mortality or death appeared in title or abstract). In addition, the references were screened from the retrieved articles.

Selection criteria

The human studies published in English or Korean languages articles were included. Two authors (S.S. and J.E.S) assessed the eligibility criteria of the study selection independently as follows: (1) cohort design, published as full-text manuscripts of the original article; (2) exposure of interest was added sugar (free sugar) intake or its sources consumption; and (3) the endpoint of interest was all-cause or CVD mortality.

Data extraction

The following information was extracted from each article: (1) the first author's last name, publication year, and country in which the study was performed; (2) cohort name and number of study participants; (3) study period; (4) study participants' age and gender; (5) dietary assessment methods; (6) exposure and its comparison groups; (7) outcome and incidence; (8) follow-up period; (9) relative risk (RR) and 95% confidence interval (CI), and *P*-trend; and (10) covariates for adjustment in the analysis. The estimates adjusted for the most covariates were used when several estimates were reported.

RESULTS

Literature search and study characteristics

A preliminary literature search yielded 276 articles through a database search. The titles and abstracts of 252 articles were screened after excluding 24 duplicated or non-English studies. Of these, 228 articles were excluded based on the selection criteria, and 24 articles were identified for the full-text review. After the full-text review, 15 articles were excluded because 10 articles were not full-text manuscripts of the original article and 5 articles were not relevant to the exposure or outcome of interest. Furthermore, 8 additional articles were identified from the reference lists of retrieved articles. Finally, 17 cohort studies were included in this systematic review (**Fig. 1**) [13,14,19-33].




Fig. 1. Flowchart for the selection of studies included in the systematic review.

Table 1 lists the characteristics of 17 articles included in the current systematic review.

All studies were based on the cohort design and adults older than 20 years. Two studies included older adults aged ≥ 65 years only [23,24]. Nine studies were conducted in US adults [13,14,20,21,25,26,29,32,33], 5 studies in European adults [19,22,27,30,31], and 3 studies in Asian adults [23,24,28]. Most studies assessed the added sugar intake or its sources using a food-frequency questionnaire (FFQ) or dietary questionnaire. The follow-up periods of the studies ranged from 6 to 34 years. The estimates obtained after adjusting for potential confounders, including nutrients, demographic, anthropometric, medical, and lifestyle variables, were considered to identify associations of added sugar intake and its sources with mortality in this review. In terms of exposure of interest, 5 articles reported the findings on the associations between added sugar intake and mortality [13,14,23,24,30], and 15 articles reported the associations between food sources of added sugar and mortality [14,19-22,25-33].

Added sugar intake, SSB consumption, and all-cause mortality

Among 17 articles, 3 studies examined the association of added or free sugar intake with the all-cause mortality risk (**Table 2**) [14,24,30]. The sugar intake was assessed in the percentages of energy from sugar intake or grams per 1,000 kcal of energy intake. After adjusting for potential covariates, only one study of older Chinese adults showed significant inverse associations of added sugar and free sugar intakes with the risk of all-cause mortality (Q5 vs. Q1: RR = 0.75 for added sugar and 0.77 for free sugar) [24]. The other two studies conducted in Western adults did not have significant associations.

The association between SSB and all-cause mortality was examined by reviewing 11 studies (**Table 3**) [14,19-21,25,27-31,33]. Although SSB consumption was evaluated using different units and comparison groups, most studies showed positive associations between SSB consumption and the risk of all-cause mortality. A study performed on 10 European studies showed an inverse association between soft drink and all-cause mortality in non-diabetic individuals but not in diabetic individuals [31]. Tasevska *et al.* [14] reported that the added sugar intake from beverages was positively associated with the all-cause mortality in women but not in men.

Table 1. Characteristics of cohort studies included in the systematic review

Author, country	Year	Cohort	No. of participants	Study period	Age at baseline (Mean ± SD or range)	Men (%)	Dietary assessment	Exposure	Outcome (case/total)	Follow-up	Confounders adjusted				
											Nutrients	Demographic	Anthrop Medical	Lifestyles	
Paganini-Hill et al., US [29]	2007	The Leisure World Cohort Study	13,624	1980–2004	74 yrs (44–101 yrs)	36.6	Survey questionnaire	• Non-alcoholic beverages (cola and other soft drinks with sugar- or artificially sweetened) • Sugar and confectionery • Cakes and cookies • Soft drinks • Juices	• All-cause mortality: 11,386/13,624	23 yrs	X	0	0	0	0
Sluik et al., 10 European countries [31]	2014	EPIC	265,295 (diabetic: 6,384, non-diabetic: 258,911)	1992–	35–70 yrs	40.3	Dietary questionnaires, semi-quantitative FFQ, and combined dietary methods of food records and questionnaires	• Added sugar • Added sugar intake from beverages • Added sugar intake from solids	• All-cause mortality in diabetic: 830/6,384 • All-cause mortality in non-diabetic: 12,135/258,911	9.9 yrs	0	0	0	X	0
Tasevska et al., US [14]	2014	The NIH-AARP Diet and Health Study	353,751	1995–2008	50–71 yrs	58.3	DHQ	• Added sugar • Added sugar intake from beverages • Added sugar intake from solids	• All-cause mortality: M, 28,617/206,371; W, 15,062/147,381	12–13 yrs	0	0	0	0	0
Vyas et al., US [32]	2015	WHI-OS cohort	59,614	1993–2005	62.8 ± 7.2 yrs (50–79 yrs)	0.0	Questionnaire	• Diet drinks (such as diet coke or diet fruit drinks, with a 12 fl oz can as a reference size) • Frequency • Added sugar	• All-cause mortality: 4,437/59,614 • CVD mortality: 942/59,614	8.7 yrs	0	0	0	0	0
Yang et al., US [13]	2014	NHANES	11,733	1988–2010	≥ 20 yrs	48.1	24-h recall	• Added sugar	• CVD mortality: 831/11,733	14.6 yrs	0	0	0	0	0
Lai et al., UK [22]	2015	UKWCS	30,458	1995–2013	35–69 yrs	0.0	FFQ	• Pure fruit juice • Orange juice (pure fruit)	• CVD mortality: 286/30,458	16.7 yrs	0	0	0	0	X
Odegaard et al., Singapore [28]	2015	SCHS	52,584	1993–2011	45–74 yrs	44.0	Semi-quantitative FFQ	• Soft drink • Juice (fruit or vegetable)	• All-cause mortality: 10,029/52,584 • CVD mortality: 3,097/52,584	16.3 person-yrs	0	0	0	0	0
Barrington and White, US [20]	2016	VITAL study	69,582	2000–2008	50–76 yrs	48.8	Semi-quantitative FFQ	• Sugar-sweetened drinks (including sugar-sweetened soda (not diet), fruit drinks (not juice), and cranberry juice) • Added sugar • Added sugar intake from sweets • Added sugar intake from cereals	• All-cause mortality: 4,187/69,582 • CVD mortality: 1,066/69,582	6.9 yrs	0	0	0	0	0
Liu et al., China [23]	2018	Mr. and Ms. OS Hong Kong	3,416	2001–2014	M: 72.4 ± 5.0 yrs, W: 72.5 ± 5.3 yrs (≥ 65 yrs)	50.0	FFQ	• Added sugar • Added sugar intake from sweets • Added sugar intake from cereals	• CVD mortality: 174/3,416	11.1 yrs	0	0	0	0	0
Liu et al., China [24]	2018	Mr. and Ms. OS Hong Kong	3,416	2001–2014	72.5 ± 5.2 yrs (≥ 65 yrs)	50.0	FFQ	• Added sugar • Free sugar	• All-cause mortality: 780/3,416	11.1 yrs	0	0	0	0	0

(continued to the next page)

Table 1. (Continued) Characteristics of cohort studies included in the systematic review

Author, country	Year	Cohort	No. of participants	Study period	Age at baseline (Mean ± SD or range)	Men (%)	Dietary assessment	Exposure	Outcome (case/total)	Follow-up	Confounders adjusted					
											Nutrients	Demographic	Anthrop Medical	Lifestyles		
Collin <i>et al.</i> , US [21]	2019	REGARDS	13,440	2003–2018	63.6 ± 9.1 yrs (>45 yrs)	59.3	FFQ	• SSB • Naturally sweet fruit juice	• All-cause mortality: 1,000/13,440 • CHD mortality: 168/13,440	6.0 yrs	0	0	0	X	0	
Malik <i>et al.</i> , US [25]	2019	HPFS	37,716	1986–2014	40–75 yrs	100.0	FFQ	• SSB • ASB	• All-cause mortality: 13,004/37,716 • CVD mortality: 3,757/37,716	28 yrs	0	0	0	0	0	0
	2019	NHS	80,647	1976–2014	30–55 yrs	0.0	FFQ	• SSB • ASB	• All-cause mortality: 23,432/80,647 • CVD mortality: 4,139/80,647	34 yrs	0	0	0	0	0	0
Mossavar-Rahmani <i>et al.</i> , US [26]	2019	WHI-OS cohort	61,204	1993–	50–79 yrs	0.0	Questionnaire	• ASB (Diet drinks such as diet coke or diet fruit drinks, with a 12 fl oz can as a reference size) • Frequency	• All-cause mortality: 9,769/61,204	11.9 yrs	0	0	0	0	0	0
Mullee <i>et al.</i> , 10 European countries [27]	2019	EPIC	451,743	1992–2013	50.8 ± 9.8 yrs	28.9	Dietary questionnaire	• Total soft drink • Sugar-sweetened soft drink • Artificially-sweetened soft drink	• All-cause mortality: 41,693/451,743 • CVD mortality: 9,106/451,743	16.4 yrs	0	0	0	0	0	0
Ramme <i>et al.</i> , Sweden [30]	2019	MDCS	24,272	1991–2014	57.6 yrs (44–73 yrs)	38.6	Modified diet history (7-d food diary + FFQ)	• Added sugar • Free sugar	• All-cause mortality: 6,309/24,272 • CVD mortality: 1,952/24,272	19–24 yrs	0	0	0	0	X	0
	2019	NSHDS	24,475	1991–2014	48.6 yrs (36–64 yrs)	46.3	FFQ	• Treats • Toppings • SSB • Added sugar • Free sugar • Treats • Toppings • SSB	• All-cause mortality: 2,881/24,475	19–24 yrs	0	0	0	0	X	0
Anderson <i>et al.</i> , UK [19]	2020	UK general population cohort	161,415	2006–2016	40–69 yrs	-	24-h recall	• SSB • ASB • Fruit or vegetable juice	• All-cause mortality: 2,311/161,415	7 yrs	0	0	0	0	0	X
Zhang <i>et al.</i> , US [33]	2021	NHANES	247,215	1999–2014	≥ 20 yrs	50.4	24-h recall	• Added sugar intake from SSB • SSB • ASB	• All-cause mortality: 3,878/247,215 • Heart disease mortality: 676/247,215	7.9 yrs	0	0	0	0	0	0

ASB, artificially-sweetened beverages; CHD, coronary heart disease; CVD, cardiovascular disease; EPIC, European Prospective Investigation into Cancer and Nutrition; MDCS, Malmo Diet and Cancer Study; NHANES, National Health and Nutrition Examination Survey; NSHDS, Northern Swedish Health and Disease Study; FFQ, food-frequency questionnaire; NHS, Nurses' Health Study; HPFS, Health Professionals Follow-Up Study; REGARDS, Reasons for Geographic and Racial Differences in Stroke; SCS, Singapore Chinese Health Study; SSB, sugar-sweetened beverages; UKWCS, UK Women's Cohort Study; VITAL, The Vitamins and Lifestyle; WHI-OS, Women's Health Initiative Observational Study.

Table 2. Association of added sugar intake with all-cause mortality risk

Exposure	Author (year)	Comparison groups	Adjusted HR	95% CI	P for trend
Free sugar intake (%kcal)	Liu <i>et al.</i> (2018) [24]	Q1: 0.87 (0.00–1.61)	1.00	[Reference]	0.023
		Q2: 2.20 (1.62–2.80)	0.90	0.73–1.12	
		Q3: 3.51 (2.81–4.31)	0.76	0.60–0.95	
		Q4: 5.28 (4.32–6.55)	0.83	0.66–1.04	
		Q5: 8.46 (6.56–54.90)	0.77	0.61–0.98	
	Ramne <i>et al.</i> (2019), MDCS [30]	< 5.0	1.22	1.09–1.37	0.511
		5.0 to < 7.5	1.05	0.96–1.14	
		7.5 to < 10.0	1.00	[Reference]	
		10.0 to < 15.0	1.03	0.97–1.11	
		15.0 to < 20.0	1.04	0.95–1.14	
	Ramne <i>et al.</i> (2019), NSHDS [30]	≥ 20.0	1.26	1.10–1.44	0.487
		< 5.0	1.07	0.94–1.21	
		5.0 to < 7.5	1.06	0.95–1.18	
		7.5 to < 10.0	1.00	[Reference]	
		10.0 to < 15.0	1.06	0.95–1.18	
Added sugar intake (%kcal)	Liu <i>et al.</i> (2018) [24]	Q1: 0.67 (0.00–1.12)	1.00	[Reference]	0.007
		Q2: 1.59 (1.12–2.03)	0.89	0.72–1.10	
		Q3: 2.50 (2.03–3.07)	0.82	0.65–1.02	
		Q4: 3.88 (3.07–4.99)	0.76	0.60–0.95	
		Q5: 6.86 (4.99–54.90)	0.75	0.59–0.95	
	Ramne <i>et al.</i> (2019), MDCS [30]	< 5.0	1.23	1.11–1.35	0.920
		5.0 to < 7.5	1.06	0.98–1.14	
		7.5 to < 10.0	1.00	[Reference]	
		10.0 to < 15.0	1.03	0.96–1.10	
		15.0 to < 20.0	1.04	0.95–1.15	
	Ramne <i>et al.</i> (2019), NSHDS [30]	≥ 20.0	1.30	1.12–1.51	0.664
		< 5.0	1.09	0.97–1.22	
		5.0 to < 7.5	1.05	0.95–1.17	
		7.5 to < 10.0	1.00	[Reference]	
		10.0 to < 15.0	1.04	0.93–1.16	
Added sugar intake (g/1000 kcal)	Tasevska <i>et al.</i> (2014), Men [14]	Q1: 9.2	1.00	[Reference]	0.190
		Q2: 14.7	0.94	0.90–0.97	
		Q3: 21.0	0.94	0.90–0.97	
		Q4: 29.4	0.92	0.89–0.96	
		Q5: 47.0	0.95	0.91–0.99	
	Tasevska <i>et al.</i> (2014), Women [14]	Q1: 10.1	1.00	[Reference]	0.300
		Q2: 15.1	0.96	0.91–1.01	
		Q3: 20.6	0.93	0.89–0.98	
		Q4: 28.6	0.97	0.92–1.02	
		Q5: 45.4	1.00	0.95–1.06	

CI, confidence interval; HR, hazard ratio; MDCS, Malmo Diet and Cancer Study; NSHDS, Northern Swedish Health, and Disease Study.

Added sugar intake, SSB consumption, and CVD mortality

Tables 4 and 5 lists the associations of added sugar intake and SSB consumption with the risk of CVD mortality, respectively. Among the 4 studies examining the association between added sugar intake and CVD mortality [13,14,23,30], 3 studies did not find any significant associations [14,23,30], but one study based on the NHANES data reported that a higher intake of added sugar was associated with an increased risk of CVD mortality in US adults (Q5 vs. Q1: RR = 2.03, 95% CI = 1.26–3.27, P for trend = 0.004) [13].

Based on a review of 8 studies on SSB consumption and CVD mortality [14,20,21,23,25,27,28,33], 5 studies did not report significant associations of SSB consumption or added sugar intake from

Added sugar intake and mortality

Table 3. Association of SSB consumption with all-cause mortality risk

Exposure	Author (year)	Comparison groups	Adjusted HR	95% CI	P for trend	
SSB						
SSB (serving/d)	Anderson <i>et al.</i> (2020) [19]	0	1.00	[Reference]	< 0.001	
		1	1.06	0.96–1.16		
		> 1 to 2	1.33	1.10–1.60		
		> 2	1.84	1.42–2.37		
Sugar-sweetened drinks (servings/wk)	Barrington and White (2016) [20]	0–0.1	1.00	[Reference]	< 0.001	
		0.2–0.4	1.08	0.98–1.19		
		0.5–2.7	1.10	1.01–1.19		
		≥ 2.8	1.19	1.08–1.30		
SSB (serving/d)	Collin <i>et al.</i> (2019) [21]	Each additional 12 oz	1.06	0.96–1.16	-	
SSB (Frequency consumption of a standard portion)	Malik <i>et al.</i> (2019), Pooled [25]	< 1 time/mo	1.00	[Reference]	< 0.001	
		1 to 4 times/mo	1.01	0.98–1.04		
		2 to 6 times/wk	1.06	1.03–1.09		
		1 to < 2 times/d	1.14	1.09–1.19		
	Malik <i>et al.</i> (2019), HPFS [25]	≥ 2 times/d	1.21	1.13–1.28	< 0.001	
		Per 1 serving/d	1.07	1.05–1.09		
		< 1 time/mo	1.00	[Reference]		
		1 to 4 times/mo	0.95	0.91–1.00		
	Malik <i>et al.</i> (2019), NHS [25]	2 to 6 times/wk	1.06	1.01–1.10	< 0.001	
		1 to < 2 times/d	1.14	1.05–1.23		
		≥ 2 times/d	1.12	1.00–1.26		
		Per 1 serving/d	1.07	1.03–1.10		
	Sugar-sweetened soft drink (1 glass = 250 mL)	Mullee <i>et al.</i> (2019), Total [27]	< 1 glass/mo	1.00	[Reference]	0.004
			1 to 4 glasses/mo	0.94	0.91–0.98	
			> 1 to 6 glasses/wk	0.96	0.93–1.00	
			1 to < 2 glasses/d	1.08	1.01–1.16	
Mullee <i>et al.</i> (2019), Men [27]		≥ 2 glasses/d	1.08	1.01–1.16	0.050	
		< 1 glass/mo	1.00	[Reference]		
		1 to 4 glasses/mo	0.96	0.91–1.01		
		> 1 to 6 glasses/wk	0.96	0.91–1.01		
Mullee <i>et al.</i> (2019), Women [27]		1 to < 2 glasses/d	1.03	0.93–1.13	0.040	
		≥ 2 glasses/d	1.09	0.99–1.20		
		< 1 glass/mo	1.00	[Reference]		
		1 to 4 glasses/mo	0.93	0.89–0.98		
Soft drink (frequency)	Odegaard <i>et al.</i> (2016) [28]	> 1 to 6 glasses/wk	0.97	0.93–1.01	0.110	
		1 to < 2 glasses/d	1.14	1.04–1.25		
		≥ 2 glasses/d	1.06	0.95–1.18		
		None	1.00	[Reference]		
		Monthly (any to < 1 time/wk)	0.93	0.87–1.01		
Cola with sugar (cans/wk)	Paganini-Hill <i>et al.</i> (2007) [29]	1 time/wk	0.94	0.84–1.04	-	
		2–6 times/wk	0.96	0.87–1.05		
		≥ 1 time/d	0.92	0.81–1.04		
Other soft drinks with sugar (cans/wk)	Paganini-Hill <i>et al.</i> (2007) [29]	None	1.00	[Reference]	-	
		≤ 1	0.95	0.89–1.01		
		> 1	1.02	0.92–1.13		
SSB (servings/wk)	Ramne <i>et al.</i> (2019), MDCS [30]	None	1.00	[Reference]	0.009	
		≤ 1	1.00	0.95–1.05		
		> 1	1.03	0.92–1.16		
		> 1–3	0.97	0.90–1.03		
		> 3–5	1.00	0.91–1.10		
> 5–8	1.09	0.98–1.21				
> 8	1.14	1.03–1.26				

(continued to the next page)

Table 3. (Continued) Association of SSB consumption with all-cause mortality risk

Exposure	Author (year)	Comparison groups	Adjusted HR	95% CI	P for trend
Soft drinks (g/d)	Ramne <i>et al.</i> (2019), NSHDS [30]	≤ 1	1.00	[Reference]	0.177
		> 1–3	0.96	0.88–1.05	
		> 3–5	0.97	0.85–1.10	
		> 5–8	1.03	0.91–1.10	
		> 8	1.10	0.90–1.35	
	Sluik <i>et al.</i> (2014) [31]	Per 10 g in diabetic	1.04	0.99–1.09	-
SSB (serving/d)	Zhang <i>et al.</i> (2021) [33]	0	1.00	[Reference]	-
		> 0 to < 1	1.09	0.98–1.21	
		1 to < 2	1.01	0.89–1.16	
		≥ 2	1.17	0.98–1.38	
		Each additional serving	1.05	1.01–1.09	
Added sugar intake from beverages					
Added sugar intake from beverages (g/1000 kcal)	Tasevska <i>et al.</i> (2014), Men [14]	Q1: 0.12	1.00	[Reference]	0.340
		Q5: 30.9	1.00	0.96–1.04	
Added sugar intake from SSB (g/1000 kcal)	Tasevska <i>et al.</i> (2014), Women [14]	Q1: 0.08	1.00	[Reference]	< 0.001
		Q5: 26.9	1.09	1.03–1.15	
	Zhang <i>et al.</i> (2021) [33]	Q1: 0	1.00	[Reference]	
		Q2: 5.59 (< 9.0)	1.05	0.92–1.20	
		Q3: 13.83 (9.0 to 19.5)	1.02	0.90–1.17	
Added sugar intake from SSB (10 g/1,000 kcal)	Zhang <i>et al.</i> (2021) [33]	Q4: 26.52 (19.6 to 35.3)	1.09	0.96–1.23	
		Q5: 50.12 (> 35.3)	1.22	1.05–1.42	
		Each additional 10 g/1,000 kcal	1.03	1.01–1.05	-

CI, confidence interval; HR, hazard ratio; MDCS, Malmo Diet and Cancer Study; NSHDS, Northern Swedish Health, and Disease Study; SSB, sugar-sweetened beverages.

SSB with CVD mortality [20,21,23,27,28]. Three US studies consistently observed that SSB intake was positively associated with CVD mortality [14,25,33]. Tasevska *et al.* [14] observed this positive association only in women, not in men.

Artificially-sweetened beverages (ASB) and juice consumption and mortality

The association of ASB consumption with the all-cause or CVD mortality was examined in seven studies (data not shown). The 6 studies reported positive associations between ASB intake and all-cause mortality [19,25–27,29,32], but 1 study did not observe a significant association [33]. Regarding the CVD mortality, 3 studies showed that a higher intake of ASB increased the risk of mortality [25,27,32], but Zhang *et al.* [33] did not observe an increase.

Four studies investigating relationships between juice consumption and all-cause mortality reported inconsistent findings: 1 study found a positive association of fruit juice [21]. Another study found no association of fruit and vegetable juices [28], and 2 studies found an inverse association (1 study addressed fruit and vegetable juice, and the other study had no information on the type of juice) [19,31]. Three studies on juice intake and CVD mortality consistently showed no significant associations, but each study defined juice consumption differently in terms of the type of juice, or the unit of the amount consumed [21,22,28].

DISCUSSION

This paper systematically reviewed studies reporting the effects of added sugar according to the source on mortality as a long-term and summary health outcome, focusing on the added sugar intake and SSB. This study observed a similar positive association of sugar intake with mortality to the studies analyzing SSB and morbidity and the intake range with low

Table 4. Association of added sugar intake with cardiovascular disease mortality risk

Exposure	Author (year)	Comparison group	Adjusted HR	95% CI	P for trend
Free sugar intake (%kcal)	Ramne <i>et al.</i> (2019), MDCS [30]	< 5.0	1.24	1.01–1.53	0.562
		5.0 to < 7.5	1.15	0.99–1.33	
		7.5 to < 10.0	1.00	[Reference]	
		10.0 to < 15.0	1.02	0.90–1.15	
		15.0 to < 20.0	0.98	0.83–1.15	
		≥ 20.0	1.31	1.03–1.66	
Added sugar intake (%kcal)	Liu <i>et al.</i> (2018) [23]	Q1: 0.67 (0.00–1.12)	1.00	[Reference]	0.055
		Q2: 1.59 (1.12–2.03)	0.69	0.27–1.73	
		Q3: 2.50 (2.03–3.07)	0.32	0.10–1.02	
		Q4: 3.88 (3.07–4.99)	0.48	0.16–1.47	
		Q5: 6.86 (4.99–54.9)	0.33	0.08–1.43	
	Ramne <i>et al.</i> (2019), MDCS [30]	< 5.0	1.22	1.02–1.47	0.744
		5.0 to < 7.5	1.09	0.95–1.25	
		7.5 to < 10.0	1.00	[Reference]	
		10.0 to < 15.0	1.01	0.89–1.13	
		15.0 to < 20.0	0.97	0.82–1.16	
	Yang <i>et al.</i> (2014) [13]	Q1: 7.4 (0.0 to < 9.6)	1.00	[Reference]	0.004
		Q2: 11.4 (9.6 to < 13.1)	1.07	1.02–1.12	
		Q3: 14.8 (13.1 to < 16.7)	1.18	1.06–1.31	
		Q4: 18.7 (16.7 to < 21.3)	1.38	1.11–1.70	
		Q5: 25.2 (≥ 21.3)	2.03	1.26–3.27	
Yang <i>et al.</i> (2014) [13]	5.0 (0.0 to < 10.0)	1.00	[Reference]	0.004	
	17.5 (10.0 to < 25.0)	1.30	1.09–1.55		
	28.7 (≥ 25.0)	2.75	1.40–5.42		
Added sugar intake (g/1,000 kcal)	Tasevska <i>et al.</i> (2014), Men [14]	Q1: 9.2	1.00	[Reference]	0.070
		Q2: 14.7	0.91	0.85–0.98	
		Q3: 21.0	0.87	0.82–0.95	
		Q4: 29.4	0.87	0.81–0.94	
		Q5: 47.0	0.91	0.84–0.98	
	Tasevska <i>et al.</i> (2014), Women [14]	Q1: 10.1	1.00	[Reference]	0.940
		Q2: 15.1	0.94	0.84–1.04	
		Q3: 20.6	0.82	0.73–0.92	
		Q4: 28.6	0.94	0.84–1.05	
		Q5: 45.4	0.96	0.86–1.08	

CI, confidence interval; HR, hazard ratio; MDCS, Malmo Diet and Cancer Study.

mortality risk. In addition, the difference in risk according to the sugar source: sugary food and beverage did not increase the mortality risk when it contained sufficient nutrients other than sugar.

The intake level of added sugar was associated with mortality in all three studies [14,24,30] that analyzed the association of the energy contribution of added sugar intake with all-cause mortality. The above three studies showed different patterns of risk according to the intake level because an increase in mortality was observed at intake levels less than 5% of the total energy, even when the intake of added sugar in the target group was low. A similar trend was observed in studies on CVD mortality. The CVD risk decreased with increasing intake level in the study that the reference intake interval was less than 5% of the total energy. Tasevska *et al.* [14] suggested an association of fructose, but not added sugars, with the mortality risk, while it may represent the influence of SSB as the main added fructose source.

Among the studies that conducted all-cause mortality as an outcome, 11 studies analyzed SSB exposure with different units. Nine studies were measured by the serving unit, among which five studies showed that all-cause mortality increased with increasing intake of SSB.

Added sugar intake and mortality

Table 5. Association of SSB consumption with cardiovascular disease mortality risk

Exposure	Author (year)	Comparison groups	Adjusted HR	95% CI	P for trend
SSB					
Sugar-sweetened drinks (servings/wk)	Barrington and White (2016) [20]	0.0–0.1	1.00	[Reference]	0.450
		0.2–0.4	0.92	0.75–1.12	
		0.5–2.7	1.06	0.89–1.25	
		≥ 2.8	1.05	0.87–1.27	
		Each additional 12 oz	1.11	0.90–1.39	
SSB (serving/d)	Collin <i>et al.</i> (2019) [21]	Each additional 12 oz	1.11	0.90–1.39	-
SSB (Frequency consumption of a standard portion)	Malik <i>et al.</i> (2019), Pooled [25]	< 1 times/mo	1.00	[Reference]	< 0.001
		1 to 4 times/mo	1.06	1.00–1.12	
		2 to 6 times/wk	1.10	1.04–1.17	
		1 to < 2 times/d	1.19	1.08–1.31	
		≥ 2 times/d	1.31	1.15–1.50	
	Malik <i>et al.</i> (2019), HPFS [25]	Per 1 serving/d	1.10	1.06–1.14	0.020
		< 1/mo	1.00	[Reference]	
		1 to 4/mo	1.04	0.95–1.14	
		2 to 6/wk	1.08	1.00–1.18	
		1 to < 2/d	1.17	1.01–1.35	
	Malik <i>et al.</i> (2019), NHS [25]	≥ 2/d	1.19	0.95–1.49	< 0.001
		Per 1 serving/d	1.07	1.01–1.14	
		< 1/mo	1.00	[Reference]	
		1 to 4/mo	1.07	0.99–1.16	
		2 to 6/wk	1.10	1.01–1.20	
Sugar-sweetened soft drink (1 glass = 250 mL)	Mullee <i>et al.</i> (2019), Total [27]	1 to < 2/d	1.21	1.06–1.37	0.160
		≥ 2/d	1.37	1.16–1.62	
		Per 1 serving/d	1.11	1.06–1.16	
		< 1 glass/mo	1.00	[Reference]	
		1 to 4 glasses/mo	0.97	0.90–1.05	
	Mullee <i>et al.</i> (2019), Men [27]	> 1 to 6 glasses/wk	0.96	0.90–1.04	0.430
		1 to < 2 glasses/d	1.06	0.92–1.22	
		≥ 2 glasses/d	1.11	0.95–1.30	
		< 1 glass/mo	1.00	[Reference]	
		1 to 4 glasses/mo	0.99	0.89–1.10	
	Mullee <i>et al.</i> (2019), Women [27]	> 1 to 6 glasses/wk	0.94	0.85–1.04	0.200
		1 to < 2 glasses/d	0.98	0.80–1.20	
		≥ 2 glasses/d	1.11	0.91–1.35	
		< 1 glass/mo	1.00	[Reference]	
		1 to 4 glasses/mo	0.97	0.87–1.07	
Soft drink (frequency)	Odegaard <i>et al.</i> (2016) [28]	> 1 to 6 glasses/wk	0.99	0.89–1.09	0.540
		1 to < 2 glasses/d	1.15	0.94–1.40	
		≥ 2 glasses/d	1.11	0.86–1.43	
		None	1.00	[Reference]	
		Monthly	0.96	0.84–1.09	
SSB (serving/d)	Zhang <i>et al.</i> (2021) [33]	1/wk	1.08	0.91–1.30	-
		2–6/wk	1.08	0.92–1.27	
		≥ 1/d	1.03	0.83–1.30	
		0	1.00	[Reference]	
		> 0 to < 1	1.02	0.81–1.28	
Added sugar intake from beverage	Liu <i>et al.</i> (2018) [23]	1 to < 2	1.38	1.05–1.81	-
		≥ 2	1.52	1.03–2.25	
		Each additional serving	1.11	1.03–1.21	
		Q1: 0.00 (0.00–0.05)	1.00	[Reference]	
		Q2: 0.17 (0.05–0.31)	1.10	0.43–2.83	
	Tasevska <i>et al.</i> (2014), Men [14]	Q3: 0.52 (0.31–0.83)	0.44	0.14–1.40	0.300
		Q4: 1.34 (0.83–2.24)	0.41	0.14–1.25	
		Q5: 4.15 (2.24–53.1)	0.55	0.15–2.07	
		Q1: 0.12	1.00	[Reference]	
		Q5: 30.9	1.01	0.94–1.09	
	Tasevska <i>et al.</i> (2014), Women [14]	Q1: 0.08	1.00	[Reference]	0.020
		Q5: 26.9	1.13	1.01–1.26	
		Q1: 0.0	1.00	[Reference]	
	Zhang <i>et al.</i> (2021) [33]	Q2: 5.59 (< 9.0)	0.91	0.65–1.25	-
		Q3: 13.83 (9.0 to 19.5)	1.15	0.89–1.47	
Q4: 26.52 (19.6 to 35.3)		1.32	0.98–1.76		
Q5: 50.12 (> 35.3)		1.45	1.06–1.97		
Each additional 10 g/1,000 kcal		1.08	1.03–1.13		
Added sugar intake from SSB (10 g/1,000 kcal)	Zhang <i>et al.</i> (2021) [33]	Each additional 10 g/1,000 kcal	1.08	1.03–1.13	-

CI, confidence interval; HR, hazard ratio; SSB, sugar-sweetened beverages.

Two studies analyzed the dose-response relationship and reported that the mortality risk increased 5% [33] and 7% [25] each for one serving per day. In one study that analyzed according to gender, a statistically significant increase in mortality risk was observed only in women [27]. In addition, 1 study that analyzed the results from 2 cohorts of different age groups [30] reported a significant association only in younger adults. Further studies on the difference in the effects according to age and gender are needed.

In 2 studies that analyzed the risk by the quintile according to the exposure level of SSB added sugar (g/1,000 kcal), the mortality was significantly higher in the top quintile. Among them, 1 study [14] reported statistical significance only in women. When the median of the top quintile was converted to %kcal, they were 10.76% [4] and 20% [33]. For every 4% increase in SSB intake, the mortality risk increased by 3% [19]. On the other hand, the consumption of ASB increased the mortality risk. One study [33] reported a decreased risk when the intake level was less than a serving per day. Therefore, it is necessary to pay attention to the dietary behavior of subjects who consume a considerable amount of beverages, including SSB when determining the correlation with health outcomes.

Liquid sources, such as SSB and ASB, increased the mortality risk, whereas solid sources decreased the risk. In addition, one study [31] analyzed the mortality risk by dividing solid sources into “sugar and confectionery” and “cakes and cookies”. They reported a 5% increased risk from “sugar and confectionery” for every 10 g increase in intake and a 3% decrease for “cakes and cookies”. These results may indicate that sugary food containing sufficient nutrients other than sugar does not increase the mortality risk.

Some studies [24,30] also suggested the mortality risk according to the intake level of free sugar. These results were similar to those of added sugar. The recommendations for reducing sugar intake address the free sugar or added sugar as the target of restrictions, but the food sources are somewhat different between free sugar and added sugar. Juice with a high sugar content is included as a free sugar food defined by the WHO [34] because it is not a whole food but a processed product with a high sugar content from fruit. On the other hand, the source of added sugar suggested by the US Department of Agriculture (USDA) [35] and US Food and Drug Administration [36] did not include 100% fruit juice. Although the USDA [37] recommended eating whole fruit rather than fruit juice in the dietary guidelines for Americans, fruit juice is also included in the fruit group.

The 4 studies analyzed according to the exposure level of juice reported inconsistent results (2 decreased, 1 increased, and 1 not significant). The large difference in the level of exposure made it difficult to conclude by summarizing the trends. According to a study on the association of whole fruit and fruit juice with the prevalence of obesity using the KNHANES data, the obesity and metabolic syndrome abnormalities were negatively associated with the whole fruit intake, but the relationship with fruit juice intake was difficult to determine.

In the Korean food guide, fruit juice is classified as a fruit group and is recommended as a healthy choice. The 2015 KDRIs referred to the WHO's recommended free sugar level when setting the added sugar recommendation. On the other hand, the term free sugar was not used to consider whether the term and its definition could be adequately communicated to the general public. Moreover, there is a lack of evidence to validate the recommendation because of the absence of a database for either added sugar or free sugar content in foods to calculate intake.

The total sugar was recently included in the standard food composition table. The total sugar intake for three years since 2016 was first reported at the national level using the 7th KNHANES data [38]. The total sugar intake for three years during the 7th survey period has been gradually decreasing, and the average daily sugar intake of Koreans was 60 g, which is approximately 12% of the average energy intake (1,988 kcal) in 2018. The intake level of added sugar might be much lower. When the WHO [34] set the recent sugar intake guidelines, the intake of free sugar was limited to less than 10% of the energy intake level, while a comment was added to ensure that the current consumption levels do not increase further rather than applying the guideline in countries where the consumption levels of sugar are currently low.

Tasevska *et al.* [14] reported the same trend as added sugar when the total sugar was analyzed as exposure. The mortality increased with increasing sugar intake, while men with a relatively lower intake than women had higher mortality, even in the lowest intake quintile (median 13.4% of energy) than in medium intake level. Although it is necessary to examine whether there is a gender-specific difference or just depending on the intake level, all-cause mortality increased with increasing added sugar intake from SSB in women, and the all-cause mortality from the solids source decreased in men. In addition, a Japanese study [39] reported that the intake level (median of 1st quintile is 4.4% of energy) was lower than that of Tasevska *et al.* [14]. On the other hand, the all-cause and CVD mortality increased with increasing total sugar intake showing the lowest risk in the lowest quintile.

After a systematic review of the association with mortality and additional evidence on morbidity since 2015 considering revision, the 2020 KDRI decided to maintain the existing recommendations because of insufficient evidence and the absence of a necessity for the revision [40]. The differences in intake levels that increase the mortality in each study may originate from differences in the method of estimating sugar intake, but may also be from the differences between countries. In addition, an accurate evaluation of decreasing trends in the total sugar intake in Koreans is required because fruit intake has been declining recently [38]. Therefore, to set sugar intake guidelines, more evidence will be needed on the comprehensive consumption data of Koreans and the consequent morbidity and mortality.

In conclusion, this systematic review revealed an appropriate range of added sugar intake to lower mortality. Although there were differences according to the study, low mortality was observed over the intake range of added sugar from 5% to 20% of energy. Accordingly, a 'U-' or 'J'-shaped relationship was estimated, but more evidence will be needed to establish a dose-response relationship. In addition, the mortality risk from sugary food and beverages was not increased when they contained sufficient nutrients other than sugar. Thus, appropriate recommendations on the total sugar intake and restricting sugary foods that are not nutritious may be a good strategy. Although the 2020 KDRI maintained the existing recommendations, more studies (i.e., comprehensive consumption and dietary behavior data of various sugary foods) will be needed to complement the current sugar guidelines and set adequate recommendations for Koreans. Academia should establish a statement toward fruit juice as a free sugar source and provide proper communication to the public for added sugar or free sugar to assist in healthy food choices.

REFERENCES

1. Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ* 2012;346:e7492.
[PUBMED](#) | [CROSSREF](#)
2. Te Morenga LA, Howatson AJ, Jones RM, Mann J. Dietary sugars and cardiometabolic risk: systematic review and meta-analyses of randomized controlled trials of the effects on blood pressure and lipids. *Am J Clin Nutr* 2014;100:65-79.
[PUBMED](#) | [CROSSREF](#)
3. Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *Am J Clin Nutr* 2013;98:1084-102.
[PUBMED](#) | [CROSSREF](#)
4. Malik VS, Popkin BM, Bray GA, Després JP, Willett WC, Hu FB. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. *Diabetes Care* 2010;33:2477-83.
[PUBMED](#) | [CROSSREF](#)
5. Ministry of Health and Welfare, Korea Disease Control and Prevention Agency. Korea Health Statistics 2019: Korea National Health and Nutrition Examination Survey (KNHANES VIII-1). Cheongju: Korea Disease Control and Prevention Agency; 2020.
6. Lim H, Lee HJ, Choue R, Wang Y. Trends in fast-food and sugar-sweetened beverage consumption and their association with social environmental status in south Korea. *J Acad Nutr Diet* 2018;118:1228-1236.e1.
[PUBMED](#) | [CROSSREF](#)
7. Chung S, Ha K, Lee HS, Kim CI, Joung H, Paik HY, Song Y. Soft drink consumption is positively associated with metabolic syndrome risk factors only in Korean women: data from the 2007-2011 Korea National Health and Nutrition Examination Survey. *Metabolism* 2015;64:1477-84.
[PUBMED](#) | [CROSSREF](#)
8. Kang Y, Kim J. Soft drink consumption is associated with increased incidence of the metabolic syndrome only in women. *Br J Nutr* 2017;117:315-24.
[PUBMED](#) | [CROSSREF](#)
9. Shin S, Kim SA, Ha J, Lim K. Sugar-sweetened beverage consumption in relation to obesity and metabolic syndrome among Korean adults: a cross-sectional study from the 2012–2016 Korean National Health and Nutrition Examination Survey (KNHANES). *Nutrients* 2018;10:1467.
[CROSSREF](#)
10. Kwak JH, Jo G, Chung HK, Shin MJ. Association between sugar-sweetened beverage consumption and incident hypertension in Korean adults: a prospective study. *Eur J Nutr* 2019;58:1009-17.
[PUBMED](#) | [CROSSREF](#)
11. Ha K, Chung S, Lee HS, Kim CI, Joung H, Paik HY, Song Y. Association of dietary sugars and sugar-sweetened beverage intake with obesity in Korean children and adolescents. *Nutrients* 2016;8:31.
[PUBMED](#) | [CROSSREF](#)
12. Seo EH, Kim H, Kwon O. Association between total sugar intake and metabolic syndrome in middle-aged Korean men and women. *Nutrients* 2019;11:2042.
[PUBMED](#) | [CROSSREF](#)
13. Yang Q, Zhang Z, Gregg EW, Flanders WD, Merritt R, Hu FB. Added sugar intake and cardiovascular diseases mortality among US adults. *JAMA Intern Med* 2014;174:516-24.
[PUBMED](#) | [CROSSREF](#)
14. Tasevska N, Park Y, Jiao L, Hollenbeck A, Subar AF, Potischman N. Sugars and risk of mortality in the NIH-AARP Diet and Health Study. *Am J Clin Nutr* 2014;99:1077-88.
[PUBMED](#) | [CROSSREF](#)
15. World Health Organization (WHO). Diet, Nutrition and the Prevention of Chronic Diseases: Report of a WHO Study Group. WHO Technical Report Series 797. Geneva: WHO; 1990.
16. World Health Organization (WHO). Diet, Nutrition and the Prevention of Chronic Diseases: Report of a Joint WHO/FAO Expert Consultation. WHO Technical Report Series, No 916. Geneva: WHO; 2003.
17. Moynihan PJ, Kelly SA. Effect on caries of restricting sugars intake: systematic review to inform WHO guidelines. *J Dent Res* 2014;93:8-18.
[PUBMED](#) | [CROSSREF](#)
18. World Health Organization (WHO). Guideline: Sugars Intake for Adults and Children. Geneva: WHO; 2015.
19. Anderson JJ, Gray SR, Welsh P, Mackay DF, Celis-Morales CA, Lyall DM, Forbes J, Sattar N, Gill JM, Pell JP. The associations of sugar-sweetened, artificially sweetened and naturally sweet juices with all-cause mortality in 198,285 UK Biobank participants: a prospective cohort study. *BMC Med* 2020;18:97.
[PUBMED](#) | [CROSSREF](#)

20. Barrington WE, White E. Mortality outcomes associated with intake of fast-food items and sugar-sweetened drinks among older adults in the Vitamins and Lifestyle (VITAL) study. *Public Health Nutr* 2016;19:3319-26.
[PUBMED](#) | [CROSSREF](#)
21. Collin LJ, Judd S, Safford M, Vaccarino V, Welsh JA. Association of sugary beverage consumption with mortality risk in US adults: a secondary analysis of data from the REGARDS study. *JAMA Netw Open* 2019;2:e193121.
[PUBMED](#) | [CROSSREF](#)
22. Lai HT, Threapleton DE, Day AJ, Williamson G, Cade JE, Burley VJ. Fruit intake and cardiovascular disease mortality in the UK Women's Cohort Study. *Eur J Epidemiol* 2015;30:1035-48.
[PUBMED](#) | [CROSSREF](#)
23. Liu ZM, Tse LA, Chan D, Wong C, Wong SY. Dietary sugar intake was associated with increased body fatness but decreased cardiovascular mortality in Chinese elderly: an 11-year prospective study of Mr and Ms OS of Hong Kong. *Int J Obes* 2018;42:808-16.
[PUBMED](#) | [CROSSREF](#)
24. Liu ZM, Tse SL, Chen B, Chan D, Wong C, Woo J, Wong SY. Dietary sugar intake does not pose any risk of bone loss and non-traumatic fracture and is associated with a decrease in all-cause mortality among Chinese elderly: Finding from an 11-year longitudinal study of Mr. and Ms. OS Hong Kong. *Bone* 2018;116:154-61.
[PUBMED](#) | [CROSSREF](#)
25. Malik VS, Li Y, Pan A, De Koning L, Schernhammer E, Willett WC, Hu FB. Long-term consumption of sugar-sweetened and artificially sweetened beverages and risk of mortality in US adults. *Circulation* 2019;139:2113-25.
[PUBMED](#) | [CROSSREF](#)
26. Mossavar-Rahmani Y, Kamensky V, Manson JE, Silver B, Rapp SR, Haring B, Beresford SA, Snetselaar L, Wassertheil-Smoller S. Artificially sweetened beverages and stroke, coronary heart disease, and all-cause mortality in the Women's Health Initiative. *Stroke* 2019;50:555-62.
[PUBMED](#) | [CROSSREF](#)
27. Mullee A, Romaguera D, Pearson-Stuttard J, Viallon V, Stepien M, Freisling H, Fagherazzi G, Mancini FR, Boutron-Ruault MC, Kühn T, et al. Association between soft drink consumption and mortality in 10 European countries. *JAMA Intern Med* 2019;179:1479-90.
[PUBMED](#) | [CROSSREF](#)
28. Odegaard AO, Koh WP, Yuan JM, Pereira MA. Beverage habits and mortality in Chinese adults. *J Nutr* 2015;145:595-604.
[PUBMED](#) | [CROSSREF](#)
29. Paganini-Hill A, Kawas CH, Corrada MM. Non-alcoholic beverage and caffeine consumption and mortality: the Leisure World Cohort Study. *Prev Med* 2007;44:305-10.
[PUBMED](#) | [CROSSREF](#)
30. Ramne S, Alves Dias J, González-Padilla E, Olsson K, Lindahl B, Engström G, Ericson U, Johansson I, Sonestedt E. Association between added sugar intake and mortality is nonlinear and dependent on sugar source in 2 Swedish population-based prospective cohorts. *Am J Clin Nutr* 2019;109:411-23.
[PUBMED](#) | [CROSSREF](#)
31. Sluik D, Boeing H, Li K, Kaaks R, Johnsen NF, Tjønneland A, Arriola L, Barricarte A, Masala G, Grioni S, et al. Lifestyle factors and mortality risk in individuals with diabetes mellitus: are the associations different from those in individuals without diabetes? *Diabetologia* 2014;57:63-72.
[PUBMED](#) | [CROSSREF](#)
32. Vyas A, Rubenstein L, Robinson J, Seguin RA, Vitolins MZ, Kazlauskaitė R, Shikany JM, Johnson KC, Snetselaar L, Wallace R. Diet drink consumption and the risk of cardiovascular events: a report from the Women's Health Initiative. *J Gen Intern Med* 2015;30:462-8.
[PUBMED](#) | [CROSSREF](#)
33. Zhang YB, Chen JX, Jiang YW, Xia PF, Pan A. Association of sugar-sweetened beverage and artificially sweetened beverage intakes with mortality: an analysis of US National Health and Nutrition Examination Survey. *Eur J Nutr* 2021;60:1945-55.
[PUBMED](#) | [CROSSREF](#)
34. World Health Organization (WHO). Information Note about Intake of Sugars Recommended in the WHO Guideline for Adults and Children (No. WHO/NMH/NHD/15.3). Geneva: WHO; 2015.
35. U.S. Department of Agriculture (USDA). What Are "Added Sugars"? Washington, D.C.: USDA; 2019 [cited 2021 September 7]. Available from: <https://ask.usda.gov/s/article/What-are-added-sugars>.

36. Office of the Federal Register (OFR) of the National Archives and Records Administration (NARA) and the U.S. Government Publishing Office (GPO). Electronic Code of Federal Regulations (eCFR). OFR/GPO; 2021 [cited 2021 September 7]. Available from: <https://www.ecfr.gov/current/title-21/chapter-I/subchapter-B/part-101>.
37. U.S. Department of Agriculture (USDA). How does United States Department of Agriculture Classify Fruit Juice and Fruit Drinks? Washington, D.C.: USDA; 2019 [cited 2021 September 7]. Available from: <https://ask.usda.gov/s/article/How-does-United-States-Department-of-Agriculture-classify-fruit-juice-and-fruit-drinks>.
38. Ministry of Health and Welfare, Korea Centers for Disease Control and Prevention. Korea Health Statistics 2018: Korea National Health and Nutrition Examination Survey (KNHANES VII-3). Cheongju: Korea Centers for Disease Control and Prevention; 2019.
39. Nagata C, Wada K, Yamakawa M, Konishi K, Goto Y, Koda S, Mizuta F, Uji T. Intake of starch and sugars and total and cause-specific mortality in a Japanese community: the Takayama Study. *Br J Nutr* 2019;122:820-8.
[PUBMED](#) | [CROSSREF](#)
40. Korean Nutrition Society. Dietary Reference Intakes for Koreans 2020. Seoul: Korean Nutrition Society; 2020.