

SPACES OF BMO TYPE

YOUNG JA PARK

ABSTRACT. It is presented a Banach space of functions of bounded mean oscillation *BMO* type.

1. Introduction

The space of functions of bounded mean oscillation, or *BMO*, naturally arises as a class of functions whose deviation from their means over cubes is bounded. In fact, the classical *BMO*-norm $\|f\|_{BMO}$ for the equivalent class of a locally integrable function f on \mathbb{R}^d ($f \in L^1_{loc}(\mathbb{R}^d)$) is defined as

$$(1.1) \quad \|f\|_{BMO} := \sup_Q \frac{1}{|Q|} \int_Q |f(x) - f_Q| dx$$

for every cube $Q \subset \mathbb{R}^d$ whose sides are parallel to the axes and

$$f_Q := \frac{1}{|Q|} \int_Q f(x) dx.$$

The space *BMO* first appeared in the work of John and Nirenberg [1] in the context of nonlinear partial differential equations that emerge in the study of minimal surfaces.

Even though the Lebesgue space L^∞ functions have the same property, there exist unbounded functions with bounded mean oscillation. Such functions are slowly growing, and typically have at most logarithmic blow-up. The space *BMO* shares similar properties with the space L^∞ , and it often serves as a substitute for it. For instance, classical singular integrals do not map L^∞ to L^∞ but L^∞ to *BMO*. In many instances the interpolation between L^p and *BMO* works just as well

Received February 25, 2022; Accepted May 01, 2022.

2010 Mathematics Subject Classification: 46E30, 42B35, 32A55.

Key words and phrases: Bounded mean oscillation, function space, Hölder inequality, John-Nirenberg inequality.

This research was supported by the Academic Research Fund of Hoseo University in 2021(20210425).

between L^p and L^∞ . Indeed, the role of the space BMO is deeper and more far-reaching than that [6]. This space crucially arises in many situations in analysis, such as in the characterization of the L^2 -boundedness of non-convolution singular integral operators with standard kernels.

Recently, we have built up a new function space in order to generalize the classical Lebesgue spaces [3, 4, 5]. The motivation of this research stems from taking a close look at the L^p -norm: $\|f\|_{L^p} = (\int_X |f(x)|^p d\mu)^{1/p}$ of the Lebesgue spaces $L^p(X)$, $1 \leq p < \infty$. It can be rewritten as

$$(1.2) \quad \|f\|_{L^p} := \alpha^{-1} \left(\int_X \alpha(|f(x)|) d\mu \right), \quad f \in L^p(X)$$

with the base function α as

$$\alpha(x) := x^p.$$

By virtue of the John-Nirenberg inequality, it is well-known that the classical BMO -norm (1.1) is equivalent to the L^p characterization of BMO -norm $\|\cdot\|_{BMO_p}$ defined by

$$(1.3) \quad \|f\|_{BMO_p} := \left(\sup_Q \frac{1}{|Q|} \int_Q |f(x) - f_Q|^p dx \right)^{\frac{1}{p}}, \quad (f \in L^1_{loc}(\mathbb{R}^d))$$

for $1 < p < \infty$.

In the same line of our research we introduce a functional

$$(1.4) \quad \|f\|_{BMO_\alpha} := \alpha^{-1} \left(\sup_Q \frac{1}{|Q|} \int_Q \alpha(|f(x) - f_Q|) dx \right), \quad (f \in L^1_{loc}(\mathbb{R}^d))$$

for an appropriate base function α . The main point of this report is to present sufficient conditions of base functions α such that $\|\cdot\|_{BMO_\alpha}$ forms a (quasi-)norm, so it constitutes a natural Banach space BMO_α .

The base functions α which we have developed include base functions of the form $\alpha(x) = x^p$, and we designate the base functions α to achieve the Minkowski type triangle inequality. This research was inspired by [2].

2. The main theorem and arguments

We have been developed appropriate base functions that permit the Hölder type inequality. In this section, we briefly introduce the concepts

of admissible base functions - the details can be found in [3, 4]. The notions presented here are modified versions without essential differences. In the following, $\bar{\mathbb{R}}_+$ represents $\{x \in \mathbb{R} : x \geq 0\}$.

Let $\alpha, \beta : \bar{\mathbb{R}}_+ \rightarrow \bar{\mathbb{R}}_+$ be strictly increasing absolutely continuous functions. The pair (α, β) is called a *pre-Hölder* pair if it obeys

$$(2.1) \quad \alpha^{-1}(x)\beta^{-1}(x) = x$$

for all $x \in \bar{\mathbb{R}}_+$. In the relation (2.1), the notations α^{-1}, β^{-1} are the inverse functions of α, β , respectively. Some examples of pre-Hölder pairs are:

$$(\alpha(x), \beta(x)) = (x^p, x^q)$$

for $p > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$, and

$$(2.2) \quad (\alpha, \beta) := (\lambda \circ A, \lambda \circ \tilde{A})$$

where we set $\lambda(x) = A^{-1}(x)\tilde{A}^{-1}(x)$ for any Orlicz N -function A together with its complementary N -function \tilde{A} .

In the following, Q stands for a cube whose sides are parallel to the axes and $|A|$ is the Lebesgue measure of the set A in \mathbb{R}^d , $d \geq 1$. We state the main theorem.

THEOREM 2.1. *Let $\hbar > 0$ be given. Suppose that (α, β) is a pre-Hölder pair such that for any positive constants $a, b > 0$, there exist constants θ_1, θ_2 and θ_f (depending on a and b) satisfying the following two conditions;*

$$\theta_1 + \theta_2 + \theta_f \leq \hbar$$

and

$$(2.3) \quad \alpha^{-1}(x)\beta^{-1}(y) \leq \theta_1 \frac{ab}{\alpha(a)} x + \theta_2 \frac{ab}{\beta(b)} y + ab\theta_f$$

for all $(x, y) \in \bar{\mathbb{R}}_+ \times \bar{\mathbb{R}}_+$. Then the functional

$$(2.4) \quad \|f\|_{BMO_\alpha} := \alpha^{-1} \left(\sup_Q \frac{1}{|Q|} \int_Q \alpha(|f(x) - f_Q|) dx \right)$$

satisfies a Minkowski type inequality : for any locally integrable functions f and g , we have

$$(2.5) \quad \|f + g\|_{BMO_\alpha} \leq \hbar \{ \|f\|_{BMO_\alpha} + \|g\|_{BMO_\alpha} \}$$

if the right hand side is finite. Also, for any constant $k \geq 0$ and for a locally integrable function f , we obtain

$$\frac{k}{\hbar} \|f\|_{BMO_\alpha} \leq \|kf\|_{BMO_\alpha} \leq k\hbar \|f\|_{BMO_\alpha}.$$

In particular, when $\hbar = 1$, we have the homogeneity:

$$\|kf\|_{BMO_\alpha} = k\|f\|_{BMO_\alpha}.$$

For example, any (convex) function satisfying

$$(2.6) \quad \alpha(x) := \begin{cases} x^p & \text{for } 0 \leq x \leq 1 \\ x^q & \text{for sufficiently large } x \end{cases}$$

($1 < p, q < \infty$) obeys the conditions in Theorem 2.1, and so are many variants of (2.6).

For a locally integrable function f on \mathbb{R}^d , we let

$$f_Q = \frac{1}{|Q|} \int_Q f(x) dx := \int_Q f(x) dx.$$

Let α be a pre-Hölder function. We denote a class of functions by

$$BMO_\alpha(\mathbb{R}^d) = \left\{ f \in L^1_{loc}(\mathbb{R}^d) : \|f\|_{BMO_\alpha} < \infty \right\},$$

where we set

$$\|f\|_{BMO_\alpha} = \alpha^{-1} \left(\sup_Q \int_Q \alpha(|f(x) - f_Q|) dx \right).$$

When α is the identity function, we write $BMO_\alpha(\mathbb{R}^d) := BMO(\mathbb{R}^d)$. In the sequel, the elements of $BMO_\alpha(\mathbb{R}^d)$ whose difference is a constant are identified. Note that even though we define $\|\cdot\|_{BMO_\alpha}$ on abstract measure spaces, we restrict our attention to the Euclidean space \mathbb{R}^d equipped with Lebesgue measure.

We now present the proof.

Proof of Theorem 2.1. We first present a Hölder type inequality: for any $f, g \in BMO_\alpha(\mathbb{R}^d)$, we have

$$(2.7) \quad \left| \int_Q f(x)g(x) dx \right| \leq \hbar \alpha^{-1} \left(\sup_Q \int_Q \alpha(|f(x)|) dx \right) \beta^{-1} \left(\sup_Q \int_Q \beta(|g(x)|) dx \right),$$

where Q is a cube whose sides are parallel to the axes.

For the proof of (2.7), we may assume that the right hand side of (2.7) is finite. We put

$$a := \alpha^{-1} \left(\sup_Q \int_Q \alpha(|f(x)|) dx \right), \quad b := \beta^{-1} \left(\sup_Q \int_Q \beta(|g(x)|) dx \right).$$

Then there exist constants θ_1, θ_2 and θ_f such that $\theta_1 + \theta_2 + \theta_f \leq \hbar$ and

$$\begin{aligned} |f(x)g(x)| &= \alpha^{-1}(\alpha(|f(x)|))\beta^{-1}(\beta(|g(x)|)) \\ (2.8) \quad &\leq \theta_1 \frac{ab}{\alpha(a)} \alpha(|f(x)|) + \theta_2 \frac{ab}{\beta(b)} \beta(|g(x)|) + ab\theta_f. \end{aligned}$$

Integrating over Q and dividing both sides by $|Q|$ yield

$$\begin{aligned} \int_Q |f(x)g(x)| dx &\leq \theta_1 \frac{ab}{\alpha(a)} \int_Q \alpha(|f(x)|) dx + \theta_2 \frac{ab}{\beta(b)} \int_Q \beta(|g(x)|) dx \\ &\quad + \theta_f ab \int_Q dx \\ &\leq \hbar \alpha^{-1} \left(\sup_Q \int_Q \alpha(|f(x)|) dx \right) \beta^{-1} \left(\sup_Q \int_Q \beta(|g(x)|) dx \right). \end{aligned}$$

This implies the Hölder type inequality (2.7).

We now verify the Minkowski type inequality (2.5). In fact, without loss of generality, we may assume that $f(x) + g(x) \neq 0$ almost every $x \in \mathbb{R}^d$ by restricting the domain \mathbb{R}^d if necessary. Applying Hölder type

inequality (2.7), we obtain

$$\begin{aligned}
& \int_Q \alpha(|f(x) + g(x) - f_Q - g_Q|) dx \\
& \leq \hbar \alpha^{-1} \left(\sup_Q \int_Q \alpha(|f(x) - f_Q|) dx \right) \\
& \quad \times \beta^{-1} \left(\sup_Q \int_Q \beta \left(\frac{\alpha(|f(x) - f_Q + g(x) - g_Q|)}{|f(x) - f_Q + g(x) - g_Q|} \right) dx \right) \\
& \quad + \hbar \beta^{-1} \left(\sup_Q \int_Q \beta(|g(x) - g_Q|) dx \right) \\
& \quad \times \beta^{-1} \left(\sup_Q \int_Q \beta \left(\frac{\alpha(|f(x) - f_Q + g(x) - g_Q|)}{|f(x) - f_Q + g(x) - g_Q|} \right) dx \right) \\
& = \hbar(\|f\|_{BMO_\alpha} + \|g\|_{BMO_\alpha}) \\
& \quad \times \beta^{-1} \left(\sup_Q \int_Q \beta \left(\frac{\alpha(|f(x) - f_Q + g(x) - g_Q|)}{|f(x) - f_Q + g(x) - g_Q|} \right) dx \right) \\
& = \hbar(\|f\|_{BMO_\alpha} + \|g\|_{BMO_\alpha})\beta^{-1} \left(\sup_Q \int_Q \alpha(|f(x) - f_Q + g(x) - g_Q|) dx \right).
\end{aligned}$$

The last equality follows from the fact that

$$(2.9) \quad \alpha(x) = \beta \left(\frac{\alpha(x)}{x} \right).$$

In fact, we solve for $\beta^{-1}(x)$ in the conjugate identity $\alpha^{-1}(x)\beta^{-1}(x) = x$ to get $\beta^{-1}(x) = \frac{x}{\alpha^{-1}(x)}$, which in turn yields

$$x = \beta \left(\frac{x}{\alpha^{-1}(x)} \right).$$

This illustrates the identity (2.9). Therefore we obtain

$$(2.10) \quad \frac{\alpha(\|f + g\|_{BMO_\alpha})}{\beta^{-1}(\alpha(\|f + g\|_{BMO_\alpha}))} \leq \|f\|_{BMO_\alpha} + \|g\|_{BMO_\alpha}.$$

From a variance of the conjugate identity: $\alpha^{-1}(x) = \frac{x}{\beta^{-1}(x)}$, we have

$$(2.11) \quad x = \frac{\alpha(x)}{\beta^{-1}(\alpha(x))}.$$

Hence from (2.10), we conclude the Minkowski type inequality (2.5):

$$\|f + g\|_{BMO_\alpha} \leq \|f\|_{BMO_\alpha} + \|g\|_{BMO_\alpha}.$$

We now verify that for any constant $k \geq 0$ and for $f \in BMO_\alpha(\mathbb{R}^d)$, we have

$$(2.12) \quad \frac{k}{\hbar} \|f\|_{BMO_\alpha} \leq \|kf\|_{BMO_\alpha} \leq k\hbar \|f\|_{BMO_\alpha}.$$

For each $f \in BMO_\alpha(\mathbb{R}^d)$, the associated operator (inhomogeneous) norm of f is defined by

$$(2.13) \quad \|f\|_* := \sup \left\{ \frac{\left| \sup_Q \int_Q (f(x) - f_Q)g(x)dx \right|}{\beta^{-1} \left(\sup_Q \int_Q \beta(|g(x)|)dx \right)} : g(x) \neq 0 \text{ almost everywhere} \right\}.$$

Then we note that for any constant $k \geq 0$,

$$(2.14) \quad \|kf\|_* = k\|f\|_*$$

and by virtue of the Hölder type inequality (2.7), we have

$$\frac{\left| \sup_Q \int_Q (f(x) - f_Q)g(x)dx \right|}{\beta^{-1} \left(\sup_Q \int_Q \beta(|g(x)|)dx \right)} \leq \hbar \|f\|_{BMO_\alpha}$$

for each measurable function g with $g(x) \neq 0$ almost everywhere. On the other hand, taking

$$g(x) := \frac{\alpha(|f(x) - f_Q|) \operatorname{sgn}(f(x) - f_Q)}{|f(x) - f_Q|},$$

we see that the identity (2.9) and its variants lead to

$$\begin{aligned} \beta^{-1} \left(\sup_Q \int_Q \beta(|g(x)|)dx \right) &= \beta^{-1} \left(\sup_Q \int_Q \alpha(|f(x) - f_Q|)dx \right) \\ &= (\beta^{-1} \circ \alpha)(\|f\|_{BMO_\alpha}) \\ &= \frac{\alpha(\|f\|_{BMO_\alpha})}{\|f\|_{BMO_\alpha}}. \end{aligned}$$

Therefore we obtain

$$\begin{aligned} \|f\|_* &\geq \frac{\left| \int_Q f_Q (f(x) - f_Q) g(x) dx \right|}{\beta^{-1} \left(\int_Q f_Q \beta(|g(x)|) dx \right)} \\ &= \frac{\left| \int_Q f_Q \alpha(|f - f_Q|) dx \right|}{\beta^{-1} \left(\int_Q f_Q \beta(|g(x)|) dx \right)} \\ &= \|f\|_{BMO_\alpha}. \end{aligned}$$

In all, we get

$$\|f\|_{BMO_\alpha} \leq \|f\|_* \leq \hbar \|f\|_{BMO_\alpha}.$$

For any constant $k \geq 0$ and for $f \in BMO_\alpha(\mathbb{R}^d)$, the identity (2.14) yields

$$\|kf\|_{BMO_\alpha} \leq k\|f\|_* \leq k\hbar\|f\|_{BMO_\alpha}$$

and

$$\hbar\|kf\|_{BMO_\alpha} \geq k\|f\|_* \geq k\|f\|_{BMO_\alpha},$$

which imply the inequalities (2.12). When $\hbar = 1$, we have the homogeneity:

$$\|kf\|_{BMO_\alpha} = k\|f\|_{BMO_\alpha}.$$

This completes the proof. \square

The functional $\|\cdot\|_{BMO_\alpha}$ on $BMO_\alpha(\mathbb{R}^d)$ may not produce a norm, since it does not always satisfy the homogeneity required for norms. Instead, by virtue of Minkowski's inequality (2.5), we may define a metric on $BMO_\alpha(\mathbb{R}^d)$ by

$$d(f, g) := \|f - g\|_{BMO_\alpha} \quad \text{for } f, g \in BMO_\alpha(\mathbb{R}^d).$$

It formulates a *complete* metric space on $BMO_\alpha(\mathbb{R}^d)$. The arguments comply the following standard procedure.

Suppose that $\{f_n\}$ is a Cauchy sequence in $BMO_\alpha(\mathbb{R}^d)$. Then there exists a subsequence $\{f_{n_k}\}$ of $\{f_n\}$ such that

$$d(f_{n_{k+1}}, f_{n_k}) \leq \frac{1}{(2\hbar)^k}, \quad k = 1, 2, \dots.$$

Setting F with

$$F(x) = |f_{n_1}(x)| + \sum_{k=1}^{\infty} |f_{n_{k+1}}(x) - f_{n_k}(x)|,$$

we can notice that $|F(x)| < \infty$ almost everywhere $x \in \mathbb{R}^d$. In fact, from the fact that

$$\begin{aligned} \|F\|_{BMO_\alpha} &\leq \hbar \|f_{n_1}\|_{BMO_\alpha} + \sum_{k=1}^{\infty} \hbar^{k+1} \|f_{n_{k+1}} - f_{n_k}\|_{BMO_\alpha} \\ &= \hbar \|f_{n_1}\|_{BMO_\alpha} + \hbar < \infty, \end{aligned}$$

there exists a null set $N \subset \mathbb{R}^d$ such that $F(x) < \infty$ for all $x \in \mathbb{R}^d \setminus N$. Therefore for any $x \in \mathbb{R}^d \setminus N$, the absolute convergence of the series

$$f_{n_1}(x) + \sum_{k=1}^{\infty} [f_{n_{k+1}}(x) - f_{n_k}(x)]$$

makes it possible to define $f(x) := \lim_{k \rightarrow \infty} f_{n_k}(x)$ on $\mathbb{R}^d \setminus N$. The fact

$$\begin{aligned} \|f - f_{n_k}\|_{BMO_\alpha} &= \left\| \sum_{j=k+1}^{\infty} f_{n_{j+1}} - f_{n_j} \right\|_{BMO_\alpha} \\ &\leq \sum_{j=k+1}^{\infty} \hbar^{j+1} \|f_{n_{j+1}} - f_{n_j}\|_{BMO_\alpha} = \frac{\hbar}{2^k} \end{aligned}$$

and the inequality

$$\|f\|_{BMO_\alpha} \leq \hbar \|f - f_{n_k}\|_{BMO_\alpha} + \hbar \|f_{n_k}\|_{BMO_\alpha}$$

yield $f \in BMO_\alpha$ and the convergence of $\{f_{n_k}\}$ to f in $BMO_\alpha(\mathbb{R}^d)$, which, in turn, implies the convergence of the original *Cauchy* sequence $\{f_n\}$ in $BMO_\alpha(\mathbb{R}^d)$.

References

- [1] F. John and L. Nirenberg, *On functions of bounded mean oscillation*, Comm. Pure and Appl. Math., **14** (1961), 415-426.
- [2] H. C. Pak, *On the John-Nirenberg inequality*, J. Inequal. Appl., **2020** (2020), no. 130.
- [3] H. C. Pak and Y. J. Park, *Trace operator and a nonlinear boundary value problem in a new space*, Boundary Value Problems, **2014** (2014), 2014:153.
- [4] H. C. Pak and Y. J. Park, *Spectrum and Singular Integrals on a New Weighted Function Space*, Acta Mathematica Sinica, English Series, **34** (2018), no. 11, 1692-1702.
- [5] Y. J. Park, *On the continuity of the Hardy-Littlewood maximal function*, J. Chung-cheong Math. Soc., **31** (2018), 43-46.
- [6] E. M. Stein, *Harmonic analysis; Real-variable methods, orthogonality, and oscillatory integrals*, Princeton University Press, 1995.

Young Ja Park
Department of Mathematics
Hoseo University
Hoseo-ro 79, Asan, Chungnam 31499
Republic of Korea
E-mail: ypark@hoseo.edu