

Nonlinear Functional Analysis and Applications
Vol. 27, No. 2 (2022), pp. 381-403

ISSN: 1229-1595(print), 2466-0973(online)

<https://doi.org/10.22771/nfaa.2022.27.02.12>

<http://nfaa.kyungnam.ac.kr/journal-nfaa>

Copyright © 2022 Kyungnam University Press



ON STRONG CONVERGENCE THEOREMS FOR A VISCOSITY-TYPE TSENG'S EXTRAGRADIENT METHODS SOLVING QUASIMONOTONE VARIATIONAL INEQUALITIES

Nopparat Wairojjana¹, Nattawut Pholasa²
and Nuttapol Pakkaranang³

¹Applied Mathematics Program, Faculty of Science and Technology
Valaya Alongkorn Rajabhat University under the Royal Patronage
Pathumthani 13180, Thailand
e-mail: nopparat@vru.ac.th

²School of Science, University of Phayao, Phayao 56000, Thailand
e-mail: nattawut_math@hotmail.com

³Mathematics and Computing Science Program, Faculty of Science and Technology
Phetchabun Rajabhat University, Phetchabun 67000, Thailand
e-mail: nuttapol.pak@pcru.ac.th

Abstract. The main goal of this research is to solve variational inequalities involving quasimonotone operators in infinite-dimensional real Hilbert spaces numerically. The main advantage of these iterative schemes is the ease with which step size rules can be designed based on an operator explanation rather than the Lipschitz constant or another line search method. The proposed iterative schemes use a monotone and non-monotone step size strategy based on mapping (operator) knowledge as a replacement for the Lipschitz constant or another line search method. The strong convergences have been demonstrated to correspond well to the proposed methods and to settle certain control specification conditions. Finally, we propose some numerical experiments to assess the effectiveness and influence of iterative methods.

⁰Received December 1, 2021. Revised January 14, 2022. Accepted March 14, 2022.

⁰2020 Mathematics Subject Classification: 47J25, 47H09, 47H06, 47J05.

⁰Keywords: Variational inequality problem, Tseng's extragradient method, strong convergence result, quasimonotone operator, Lipschitz continuity.

⁰Corresponding author: Nuttapol Pakkaranang(nuttapol.pak@pcru.ac.th).

1. INTRODUCTION

The main objective of this study is to examine the iterative methods employed to determine the solution to variational inequality problem (shortly, VIP) [34] involving quasimonotone operators in any real Hilbert space. Consider that Σ is a real Hilbert space and Δ is a nonempty, closed and convex subset of Σ . Look at the operator $\Gamma : \Sigma \rightarrow \Sigma$. The variational inequality problem for Γ on Δ is described in the subsequent fashion:

$$\text{Find } \omega^* \in \Delta \text{ such that } \langle \Gamma(\omega^*), y - \omega^* \rangle \geq 0, \forall y \in \Delta. \quad (\text{VIP})$$

The mathematical model of the variational inequality problem is a key problem in nonlinear analysis. It is a remarkable mathematical design that consolidates a lot of essential notions in applied mathematics, such as a nonlinear system of equation, optimization conditions for problems with the optimization process, the complementarity problems, network equilibrium problems and finance (see for more details [8, 11, 12, 13, 14, 15, 23]). As a consequence, this notion has various applications in the fields of mathematical programming, engineering, transport analysis, network economics, game theory and computer science. The regularized method and the projection method are two prominent and general procedures for finding a solution to variational inequalities. It is also noted that the first approach is most commonly used to deal with the variational inequalities accompanied by the class of monotone operators. The regularized sub-problem in this method is strongly monotone, and its unique solution is found more conveniently than the initial problem.

In this study, we discuss the projection methods that are well known for their simpler numerical computing. Many authors have dedicated themselves to studying not only the theory of existence and stability of solutions but also iterative methods for solving variational inequality problems. In addition, projection methods are useful to approximate the numerical solution of variational inequalities. Many researchers have established different variants of projection methods to solve such problems (see for more details [3, 5, 6, 7, 10, 16, 20, 22, 24, 33, 35, 38, 39, 41]) and others in [4, 9, 21, 25, 26, 27, 28, 29, 30, 31, 32, 36, 40, 42]. Almost all methods for solving the problem (VIP) are based on the computation of a projection on the feasible set Δ . Korpelevich [16] and Antipin [1] introduced the following extragradient method. Their method takes the following form:

$$\begin{cases} u_1 \in \Delta, \\ y_n = P_\Delta[u_n - \alpha\Gamma(u_n)], \\ u_{n+1} = P_\Delta[u_n - \alpha\Gamma(y_n)], \end{cases} \quad (1.1)$$

where $0 < \varkappa < \frac{1}{L}$. In view of the above method, we have used two projections on the underlying set Δ for each iteration. This, of course, can affect the computational effectiveness of the method if the feasible set Δ has a complicated structure.

Here, we present some methods which can remove this drawback. The first is the following subgradient extragradient method introduced by Censor et al. [6]. This method takes the following form:

$$\begin{cases} u_1 \in \Delta, \\ y_n = P_{\Delta}[u_n - \varkappa\Gamma(u_n)], \\ u_{n+1} = P_{\Sigma_n}[u_n - \varkappa\Gamma(y_n)], \end{cases} \quad (1.2)$$

where $0 < \varkappa < \frac{1}{L}$ and

$$\Sigma_n = \{z \in \Sigma : \langle u_n - \varkappa\Gamma(u_n) - y_n, z - y_n \rangle \leq 0\}.$$

In this article, our main focus on the Tseng's extragradient method [35] that uses only one projection for each iteration. This method takes the following form:

$$\begin{cases} u_1 \in \Delta, \\ y_n = P_{\Delta}[u_n - \varkappa\Gamma(u_n)], \\ u_{n+1} = y_n + \varkappa[\Gamma(u_n) - \Gamma(y_n)], \end{cases} \quad (1.3)$$

where $0 < \varkappa < \frac{1}{L}$. It is important to note that the above mentioned methods have two major flaws: a fixed constant step size rule that is dependent on the Lipschitz constant of mapping and generates a weakly convergent iterative sequence. The Lipschitz constant is generally unknown or difficult to compute. From a computational point of view, it can be difficult to consider a fixed step size constraint that affects the method's efficiency and rate of convergence. In addition, the study of a strongly convergent iterative sequence is important in the context of an infinite-dimensional Hilbert space.

The main objective of this paper is to introduce a new strongly convergent method by using viscosity and Tseng's extragradient-type method, including a monotonic and non-monotonic variable step size rule to solve variational inequalities involving the quasimonotone operator. Furthermore, to show that the iterative sequences generated by all four subgradient extragradient algorithms strongly converge to a solution. Both the monotone and non-monotone variable step size rules are used in subgradient and extragradient algorithms.

The paper is arranged in the following way: In Sect. 2, preliminary results were presented. Sect. 3 gives all new algorithms and their convergence analysis. Finally, Sect. 4 gives some numerical results to explain the practical efficiency of the proposed methods.

2. PRELIMINARIES

This section contains a number of important identities and relevant lemmas. For any $u, y \in \Sigma$, we have

$$\|u + y\|^2 = \|u\|^2 + 2\langle u, y \rangle + \|y\|^2.$$

Lemma 2.1. ([2]) *For any $y_1, y_2 \in \Sigma$ and $\ell \in \mathbb{R}$, The following inequalities are hold.*

- (i) $\|\ell y_1 + (1 - \ell)y_2\|^2 = \ell\|y_1\|^2 + (1 - \ell)\|y_2\|^2 - \ell(1 - \ell)\|y_1 - y_2\|^2.$
- (ii) $\|y_1 + y_2\|^2 \leq \|y_1\|^2 + 2\langle y_2, y_1 + y_2 \rangle.$

A metric projection $P_\Delta(y_1)$ of $y_1 \in \Sigma$ is defined by

$$P_\Delta(y_1) = \arg \min\{\|y_1 - y_2\| : y_2 \in \Delta\}.$$

First, we list some of the important features of projection mapping.

Lemma 2.2. ([2]) *Let $P_\Delta : \Sigma \rightarrow \Delta$ be a metric projection. Then, the following conditions are satisfied.*

- (i) $y_3 = P_\Delta(y_1)$ if and only if $\langle y_1 - y_3, y_2 - y_3 \rangle \leq 0, \forall y_2 \in \Delta,$
- (ii) $\|y_1 - P_\Delta(y_2)\|^2 + \|P_\Delta(y_2) - y_2\|^2 \leq \|y_1 - y_2\|^2, y_1 \in \Delta, y_2 \in \Sigma,$
- (iii) $\|y_1 - P_\Delta(y_1)\| \leq \|y_1 - y_2\|, y_2 \in \Delta, y_1 \in \Sigma.$

Lemma 2.3. ([37]) *Let $\{e_n\} \subset [0, +\infty)$ be a sequence satisfies the following condition*

$$e_{n+1} \leq (1 - f_n)e_n + f_n g_n, \forall n \in \mathbb{N}.$$

In addition, two sequences $\{f_n\} \subset (0, 1)$ and $\{g_n\} \subset \mathbb{R}$ satisfies the following conditions:

$$\lim_{n \rightarrow +\infty} f_n = 0, \sum_{n=1}^{+\infty} f_n = +\infty \text{ and } \limsup_{n \rightarrow +\infty} g_n \leq 0.$$

Then, $\lim_{n \rightarrow +\infty} e_n = 0.$

Lemma 2.4. ([19]) *Let $\{e_n\} \subset \mathbb{R}$ be a sequence and there exists a subsequence $\{n_i\}$ of $\{n\}$ such that*

$$e_{n_i} < e_{n_{i+1}}, \forall i \in \mathbb{N}.$$

Then, there exists a nondecreasing sequence $m_k \subset \mathbb{N}$ such that $m_k \rightarrow +\infty$ as $k \rightarrow +\infty$ with

$$e_{m_k} \leq e_{m_{k+1}} \text{ and } e_k \leq e_{m_{k+1}}, \forall k \in \mathbb{N}.$$

Indeed, $m_k = \max\{j \leq k : e_j \leq e_{j+1}\}.$

3. MAIN RESULTS

In this section, we propose a class of iterative algorithms for solving quasi-monotone variational inequalities based on Tseng's extragradient method. In the following text, all methods are described in detail. The following conditions are assumed to be met in order to verify the strong convergence:

(Γ1) the solution set for problem (VIP) is denoted by $VI(\Delta, \Gamma)$ and it is nonempty;

(Γ2) an operator $\Gamma : \Sigma \rightarrow \Sigma$ is said to be quasimonotone if

$$\langle \Gamma(u), y - u \rangle > 0 \implies \langle \Gamma(y), y - u \rangle \geq 0, \quad \forall u, y \in \Delta; \quad (\text{QM})$$

(Γ3) an operator $\Gamma : \Sigma \rightarrow \Sigma$ is said to be Lipschitz continuous if there exists a constant $L > 0$ such that

$$\|\Gamma(u) - \Gamma(y)\| \leq L\|u - y\|, \quad \forall u, y \in \Delta; \quad (\text{LC})$$

(Γ4) an operator $\Gamma : \Sigma \rightarrow \Sigma$ is weakly sequentially continuous if $\{\Gamma(u_n)\}$ weakly converges to $\Gamma(u)$ for every sequence $\{u_n\}$ weakly converges to u .

In this part, we present an iterative scheme for solving quasimonotone variational inequality problems that is based on Tseng's extragradient method [35] and viscosity scheme [20]. It is important to note that the proposed method has a straightforward structure for achieving strong convergence. Suppose that $g : \Sigma \rightarrow \Sigma$ is a strict contraction function with constant $\xi \in [0, 1)$. The main algorithm has been presented as follows:

Algorithm 3.1. (Viscosity Extragradient Method With Fixed Step Size Rule)

Step 0. Let $u_1 \in \Delta$, $0 < \varkappa < \frac{1}{L}$ and $\{\vartheta_n\} \subset (0, 1)$ meet the following conditions:

$$\lim_{n \rightarrow +\infty} \vartheta_n = 0 \quad \text{and} \quad \sum_{n=1}^{+\infty} \vartheta_n = +\infty.$$

Step 1. Compute

$$y_n = P_\Delta(u_n - \varkappa\Gamma(u_n)).$$

If $u_n = y_n$, Stop. Otherwise, go to Step 2.

Step 2. Compute

$$z_n = y_n + \varkappa[\Gamma(u_n) - \Gamma(y_n)].$$

Step 3. Compute

$$u_{n+1} = \vartheta_n g(u_n) + (1 - \vartheta_n)z_n.$$

Set $n := n + 1$ and go back to Step 1.

Algorithm 3.2. (Monotonic Explicit Viscosity Extragradient Method)

Step 0. Let $u_1 \in \Delta$, $\varkappa_1 > 0$, $\chi \in (0, 1)$ and $\{\vartheta_n\} \subset (0, 1)$ meet the following conditions:

$$\lim_{n \rightarrow +\infty} \vartheta_n = 0 \quad \text{and} \quad \sum_{n=1}^{+\infty} \vartheta_n = +\infty.$$

Step 1. Compute

$$y_n = P_{\Delta}(u_n - \varkappa_n \Gamma(u_n)).$$

If $u_n = y_n$, STOP. Otherwise, go to Step 2.

Step 2. Compute

$$z_n = y_n + \varkappa_n [\Gamma(u_n) - \Gamma(y_n)].$$

Step 3. Compute $u_{n+1} = \vartheta_n g(u_n) + (1 - \vartheta_n)z_n$.

Step 4. Compute

$$\varkappa_{n+1} = \begin{cases} \min \left\{ \varkappa_n, \frac{\chi \|u_n - y_n\|}{\|\Gamma(u_n) - \Gamma(y_n)\|} \right\} & \text{if } \Gamma(u_n) - \Gamma(y_n) \neq 0, \\ \varkappa_n & \text{otherwise.} \end{cases} \quad (3.1)$$

Set $n := n + 1$ and go back to Step 1.

Lemma 3.3. *The sequence $\{\varkappa_n\}$ generated by (3.1) is decreasing monotonically and converges to $\varkappa > 0$.*

Proof. It is given that Γ is Lipschitz-continuous with constant $L > 0$. Let $\Gamma(u_n) \neq \Gamma(y_n)$ such that

$$\frac{\chi \|u_n - y_n\|}{\|\Gamma(u_n) - \Gamma(y_n)\|} \geq \frac{\chi \|u_n - y_n\|}{L \|u_n - y_n\|} \geq \frac{\chi}{L}. \quad (3.2)$$

The above expression implies that $\lim_{n \rightarrow +\infty} \varkappa_n = \varkappa$. \square

Algorithm 3.4. (Non-Monotonic Explicit Viscosity Extragradient Method)

Step 0. Let $u_1 \in \Delta$, $\varkappa_1 > 0$, $\chi \in (0, 1)$ and sequence $\{\varphi_n\}$ satisfying $\sum_{n=1}^{+\infty} \varphi_n < +\infty$. Moreover, $\{\vartheta_n\} \subset (0, 1)$ satisfying the following conditions:

$$\lim_{n \rightarrow +\infty} \vartheta_n = 0 \quad \text{and} \quad \sum_{n=1}^{+\infty} \vartheta_n = +\infty.$$

Step 1. Compute

$$y_n = P_{\Delta}(u_n - \varkappa_n \Gamma(u_n)).$$

If $u_n = y_n$, STOP. Otherwise, go to Step 2.

Step 2. Compute

$$z_n = y_n + \varkappa_n [\Gamma(u_n) - \Gamma(y_n)].$$

Step 3. Compute $u_{n+1} = \vartheta_n g(u_n) + (1 - \vartheta_n)z_n$.

Step 4. Compute

$$\varkappa_{n+1} = \begin{cases} \min \left\{ \varkappa_n + \varphi_n, \frac{\chi \|u_n - y_n\|}{\|\Gamma(u_n) - \Gamma(y_n)\|} \right\} & \text{if } \Gamma(u_n) - \Gamma(y_n) \neq 0, \\ \varkappa_n + \varphi_n & \text{otherwise.} \end{cases} \quad (3.3)$$

Set $n := n + 1$ and go back to Step 1.

Lemma 3.5. *A sequence $\{\varkappa_n\}$ generated by (3.3) is convergent to \varkappa and satisfying the following inequality*

$$\min \left\{ \frac{\chi}{L}, \varkappa_1 \right\} \leq \varkappa_n \leq \varkappa_1 + P \quad \text{where } P = \sum_{n=1}^{+\infty} \varphi_n.$$

Proof. It is given that Γ is Lipschitz-continuous with constant $L > 0$. Let $\Gamma(u_n) \neq \Gamma(y_n)$ such that

$$\frac{\chi \|u_n - y_n\|}{\|\Gamma(u_n) - \Gamma(y_n)\|} \geq \frac{\chi \|u_n - y_n\|}{L \|u_n - y_n\|} \geq \frac{\chi}{L}. \quad (3.4)$$

By using mathematical induction on the definition of \varkappa_{n+1} , we have

$$\min \left\{ \frac{\chi}{L}, \varkappa_1 \right\} \leq \varkappa_n \leq \varkappa_1 + P.$$

Let

$$[\varkappa_{n+1} - \varkappa_n]^+ = \max \{0, \varkappa_{n+1} - \varkappa_n\}$$

and

$$[\varkappa_{n+1} - \varkappa_n]^- = \max \{0, -(\varkappa_{n+1} - \varkappa_n)\}.$$

Then, from the definition of $\{\varkappa_n\}$, we have

$$\sum_{n=1}^{+\infty} [\varkappa_{n+1} - \varkappa_n]^+ = \sum_{n=1}^{+\infty} \max \{0, \varkappa_{n+1} - \varkappa_n\} \leq P < +\infty. \quad (3.5)$$

That means that the series $\sum_{n=1}^{+\infty} [\varkappa_{n+1} - \varkappa_n]^+$ is convergent.

Next, we need to prove the convergence of $\sum_{n=1}^{+\infty} [\varkappa_{n+1} - \varkappa_n]^-$.

Let $\sum_{n=1}^{+\infty} [\varkappa_{n+1} - \varkappa_n]^- = +\infty$. Due to the reason that $\varkappa_{n+1} - \varkappa_n = (\varkappa_{n+1} - \varkappa_n)^+ - (\varkappa_{n+1} - \varkappa_n)^-$, we have

$$\varkappa_{k+1} - \varkappa_1 = \sum_{n=0}^k (\varkappa_{n+1} - \varkappa_n) = \sum_{n=0}^k [\varkappa_{n+1} - \varkappa_n]^+ - \sum_{n=0}^k [\varkappa_{n+1} - \varkappa_n]^-. \quad (3.6)$$

By allowing $k \rightarrow +\infty$ in (3.6), we have $\varkappa_k \rightarrow -\infty$ as $k \rightarrow +\infty$. This is a contradiction. Due to the convergence of the series $\sum_{n=0}^k [\varkappa_{n+1} - \varkappa_n]^+$ and $\sum_{n=0}^k [\varkappa_{n+1} - \varkappa_n]^-$ taking $k \rightarrow +\infty$ in (3.6), we obtain $\lim_{n \rightarrow +\infty} \varkappa_n = \varkappa$. This completes the proof. \square

Lemma 3.6. *Suppose that $\Gamma : \Sigma \rightarrow \Sigma$ satisfies the conditions $(\Gamma 1)$ - $(\Gamma 4)$ and sequence $\{u_n\}$ generated by Algorithm 3.1. Then, we have*

$$\|u_{n+1} - \omega^*\|^2 \leq \|u_n - \omega^*\|^2 - (1 - \varkappa^2 L^2) \|u_n - y_n\|^2.$$

Proof. Since $\omega^* \in VI(\Delta, \Gamma)$, we have

$$\begin{aligned} & \|u_{n+1} - \omega^*\|^2 \\ &= \|y_n + \varkappa[\Gamma(u_n) - \Gamma(y_n)] - \omega^*\|^2 \\ &= \|y_n - \omega^*\|^2 + \varkappa^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 + 2\varkappa \langle y_n - \omega^*, \Gamma(u_n) - \Gamma(y_n) \rangle \\ &= \|y_n + u_n - u_n - \omega^*\|^2 + \varkappa^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 + 2\varkappa \langle y_n - \omega^*, \Gamma(u_n) - \Gamma(y_n) \rangle \\ &= \|y_n - u_n\|^2 + \|u_n - \omega^*\|^2 + 2\langle y_n - u_n, u_n - \omega^* \rangle \\ &\quad + \varkappa^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 + 2\varkappa \langle y_n - \omega^*, \Gamma(u_n) - \Gamma(y_n) \rangle \\ &= \|u_n - \omega^*\|^2 + \|y_n - u_n\|^2 + 2\langle y_n - u_n, y_n - \omega^* \rangle + 2\langle y_n - u_n, u_n - y_n \rangle \\ &\quad + \varkappa^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 + 2\varkappa \langle y_n - \omega^*, \Gamma(u_n) - \Gamma(y_n) \rangle. \end{aligned} \tag{3.7}$$

It is given that $y_n = P_\Delta[u_n - \varkappa\Gamma(u_n)]$ and it gives that

$$\langle u_n - \varkappa\Gamma(u_n) - y_n, y_n - y_n \rangle \leq 0, \quad \forall y \in \Delta. \tag{3.8}$$

Thus, we have

$$\langle u_n - y_n, \omega^* - y_n \rangle \leq \varkappa \langle \Gamma(u_n), \omega^* - y_n \rangle. \tag{3.9}$$

Combining expressions (3.7) and (3.9), we have

$$\begin{aligned} & \|u_{n+1} - \omega^*\|^2 \\ &\leq \|u_n - \omega^*\|^2 + \|y_n - u_n\|^2 + 2\varkappa \langle \Gamma(u_n), \omega^* - y_n \rangle - 2\langle u_n - y_n, u_n - y_n \rangle \\ &\quad + \varkappa^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 - 2\varkappa \langle \Gamma(u_n) - \Gamma(y_n), \omega^* - y_n \rangle \\ &= \|u_n - \omega^*\|^2 - \|u_n - y_n\|^2 + \varkappa^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 - 2\varkappa \langle \Gamma(y_n), y_n - \omega^* \rangle. \end{aligned} \tag{3.10}$$

It is given that ω^* is the solution of the problem (VIP) implies that

$$\langle \Gamma(\omega^*), y - \omega^* \rangle \geq 0, \quad \forall y \in \Delta.$$

It implies that

$$\langle \Gamma(y), y - \omega^* \rangle \geq 0, \quad \forall y \in \Delta.$$

By substituting $y = y_n \in \Delta$, we have

$$\langle \Gamma(y_n), y_n - \omega^* \rangle \geq 0. \quad (3.11)$$

From expressions (3.10) and (3.11), we obtain

$$\begin{aligned} \|u_{n+1} - \omega^*\|^2 &\leq \|u_n - \omega^*\|^2 - \|u_n - y_n\|^2 + \varkappa^2 L^2 \|u_n - y_n\|^2 \\ &= \|u_n - \omega^*\|^2 - (1 - \varkappa^2 L^2) \|u_n - y_n\|^2. \end{aligned} \quad (3.12)$$

□

Lemma 3.7. *Assume that $\Gamma : \Sigma \rightarrow \Sigma$ satisfies the conditions $(\Gamma 1)$ - $(\Gamma 4)$. Let $\{u_n\}$ be a sequence is generated by Algorithm 3.2 and 3.4. Then, for each $\omega^* \in VI(\Delta, \Gamma)$, we have*

$$\|u_{n+1} - \omega^*\|^2 \leq \|u_n - \omega^*\|^2 - \left(1 - \chi^2 \frac{\varkappa_n^2}{\varkappa_{n+1}^2}\right) \|u_n - y_n\|^2.$$

Proof. Let $\omega^* \in VI(\Delta, \Gamma)$ and by definition of u_{n+1} , we have

$$\begin{aligned} &\|u_{n+1} - \omega^*\|^2 \\ &= \|y_n + \varkappa_n [\Gamma(u_n) - \Gamma(y_n)] - \omega^*\|^2 \\ &= \|y_n - \omega^*\|^2 + \varkappa_n^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 + 2\varkappa_n \langle y_n - \omega^*, \Gamma(u_n) - \Gamma(y_n) \rangle \\ &= \|y_n + u_n - u_n - \omega^*\|^2 + \varkappa_n^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 \\ &\quad + 2\varkappa_n \langle y_n - \omega^*, \Gamma(u_n) - \Gamma(y_n) \rangle \\ &= \|y_n - u_n\|^2 + \|u_n - \omega^*\|^2 + 2\langle y_n - u_n, u_n - \omega^* \rangle \\ &\quad + \varkappa_n^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 + 2\varkappa_n \langle y_n - \omega^*, \Gamma(u_n) - \Gamma(y_n) \rangle \\ &= \|u_n - \omega^*\|^2 + \|y_n - u_n\|^2 + 2\langle y_n - u_n, y_n - \omega^* \rangle + 2\langle y_n - u_n, u_n - y_n \rangle \\ &\quad + \varkappa_n^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 + 2\varkappa_n \langle y_n - \omega^*, \Gamma(u_n) - \Gamma(y_n) \rangle. \end{aligned} \quad (3.13)$$

It is given that $y_n = P_\Delta[u_n - \varkappa_n \Gamma(u_n)]$ and it further implies that

$$\langle u_n - \varkappa_n \Gamma(u_n) - y_n, y - y_n \rangle \leq 0, \quad \forall y \in \Delta \quad (3.14)$$

or equivalently for some $\omega^* \in VI(\Delta, \Gamma)$, we can write

$$\langle u_n - y_n, \omega^* - y_n \rangle \leq \varkappa_n \langle \Gamma(u_n), \omega^* - y_n \rangle. \quad (3.15)$$

Combining expressions (3.13) and (3.15), we have

$$\begin{aligned}
& \|u_{n+1} - \omega^*\|^2 \\
& \leq \|u_n - \omega^*\|^2 + \|y_n - u_n\|^2 + 2\alpha_n \langle \Gamma(u_n), \omega^* - y_n \rangle - 2 \langle u_n - y_n, u_n - y_n \rangle \\
& \quad + \alpha_n^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 - 2\alpha_n \langle \Gamma(u_n) - \Gamma(y_n), \omega^* - y_n \rangle \\
& = \|u_n - \omega^*\|^2 - \|u_n - y_n\|^2 + \alpha_n^2 \|\Gamma(u_n) - \Gamma(y_n)\|^2 - 2\alpha_n \langle \Gamma(y_n), y_n - \omega^* \rangle.
\end{aligned} \tag{3.16}$$

It is given that ω^* is the solution of the problem (VIP), implies that

$$\langle \Gamma(\omega^*), y - \omega^* \rangle > 0, \quad \forall y \in \Delta.$$

Due to the property of Γ on Δ , we obtain

$$\langle \Gamma(y), y - \omega^* \rangle \geq 0, \quad \forall y \in \Delta.$$

Substituting $y = y_n \in \Delta$, we have

$$\langle \Gamma(y_n), y_n - \omega^* \rangle \geq 0. \tag{3.17}$$

Combining expressions (3.16) and (3.17), we obtain

$$\begin{aligned}
\|u_{n+1} - \omega^*\|^2 & \leq \|u_n - \omega^*\|^2 - \|u_n - y_n\|^2 + \chi^2 \frac{\alpha_n^2}{\alpha_{n+1}^2} \|u_n - y_n\|^2 \\
& = \|u_n - \omega^*\|^2 - \left(1 - \chi^2 \frac{\alpha_n^2}{\alpha_{n+1}^2}\right) \|u_n - y_n\|^2.
\end{aligned} \tag{3.18}$$

□

Lemma 3.8. *Let $\Gamma : \Sigma \rightarrow \Sigma$ be an operator satisfies the conditions (Γ1)–(Γ4). If there exists a weakly convergent subsequence $\{u_{n_k}\}$ to \hat{u} and $\lim_{k \rightarrow +\infty} \|u_{n_k} - y_{n_k}\| = 0$. Then, $\hat{u} \in VI(\Delta, \Gamma)$.*

Proof. Since $\{u_{n_k}\}$ weakly convergent to \hat{u} and due to $\lim_{k \rightarrow +\infty} \|u_{n_k} - y_{n_k}\| = 0$, the sequence $\{y_{n_k}\}$ also weakly convergent to \hat{u} . Next, we need to prove that $\hat{u} \in VI(\Delta, \Gamma)$. By value of y_n , we have

$$y_{n_k} = P_\Delta[u_{n_k} - \alpha_{n_k} \Gamma(u_{n_k})],$$

that is equivalent to

$$\langle u_{n_k} - \alpha_{n_k} \Gamma(u_{n_k}) - y_{n_k}, y - y_{n_k} \rangle \leq 0, \quad \forall y \in \Delta. \tag{3.19}$$

The above inequality implies that

$$\langle u_{n_k} - y_{n_k}, y - y_{n_k} \rangle \leq \alpha_{n_k} \langle \Gamma(u_{n_k}), y - y_{n_k} \rangle, \quad \forall y \in \Delta. \tag{3.20}$$

Thus, we obtain

$$\frac{1}{\varkappa_{n_k}} \langle u_{n_k} - y_{n_k}, y - y_{n_k} \rangle + \langle \Gamma(u_{n_k}), y_{n_k} - u_{n_k} \rangle \leq \langle \Gamma(u_{n_k}), y - u_{n_k} \rangle, \quad \forall y \in \Delta. \quad (3.21)$$

By use of $\lim_{k \rightarrow +\infty} \|u_{n_k} - y_{n_k}\| = 0$ and $k \rightarrow +\infty$ in (3.21), we have

$$\liminf_{k \rightarrow +\infty} \langle \Gamma(u_{n_k}), y - u_{n_k} \rangle \geq 0, \quad \forall y \in \Delta. \quad (3.22)$$

Furthermore, it implies that

$$\begin{aligned} \langle \Gamma(y_{n_k}), y - y_{n_k} \rangle &= \langle \Gamma(y_{n_k}) - \Gamma(u_{n_k}), y - u_{n_k} \rangle \\ &\quad + \langle \Gamma(u_{n_k}), y - u_{n_k} \rangle + \langle \Gamma(y_{n_k}), u_{n_k} - y_{n_k} \rangle. \end{aligned} \quad (3.23)$$

Since $\lim_{k \rightarrow +\infty} \|u_{n_k} - y_{n_k}\| = 0$, we have

$$\lim_{k \rightarrow +\infty} \|\Gamma(u_{n_k}) - \Gamma(y_{n_k})\| = 0, \quad (3.24)$$

which together with (3.23) and (3.24), we obtain

$$\liminf_{k \rightarrow +\infty} \langle \Gamma(y_{n_k}), y - y_{n_k} \rangle \geq 0, \quad \forall y \in \Delta. \quad (3.25)$$

Moreover, let us take a positive sequence $\{\epsilon_k\}$ that is decreasing and convergent to zero. For each $\{\epsilon_k\}$ there exists a least positive integer denoted by m_k such that

$$\langle \Gamma(u_{n_i}), y - u_{n_i} \rangle + \epsilon_k > 0, \quad \forall i \geq m_k. \quad (3.26)$$

Since $\{\epsilon_k\}$ is decreasing sequence, it is easy to see that the sequence $\{m_k\}$ is increasing. If there exists a natural number $N_0 \in \mathbb{N}$ such that for all $\Gamma(u_{n_{m_k}}) \neq 0$, $n_{m_k} \geq N_0$. Consider that

$$\aleph_{n_{m_k}} = \frac{\Gamma(u_{n_{m_k}})}{\|\Gamma(u_{n_{m_k}})\|^2}, \quad \forall n_{m_k} \geq N_0. \quad (3.27)$$

Due to the above definition, we have

$$\langle \Gamma(u_{n_{m_k}}), \aleph_{n_{m_k}} \rangle = 1, \quad \forall n_{m_k} \geq N_0. \quad (3.28)$$

Moreover, from expressions (3.26) and (3.28) for all $n_{m_k} \geq N_0$, we have

$$\langle \Gamma(u_{n_{m_k}}), y + \epsilon_k \aleph_{n_{m_k}} - u_{n_{m_k}} \rangle > 0. \quad (3.29)$$

By the definition of quasimonotone, we have

$$\langle \Gamma(y + \epsilon_k \aleph_{n_{m_k}}), y + \epsilon_k \aleph_{n_{m_k}} - u_{n_{m_k}} \rangle > 0. \quad (3.30)$$

For all $n_{m_k} \geq N_0$, we have

$$\langle \Gamma(y), y - u_{n_{m_k}} \rangle \geq \langle \Gamma(y) - \Gamma(y + \epsilon_k \aleph_{n_{m_k}}), y + \epsilon_k \aleph_{n_{m_k}} - u_{n_{m_k}} \rangle - \epsilon_k \langle \Gamma(y), \aleph_{n_{m_k}} \rangle. \quad (3.31)$$

Due to $\{u_{n_k}\}$ converges weakly to $\hat{u} \in \Delta$ with Γ is weakly sequentially continuous on the set Δ , we obtain $\{\Gamma(u_{n_k})\}$ converges weakly to $\Gamma(\hat{u})$. Let $\Gamma(\hat{u}) \neq 0$. Then

$$\|\Gamma(\hat{u})\| \leq \liminf_{k \rightarrow +\infty} \|\Gamma(u_{n_k})\|. \quad (3.32)$$

Since $\{u_{n_{m_k}}\} \subset \{u_{n_k}\}$ and $\lim_{k \rightarrow +\infty} \epsilon_k = 0$, we have

$$0 \leq \lim_{k \rightarrow +\infty} \|\epsilon_k \aleph_{n_{m_k}}\| = \lim_{k \rightarrow +\infty} \frac{\epsilon_k}{\|\Gamma(u_{n_{m_k}})\|} \leq \frac{0}{\|\Gamma(\hat{u})\|} = 0. \quad (3.33)$$

By letting $k \rightarrow +\infty$ in (3.31), we obtain

$$\langle \Gamma(y), y - \hat{u} \rangle \geq 0, \quad \forall y \in \Delta. \quad (3.34)$$

Let $u \in \Delta$ be arbitrary element and for $0 < \varkappa \leq 1$, let

$$\hat{u}_\varkappa = \varkappa u + (1 - \varkappa)\hat{u}. \quad (3.35)$$

Then $\hat{u}_\varkappa \in \Delta$ and from (3.34) we have

$$\varkappa \langle \Gamma(\hat{u}_\varkappa), u - \hat{u} \rangle \geq 0. \quad (3.36)$$

Hence

$$\langle \Gamma(\hat{u}_\varkappa), u - \hat{u} \rangle \geq 0. \quad (3.37)$$

Let $\varkappa \rightarrow 0$. Then $\hat{u}_\varkappa \rightarrow \hat{u}$ along a line segment. By the continuity of an operator, $\Gamma(\hat{u}_\varkappa)$ converges to $\Gamma(\hat{u})$ as $\varkappa \rightarrow 0$. It follows from (3.37) that

$$\langle \Gamma(\hat{u}), u - \hat{u} \rangle \geq 0. \quad (3.38)$$

Therefore \hat{u} is a solution of problem (VIP). \square

Theorem 3.9. *Assume that an operator $\Gamma : \Delta \rightarrow \Sigma$ satisfies the conditions (G1)–(G4) and ω^* belongs to the solution set $VI(\Delta, \Gamma)$. Moreover, sequence $\{\vartheta_n\} \subset (0, 1)$ satisfying the following conditions:*

$$\lim_{n \rightarrow +\infty} \vartheta_n = 0 \quad \text{and} \quad \sum_{n=1}^{+\infty} \vartheta_n = +\infty.$$

Then, the sequences $\{u_n\}$, $\{y_n\}$ and $\{z_n\}$ generated by Algorithm 3.4 converge strongly to $\omega^ = P_{VI(\Delta, \Gamma)} \circ g(\omega^*)$.*

Proof. By using Lemma 3.7, we have

$$\|z_n - \omega^*\|^2 \leq \|u_n - \omega^*\|^2 - \left(1 - \chi^2 \frac{\varkappa_n^2}{\varkappa_{n+1}^2}\right) \|u_n - y_n\|^2. \quad (3.39)$$

Given that $\varkappa_n \rightarrow \varkappa$, so there exists a fixed number $\epsilon \in (0, 1 - \chi^2)$ such that

$$\lim_{n \rightarrow +\infty} \left(1 - \chi^2 \frac{\varkappa_n^2}{\varkappa_{n+1}^2}\right) = 1 - \chi^2 > \epsilon > 0.$$

Thus, there exists a fixed number $N_1 \in \mathbb{N}$ such that

$$\left(1 - \chi^2 \frac{\chi_n^2}{\chi_{n+1}^2}\right) > \epsilon > 0, \quad \forall n \geq N_1. \quad (3.40)$$

Therefore, we obtain

$$\|z_n - \omega^*\|^2 \leq \|u_n - \omega^*\|^2, \quad \forall n \geq N_1. \quad (3.41)$$

It is given that $\omega^* \in VI(\Delta, \Gamma)$. From sequence $\{u_{n+1}\}$ and the reason that g is a contraction with constant $\xi \in [0, 1)$ and $n \geq N_1$, we have

$$\begin{aligned} \|u_{n+1} - \omega^*\| &= \|\vartheta_n g(u_n) + (1 - \vartheta_n)z_n - \omega^*\| \\ &= \|\vartheta_n [g(u_n) - \omega^*] + (1 - \vartheta_n)[z_n - \omega^*]\| \\ &= \|\vartheta_n [g(u_n) + g(\omega^*) - g(\omega^*) - \omega^*] + (1 - \vartheta_n)[z_n - \omega^*]\| \\ &\leq \vartheta_n \|g(u_n) - g(\omega^*)\| + \vartheta_n \|g(\omega^*) - \omega^*\| + (1 - \vartheta_n) \|z_n - \omega^*\| \\ &\leq \vartheta_n \xi \|u_n - \omega^*\| + \vartheta_n \|g(\omega^*) - \omega^*\| + (1 - \vartheta_n) \|z_n - \omega^*\|. \end{aligned} \quad (3.42)$$

Combining expressions (3.41) with (3.42) and $\vartheta_n \in (0, 1)$, we deduce that

$$\begin{aligned} \|u_{n+1} - \omega^*\| &\leq \vartheta_n \xi \|u_n - \omega^*\| + \vartheta_n \|g(\omega^*) - \omega^*\| + (1 - \vartheta_n) \|u_n - \omega^*\| \\ &= [1 - \vartheta_n + \xi \vartheta_n] \|u_n - \omega^*\| + \vartheta_n (1 - \xi) \frac{\|g(\omega^*) - \omega^*\|}{(1 - \xi)} \\ &\leq \max \left\{ \|u_n - \omega^*\|, \frac{\|g(\omega^*) - \omega^*\|}{(1 - \xi)} \right\} \\ &\leq \max \left\{ \|u_{N_1} - \omega^*\|, \frac{\|g(\omega^*) - \omega^*\|}{(1 - \xi)} \right\}. \end{aligned} \quad (3.43)$$

Therefore, we deduce that $\{u_n\}$ is a bounded sequence. Due to the continuity and monotonicity of the operator Γ implies that the solution set $VI(\Delta, \Gamma)$ is a closed and convex set (for more details see [17, 18]). Since the mapping is a contraction and so does $P_{VI(\Delta, \Gamma)} \circ g$.

Now, we are in position to use the Banach contraction theorem for the existence of a fixed point of $\omega^* \in VI(\Delta, \Gamma)$ such that

$$\omega^* = P_{VI(\Delta, \Gamma)}(g(\omega^*)).$$

By using Lemma 2.2 (ii), we have

$$\langle g(\omega^*) - \omega^*, y - \omega^* \rangle \leq 0, \quad \forall y \in VI(\Delta, \Gamma). \quad (3.44)$$

It is given that $u_{n+1} = \vartheta_n g(u_n) + (1 - \vartheta_n)z_n$, and using Lemma 2.1 (i) and Lemma 3.7, we have

$$\begin{aligned}
& \|u_{n+1} - \omega^*\|^2 \\
&= \|\vartheta_n g(u_n) + (1 - \vartheta_n)z_n - \omega^*\|^2 \\
&= \|\vartheta_n [g(u_n) - \omega^*] + (1 - \vartheta_n)[z_n - \omega^*]\|^2 \\
&= \vartheta_n \|g(u_n) - \omega^*\|^2 + (1 - \vartheta_n) \|z_n - \omega^*\|^2 - \vartheta_n(1 - \vartheta_n) \|g(u_n) - z_n\|^2 \\
&\leq \vartheta_n \|g(u_n) - \omega^*\|^2 + (1 - \vartheta_n) \left[\|u_n - \omega^*\|^2 - \left(1 - \chi^2 \frac{\varkappa_n^2}{\varkappa_{n+1}^2}\right) \|u_n - y_n\|^2 \right] \\
&\quad - \vartheta_n(1 - \vartheta_n) \|g(u_n) - z_n\|^2 \\
&\leq \vartheta_n \|g(u_n) - \omega^*\|^2 + \|u_n - \omega^*\|^2 - (1 - \vartheta_n) \left(1 - \chi^2 \frac{\varkappa_n^2}{\varkappa_{n+1}^2}\right) \|u_n - y_n\|^2.
\end{aligned} \tag{3.45}$$

The remainder of the proof shall be divided into the following two parts:

Case 1: Assume that there is a fixed number $N_2 \in \mathbb{N}$ ($N_2 \geq N_1$) such that

$$\|u_{n+1} - \omega^*\| \leq \|u_n - \omega^*\|, \quad \forall n \geq N_2. \tag{3.46}$$

Then, $\lim_{n \rightarrow +\infty} \|u_n - \omega^*\|$ exists and let $\lim_{n \rightarrow +\infty} \|u_n - \omega^*\| = l$. From expression (3.45), we have

$$\begin{aligned}
(1 - \vartheta_n) \left(1 - \chi^2 \frac{\varkappa_n^2}{\varkappa_{n+1}^2}\right) \|u_n - y_n\|^2 &\leq \vartheta_n \|g(u_n) - \omega^*\|^2 \\
&\quad + \|u_n - \omega^*\|^2 - \|u_{n+1} - \omega^*\|^2.
\end{aligned} \tag{3.47}$$

From the existence of $\lim_{n \rightarrow +\infty} \|u_n - \omega^*\| = l$, and $\vartheta_n \rightarrow 0$, we infer that

$$\lim_{n \rightarrow +\infty} \|u_n - y_n\| = 0. \tag{3.48}$$

It follows that

$$\|z_n - y_n\| = \|y_n + \varkappa_n [\Gamma(u_n) - \Gamma(y_n)] - y_n\| \leq \varkappa_0 L \|u_n - y_n\|.$$

This implies that

$$\lim_{n \rightarrow +\infty} \|z_n - y_n\| = 0. \tag{3.49}$$

It follows that

$$\lim_{n \rightarrow +\infty} \|u_n - z_n\| \leq \lim_{n \rightarrow +\infty} \|u_n - y_n\| + \lim_{n \rightarrow +\infty} \|y_n - z_n\| = 0. \tag{3.50}$$

We can also obtain

$$\begin{aligned}
\|u_{n+1} - u_n\| &= \|\vartheta_n g(u_n) + (1 - \vartheta_n)z_n - u_n\| \\
&= \|\vartheta_n[g(u_n) - u_n] + (1 - \vartheta_n)[z_n - u_n]\| \\
&\leq \vartheta_n \|g(u_n) - u_n\| + (1 - \vartheta_n)\|z_n - u_n\| \\
&\longrightarrow 0, \quad \text{as } n \rightarrow +\infty.
\end{aligned} \tag{3.51}$$

Hence we have

$$\lim_{n \rightarrow +\infty} \|u_{n+1} - u_n\| = 0. \tag{3.52}$$

Since the sequence $\{u_n\}$ is bounded, the sequences $\{y_n\}$ and $\{z_n\}$ are also bounded. Thus, we can take a subsequence $\{u_{n_k}\}$ of $\{u_n\}$ such that $\{u_{n_k}\}$ weakly converges to some $\hat{u} \in \Sigma$. Moreover, from $\|u_n - y_n\| \rightarrow 0$, we have $\hat{u} \in VI(\Delta, \Gamma)$. It follows that

$$\begin{aligned}
\limsup_{n \rightarrow +\infty} \langle g(\omega^*) - \omega^*, u_n - \omega^* \rangle &= \limsup_{k \rightarrow +\infty} \langle g(\omega^*) - \omega^*, u_{n_k} - \omega^* \rangle \\
&= \langle g(\omega^*) - \omega^*, \hat{u} - \omega^* \rangle \\
&\leq 0.
\end{aligned} \tag{3.53}$$

Since $\lim_{n \rightarrow +\infty} \|u_{n+1} - u_n\| = 0$. It follows that

$$\begin{aligned}
\limsup_{n \rightarrow +\infty} \langle g(\omega^*) - \omega^*, u_{n+1} - \omega^* \rangle &\leq \limsup_{n \rightarrow +\infty} \langle g(\omega^*) - \omega^*, u_{n+1} - u_n \rangle \\
&\quad + \limsup_{n \rightarrow +\infty} \langle g(\omega^*) - \omega^*, u_n - \omega^* \rangle \\
&\leq 0.
\end{aligned} \tag{3.54}$$

From Lemma 2.1 (ii) and Lemma 3.7 for all $n \geq N_2$, we obtain

$$\begin{aligned}
\|u_{n+1} - \omega^*\|^2 &= \|\vartheta_n g(u_n) + (1 - \vartheta_n)z_n - \omega^*\|^2 \\
&= \|\vartheta_n[g(u_n) - \omega^*] + (1 - \vartheta_n)[z_n - \omega^*]\|^2 \\
&\leq (1 - \vartheta_n)^2 \|z_n - \omega^*\|^2 \\
&\quad + 2\vartheta_n \langle g(u_n) - \omega^*, (1 - \vartheta_n)[z_n - \omega^*] + \vartheta_n[g(u_n) - \omega^*] \rangle \\
&= (1 - \vartheta_n)^2 \|z_n - \omega^*\|^2 \\
&\quad + 2\vartheta_n \langle g(u_n) - g(\omega^*) + g(\omega^*) - \omega^*, u_{n+1} - \omega^* \rangle
\end{aligned}$$

$$\begin{aligned}
&= (1 - \vartheta_n)^2 \|z_n - \omega^*\|^2 + 2\vartheta_n \langle g(u_n) - g(\omega^*), u_{n+1} - \omega^* \rangle \\
&\quad + 2\vartheta_n \langle g(\omega^*) - \omega^*, u_{n+1} - \omega^* \rangle \\
&\leq (1 - \vartheta_n)^2 \|z_n - \omega^*\|^2 + 2\vartheta_n \xi \|u_n - \omega^*\| \|u_{n+1} - \omega^*\| \\
&\quad + 2\vartheta_n \langle g(\omega^*) - \omega^*, u_{n+1} - \omega^* \rangle \\
&\leq (1 + \vartheta_n^2 - 2\vartheta_n) \|u_n - \omega^*\|^2 + 2\vartheta_n \xi \|u_n - \omega^*\|^2 \\
&\quad + 2\vartheta_n \langle g(\omega^*) - \omega^*, u_{n+1} - \omega^* \rangle \\
&= (1 - 2\vartheta_n) \|u_n - \omega^*\|^2 + \vartheta_n^2 \|u_n - \omega^*\|^2 + 2\vartheta_n \xi \|u_n - \omega^*\|^2 \\
&\quad + 2\vartheta_n \langle g(\omega^*) - \omega^*, u_{n+1} - \omega^* \rangle \\
&= [1 - 2\vartheta_n(1 - \xi)] \|u_n - \omega^*\|^2 \\
&\quad + 2\vartheta_n(1 - \xi) \left[\frac{\vartheta_n \|u_n - \omega^*\|^2}{2(1 - \xi)} + \frac{\langle g(\omega^*) - \omega^*, u_{n+1} - \omega^* \rangle}{1 - \xi} \right]. \tag{3.55}
\end{aligned}$$

It follows from expressions (3.54) and (3.55) that

$$\limsup_{n \rightarrow +\infty} \left[\frac{\vartheta_n \|u_n - \omega^*\|^2}{2(1 - \xi)} + \frac{\langle g(\omega^*) - \omega^*, u_{n+1} - \omega^* \rangle}{1 - \xi} \right] \leq 0. \tag{3.56}$$

Let choose $n \geq N_3 \in \mathbb{N}$ ($N_3 \geq N_2$) large enough such that

$$2\vartheta_n(1 - \xi) < 1.$$

By using (3.55) and (3.56), and applying Lemma 2.3, we conclude that $\|u_n - \omega^*\| \rightarrow 0$, as $n \rightarrow +\infty$.

Case 2: Assume there is a subsequence $\{n_i\}$ of $\{n\}$ such that

$$\|u_{n_i} - \omega^*\| \leq \|u_{n_{i+1}} - \omega^*\|, \quad \forall i \in \mathbb{N}.$$

Then, by Lemma 2.4, there is a sequence $\{m_k\} \subset \mathbb{N}$ as $m_k \rightarrow +\infty$, such that

$$\|u_{m_k} - \omega^*\| \leq \|u_{m_{k+1}} - \omega^*\| \quad \text{and} \quad \|u_k - \omega^*\| \leq \|u_{m_{k+1}} - \omega^*\|, \quad \text{for all } k \in \mathbb{N}. \tag{3.57}$$

As similar to Case 1, from (3.45), we have

$$\begin{aligned}
&(1 - \vartheta_{m_k}) \left(1 - \chi^2 \frac{\chi_{m_k}^2}{\chi_{m_{k+1}}^2} \right) \|u_{m_k} - y_{m_k}\|^2 \\
&\leq \vartheta_{m_k} \|g(u_{m_k}) - \omega^*\|^2 + \|u_{m_k} - \omega^*\|^2 - \|u_{m_{k+1}} - \omega^*\|^2. \tag{3.58}
\end{aligned}$$

Due to $\vartheta_{m_k} \rightarrow 0$, we deduce the following:

$$\lim_{k \rightarrow +\infty} \|u_{m_k} - y_{m_k}\| = 0. \tag{3.59}$$

Similar to above case we can prove that

$$\lim_{k \rightarrow +\infty} \|u_{m_k} - z_{m_k}\| = \lim_{k \rightarrow +\infty} \|y_{m_k} - z_{m_k}\| = 0. \quad (3.60)$$

Also, we obtain

$$\begin{aligned} \|u_{m_k+1} - u_{m_k}\| &= \|\vartheta_{m_k} g(u_{m_k}) + (1 - \vartheta_{m_k})z_{m_k} - u_{m_k}\| \\ &= \|\vartheta_{m_k}[g(u_{m_k}) - u_{m_k}] + (1 - \vartheta_{m_k})[z_{m_k} - u_{m_k}]\| \\ &\leq \vartheta_{m_k} \|g(u_{m_k}) - u_{m_k}\| + (1 - \vartheta_{m_k}) \|z_{m_k} - u_{m_k}\| \\ &\rightarrow 0. \end{aligned} \quad (3.61)$$

We have to use the same justification as in the Case 1 such that

$$\limsup_{k \rightarrow +\infty} \langle g(\omega^*) - \omega^*, u_{m_k+1} - \omega^* \rangle \leq 0. \quad (3.62)$$

By the use of expressions (3.55) and (3.57), we have

$$\begin{aligned} &\|u_{m_k+1} - \omega^*\|^2 \\ &\leq [1 - 2\vartheta_{m_k}(1 - \xi)] \|u_{m_k} - \omega^*\|^2 \\ &\quad + 2\vartheta_{m_k}(1 - \xi) \left[\frac{\vartheta_{m_k} \|u_{m_k} - \omega^*\|^2}{2(1 - \xi)} + \frac{\langle g(\omega^*) - \omega^*, u_{m_k+1} - \omega^* \rangle}{1 - \xi} \right] \\ &\leq [1 - 2\vartheta_{m_k}(1 - \xi)] \|u_{m_k+1} - \omega^*\|^2 \\ &\quad + 2\vartheta_{m_k}(1 - \xi) \left[\frac{\vartheta_{m_k} \|u_{m_k} - \omega^*\|^2}{2(1 - \xi)} + \frac{\langle g(\omega^*) - \omega^*, u_{m_k+1} - \omega^* \rangle}{1 - \xi} \right]. \end{aligned} \quad (3.63)$$

It follows that

$$\|u_{m_k+1} - \omega^*\|^2 \leq \frac{\vartheta_{m_k} \|u_{m_k} - \omega^*\|^2}{2(1 - \xi)} + \frac{\langle g(\omega^*) - \omega^*, u_{m_k+1} - \omega^* \rangle}{1 - \xi}. \quad (3.64)$$

Since $\vartheta_{m_k} \rightarrow 0$, as $k \rightarrow +\infty$ and $\|u_{m_k} - \omega^*\|$ is a bounded sequence, it follows from (3.62) and (3.64) that

$$\|u_{m_k+1} - \omega^*\|^2 \rightarrow 0, \text{ as } k \rightarrow +\infty. \quad (3.65)$$

The above expression with (3.57) implies that

$$\lim_{k \rightarrow +\infty} \|u_k - \omega^*\|^2 \leq \lim_{k \rightarrow +\infty} \|u_{m_k+1} - \omega^*\|^2 \leq 0. \quad (3.66)$$

Consequently, we have $u_n \rightarrow \omega^*$ as $n \rightarrow +\infty$. This completes the proof of the theorem. \square

4. NUMERICAL ILLUSTRATION

This section describes the numerical performance of the proposed algorithms, in contrast to some related work in the literature, as well as the analysis of how variations in control parameters affect the numerical effectiveness of the proposed algorithms.

All computations are done in MATLAB R2018b and run on HP i-5 Core(TM) i5-6200 8.00 GB (7.78 GB usable) RAM laptop.

Example 4.1. Let $\Sigma = l_2$ be a real Hilbert space with the sequences of real numbers satisfying the following condition:

$$\|u_1\|^2 + \|u_2\|^2 + \cdots + \|u_n\|^2 + \cdots < +\infty. \quad (4.1)$$

Assume that a mapping $\Gamma : \Delta \rightarrow \Delta$ is defined by

$$\Gamma(u) = (5 - \|u\|)u, \quad \forall u \in \Sigma,$$

where $\Delta = \{u \in \Sigma : \|u\| \leq 3\}$. Then, we can easily see that Γ is weakly sequentially continuous on Σ and the solution set is $VI(\Delta, \Gamma) = \{0\}$.

For any $u, y \in \Sigma$, we have

$$\begin{aligned} \|\Gamma(u) - \Gamma(y)\| &= \|(5 - \|u\|)u - (5 - \|y\|)y\| \\ &= \|5(u - y) - \|u\|(u - y) - (\|u\| - \|y\|)y\| \\ &\leq 5\|u - y\| + \|u\|\|u - y\| + \|\|u\| - \|y\|\|\|y\| \\ &\leq 5\|u - y\| + 3\|u - y\| + 3\|u - y\| \\ &\leq 11\|u - y\|. \end{aligned} \quad (4.2)$$

Hence Γ is L -Lipschitz continuous with $L = 11$.

And, for any $u, y \in \Sigma$ and let $\langle \Gamma(u), y - u \rangle > 0$, such that

$$(5 - \|u\|)\langle u, y - u \rangle > 0.$$

Since $\|u\| \leq 3$ and it implies that

$$\langle u, y - u \rangle > 0.$$

Consider that

$$\begin{aligned} \langle \Gamma(y), y - u \rangle &= (5 - \|y\|)\langle y, y - u \rangle \\ &\geq (5 - \|y\|)\langle y, y - u \rangle - (5 - \|y\|)\langle u, y - u \rangle \\ &\geq 2\|u - y\|^2 \geq 0. \end{aligned} \quad (4.3)$$

Hence a mapping Γ is quasimonotone on Δ .

Let $u = (\frac{5}{2}, 0, 0, \dots, 0, \dots)$ and $y = (3, 0, 0, \dots, 0, \dots)$ such that

$$\langle \Gamma(u) - \Gamma(y), u - y \rangle = \left(\frac{5}{2} - 3\right)^3 < 0.$$

Let considered the following projection formula:

$$P_{\Delta}(u) = \begin{cases} u & \text{if } \|u\| \leq 3, \\ \frac{3u}{\|u\|}, & \text{otherwise.} \end{cases}$$

Figure 1–2 and Table 1 show numerical results. The control conditions are taken in the following way:

- (i) Algorithm 3.1 (shortly, **Alg11**): $\varkappa = \frac{0.7}{L}, \vartheta_n = \frac{1}{(n+2)}, g(u) = \frac{u}{2}, D_n = \|u_n - u_y\|;$
- (ii) Algorithm 3.2 (shortly, **Alg22**): $\varkappa_1 = 0.22, \chi = 0.44, \vartheta_n = \frac{1}{(n+2)}, g(u) = \frac{u}{2}, D_n = \|u_n - y_n\|;$
- (iii) Algorithm 3.4 (shortly, **Alg33**): $\varkappa_1 = 0.22, \chi = 0.44, \varphi_n = \frac{100}{(n+1)^2}, \vartheta_n = \frac{1}{(n+2)}, g(u) = \frac{u}{2}, D_n = \|u_n - y_n\|.$




FIGURE 1. Numerical illustration of Algorithm 3.1 and Algorithm 3.2 with Algorithm 3.4 while $u_1 = (1, 1, \dots, 1_{5000}, 0, 0, \dots)$.




FIGURE 2. Numerical illustration of Algorithm 3.1 and Algorithm 3.2 with Algorithm 3.4 while $u_1 = (1, 1, \dots, 1_{5000}, 0, 0, \dots)$.

TABLE 1. Numerical explanation for Example 4.1.

u_1	Number of iterations			Execution time in seconds		
	Alg11	Alg22	Alg33	Alg11	Alg22	Alg33
$(1, 1, \dots, 1_{5000}, 0, 0, \dots)$	35	33	27	2.0040884	2.2295634	1.8277622
$(1, 2, \dots, 10_{10000}, 0, 0, \dots)$	37	34	27	2.2736534	2.5582612	1.9910394
$(10, 10, \dots, 10_{10000}, 0, 0, \dots)$	43	39	34	3.2163693	2.9575764	2.2758355
$(20, 20, \dots, 20_{10000}, 0, 0, \dots)$	49	43	38	4.5876948	3.8674743	3.1847685

CONCLUSION

We developed various modified extragradient type methods to provide a numerical solution to quasimonotone variational inequality problems in real

Hilbert space. Despite the fact that each sequence is generated by a different step size rule, all sequences generated by the proposed method are strongly convergent to the solution. The numerical results are given to demonstrate the numerical effectiveness of our algorithm. These numerical investigations have demonstrated that the variable step size impacts the efficiency of the iterative sequence in this context.

Acknowledgements: Nattawut Pholasa would like to thank University of Phayao and Thailand Science Research and Innovation grant no. FF65-RIM072 and FF65-UoE001. Nuttapol Pakkaranang would like to thank Phetchabun Rajabhat University.

REFERENCES

- [1] A.S. Antipin, *On a method for convex programs using a symmetrical modification of the Lagrange function*, *Economika i Matem. Metody.*, **12** (1976), 1164–1173.
- [2] H.H. Bauschke and P.L. Combettes, *Convex analysis and monotone operator theory in Hilbert spaces*, Springer, New York, 2011.
- [3] L.C. Ceng, A. Petruşel and J.C. Yao, *On mann viscosity subgradient extragradient algorithms for fixed point problems of finitely many strict pseudocontractions and variational inequalities*, *Mathematics*, **7** (2019), Article ID 925.
- [4] L.C. Ceng and M. Shang, *Hybrid inertial subgradient extragradient methods for variational inequalities and fixed point problems involving asymptotically nonexpansive mappings*, *Optimization*, **70** (2019), 715–740.
- [5] L.C. Ceng, X. Qin, Y. Shehu and J.C. Yao, *Mildly inertial subgradient extragradient method for variational inequalities involving an asymptotically nonexpansive and finitely many nonexpansive mappings*, *Mathematics*, **7** (2019), Article ID 881.
- [6] Y. Censor, A. Gibali and S. Reich, *The subgradient extragradient method for solving variational inequalities in Hilbert space*, *J. Optim. Theory Appl.*, **148** (2011), 318–335.
- [7] Y. Censor, A. Gibali and S. Reich, *Extensions of korpelevich extragradient method for the variational inequality problem in euclidean space*, *Optimization*, **61** (2012), 1119–1132.
- [8] C.M. Elliott, *Variational and quasivariational inequalities: applications to free-boundary problems (Claudio Baiocchi and antônio Capelo)*, *SIAM Rev.*, **29**(2) (1987), 314–315.
- [9] H.A. Hammad, H. Rehman and M.D. la Sen, *Advanced algorithms and common solutions to variational inequalities*, *Symmetry*, **12** (2020), Article ID 1198.
- [10] A.N. Iusem and B.F. Svaiter, *A variant of korpelevich's method for variational inequalities with a new search strategy*, *Optimization*, **42** (1997), 309–321.
- [11] G. Kassay, J. Kolumbán and Z. Páles, *On nash stationary points*, *Publ. Math. Debrecen*, **54**(3-4) (1999), 267–279.
- [12] G. Kassay, J. Kolumbán and Z. Páles, *Factorization of minty and stampanchia variational inequality systems*, *European J. Oper. Res.*, **143** (2002), 377–389.
- [13] J.K. Kim and Salahuddin, *Extragradient methods for generalized mixed equilibrium problems and fixed point problems in Hilbert spaces*, *Nonlinear Funct. Anal. Appl.*, **22**(4) (2017), 693–709.
- [14] D. Kinderlehrer and G. Stampacchia, *An introduction to variational inequalities and their applications*, *Classic in Applied Mathematics*, Society for Industrial and Applied Mathematics, (SIAM), Philadelphia, PA, 2000.
- [15] I. Konnov, *Equilibrium models and variational inequalities*, **210**, Elsevier, 2007.

- [16] G. Korpelevich, *The extragradient method for finding saddle points and other problems*, *Ékonom. i Mat. Metody*, **12**(4) (1976), 747–756.
- [17] Z. Liu, S. Migórski and S. Zeng, *Partial differential variational inequalities involving nonlocal boundary conditions in Banach spaces*, *J. Differential Equ.*, **263** (2017), 3989–4006.
- [18] Z. Liu, S. Zeng and D. Motreanu, *Evolutionary problems driven by variational inequalities*, *J. Differential Equ.*, **260** (2016), 6787–6799.
- [19] P.E. Maingé, *Strong convergence of projected subgradient methods for nonsmooth and nonstrictly convex minimization*, *Set-Valued Anal.*, **16** (2008), 899–912.
- [20] A. Moudafi, *Viscosity approximation methods for fixed-points problems*, *J. Math. Anal. Appl.*, **241** (2000), 46–55.
- [21] K. Muangchoo, H. Rehman and P. Kumam, *Two strongly convergent methods governed by pseudo-monotone bi-function in a real Hilbert space with applications*, *J. Appl. Math. Comput.*, **67**(1-2) (2021), 891–917.
- [22] K. Muangchoo, *A Viscosity type Projection Method for Solving Pseudomonotone Variational Inequalities*, *Nonlinear Funct. Anal. Appl.*, **26**(2) (2021), 347–371.
- [23] A. Nagurney, *Network economics: a variational inequality approach*, Kluwer Academic Publishers Group, Dordrecht, 1993. <https://doi.org/10.1007/978-94-011-2178-1>
- [24] M.A. Noor, *Some iterative methods for nonconvex variational inequalities*, *Math. Comput. Modelling*, **54** (11-12) (2011), 2955–2961.
- [25] H. Rehman, A. Gibali, P. Kumam and K. Sitthithakerngkiet, *Two new extragradient methods for solving equilibrium problems*, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat.*, **115**(2) (2021), Article ID 75.
- [26] H. Rehman, P. Kumam, A.B. Abubakar and Y.J. Cho, *The extragradient algorithm with inertial effects extended to equilibrium problems*, *Comput. Appl. Math.*, **39**(2) (2020), Article ID 100.
- [27] H. Rehman, P. Kumam, I.K. Argyros, W. Deebani and W. Kumam, *Inertial extragradient method for solving a family of strongly pseudomonotone equilibrium problems in real Hilbert spaces with application in variational inequality problem*, *Symmetry*, **12** (2020), Article ID 503.
- [28] H. Rehman, P. Kumam, Y.J. Cho, Y.I. Suleiman and W. Kumam, *Modified popov's explicit iterative algorithms for solving pseudomonotone equilibrium problems*, *Optim. Methods Softw.*, **36**(1) (2021), 82–113.
- [29] H. Rehman, P. Kumam, Y.J. Cho and P. Yordsorn, *Weak convergence of explicit extragradient algorithms for solving equilibrium problems*, *J. Inequal. Appl.*, **2019** (2019), Article ID 282.
- [30] H. Rehman, P. Kumam, Q.L. Dong and Y.J. Cho, *A modified self-adaptive extragradient method for pseudomonotone equilibrium problem in a real Hilbert space with applications*, *Math. Methods Appl. Sci.*, **44**(5) (2020), 3527–3547.
- [31] H. Rehman, P. Kumam, A. Gibali and W. Kumam, *Convergence analysis of a general inertial projection-type method for solving pseudomonotone equilibrium problems with applications*, *J. Inequal. Appl.*, **2021** (2021), Article ID 63.
- [32] H. Rehman, P. Kumam, W. Kumam, M. Shutaywi and W. Jirakitpuwapat, *The inertial sub-gradient extra-gradient method for a class of pseudo-monotone equilibrium problems*, *Symmetry*, **12**(3) (2020), Article ID 463.
- [33] H. Rehman, N. Pakkaranang, A. Hussain and N. Wairojjana, *A modified extra-gradient method for a family of strongly pseudomonotone equilibrium problems in real Hilbert spaces*, *J. Math. Comput. Sci.*, **22**(1) (2020), 38–48.

- [34] G. Stampacchia, *Formes bilinéaires coercitives sur les ensembles convexes*, C. R. Acad. Sci. Paris, **258** (1964), 4413–4416.
- [35] P. Tseng, *A modified forward-backward splitting method for maximal monotone mappings*, SIAM J. Control Optim., **38**(2) (2000), 431–446.
- [36] N. Wairojjana and N. Pakkaranang, *Halpern Tseng's extragradient methods for solving variational inequalities involving semistrictly quasimonotone operator*, Nonlinear Funct. Anal. Appl., **27**(1) (2022), 121–140.
- [37] H.K. Xu, *Another control condition in an iterative method for nonexpansive mappings*, Bull. Austral. Math. Soc., **65**(1) (2002), 109–113.
- [38] J. Yang and H. Liu, *A modified projected gradient method for monotone variational inequalities*, J. Optim. Theory Appl., **179** (1) (2018), 197–211.
- [39] J. Yang, H. Liu and Z. Liu, *Modified subgradient extragradient algorithms for solving monotone variational inequalities*, Optimization, **67** (2018), 2247–2258.
- [40] P. Yordsorn, P. Kumam, H. Rehman and A.H. Ibrahim, *A weak convergence self-adaptive method for solving pseudomonotone equilibrium problems in a real Hilbert space*, Mathematics, **8** (2020), Article ID 1165.
- [41] L. Zhang, C. Fang and S. Chen, *An inertial subgradient-type method for solving single-valued variational inequalities and fixed point problems*, Numer. Algorithms, **79**(3) (2018), 941–956.
- [42] T.Y. Zhao, D.Q. Wang, L.C. Ceng, L. He, C.Y. Wang and H.L. Fan, *Quasi-inertial Tseng's extragradient algorithms for pseudomonotone variational inequalities and fixed point problems of quasi-nonexpansive operators*, Numer. Funct. Anal. Optim., **42**(1) (2021), 69–90.