



Galectin-1 from redlip mullet *Liza haematocheilia*: identification, immune responses, and functional characterization as pattern recognition receptors (PRRs) in host immune defense system

Chaehyeon Lim¹, Hyukjae Kwon^{1,2}, Jehee Lee^{1,2,*}

¹ Department of Marine Life Sciences & Fish Vaccine Research Center, Jeju National University, Jeju 63243, Korea

² Marine Science Institute, Jeju National University, Jeju 63333, Korea

Abstract

Galectins, a family of β -galactoside-binding lectins, have emerged as soluble mediators in infected cells and pattern recognition receptors (PRRs) responsible for evoking and regulating innate immunity. The present study aimed to evaluate the role of galectin-1 in the host immune response of redlip mullet (*Liza haematocheilia*). We established a cDNA database for redlip mullet, and the cDNA sequence of galectin-1 (LhGal-1) was characterized. In silico analysis was performed, and the spatial and temporal expression patterns in gills and blood in response to lipopolysaccharide polyinosinic:polycytidylic acid, and *Lactococcus garvieae* were estimated via quantitative real-time PCR. Functional assays were conducted using recombinant protein to investigate carbohydrate binding, bacterial binding, and bacterial agglutination activity. LhGal-1 was composed of 135 amino acids. Conserved motifs (H-NPR, -N- and -W-E-R) within the carbohydrate recognition domain were found in LhGal-1. The tissue distribution revealed that the healthy stomach expressed high levels of LhGal-1. The temporal monitoring of *LhGal-1* mRNA expression in the gill and blood showed its significant upregulation in response to immune challenges with different stimulants. rLhGal-1 exhibited binding activity in response to carbohydrates and bacteria. Moreover, the agglutination of rLhGal-1 against *Escherichia coli* was observed. Collectively, our findings suggest that LhGal-1 may function as a PRR in redlip mullet. Furthermore, *LhGal-1* can be considered a significant gene to play a protective role in redlip mullet immune system.

Keywords: Galectin-1, Carbohydrate recognition domain, mRNA expression pattern, Pattern recognition receptor, Redlip mullet

Introduction

The innate immune system constitutes the first line of defense

against pathogens penetrating marine organisms. Pattern recognition receptors (PRRs) initiate appropriate functions of the innate immune system and play vital roles in recognizing patho-

Received: Jun 28, 2022 Revised: Oct 17, 2022 Accepted: Oct 19, 2022

*Corresponding author: Jehee Lee

Department of Marine Life Sciences & Fish Vaccine Research Center, Jeju National University, Jeju 63243, Korea

Tel: +82-64-754-3472, E-mail: jehee@jejunu.ac.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2022 The Korean Society of Fisheries and Aquatic Science

gen-associated molecular patterns (PAMPs) present on the surface of microbial pathogens (Kumagai et al., 2008). Among the PRRs, lectins are a group of carbohydrate-binding proteins that are primarily involved in pathogen recognition and triggering proinflammatory responses via protein-carbohydrate interactions (Vasta et al., 2011). Lectins are broadly classified into C-type, F-type, P-type, ficolins, pentraxins, and galectins.

Galectins are a conserved family of carbohydrate-binding proteins that include at least one carbohydrate recognition domain (CRD). These are widely distributed in mammals, birds, amphibians, fish, nematodes, sponges, and some fungi. In contrast to other lectins, galectins function in a Ca^{2+} -independent manner and have two basic properties; the characteristic affinity for β -galactosides, and the presence of a conserved CRD sequence motif. Evolutionarily conserved galectins have been classified into proto, chimera, and tandem-repeat types based on the structural differences of CRDs (Hirabayashi & Kasai, 1993). Proto-type galectins include one CRD with non-covalently linked homodimers. Chimera-type galectins comprise a C-terminal CRD and an N-terminal domain rich in proline and glycine. Tandem-repeat type galectins are embodied with two CRDs and a functional linker peptide that is required for joining the two domains (Vasta, 2012).

Galectin is a multifunctional protein involved in processes of cell development and migration, agglutination of erythrocytes and bacteria, and the regulation of adaptive immune responses. Previous studies reported that galectins act as mediators in developmental processes including cell differentiation, tissue organization, and regulation of immune homeostasis by binding endogenous (self) glycans (Leffler et al., 2002). Moreover, galectins can bind to exogenous (non-self) glycans present on the surface of viruses, bacteria, parasites, and fungi, as well as other PAMPs, as PRRs in the innate immune system (Chen et al., 2014; Shi et al., 2018; Wang et al., 2020).

Redlip mullets (*Liza haematocheilia*) belong to the Mugilidae family, which consists of more than 72 species from 17 fish genera distributed worldwide (Minos et al., 2009). Among them, redlip mullet is the only species being cultured and is regarded as a valuable aquaculture species in South Korea, China, Japan, and the Northwest Pacific Ocean. The statistics published by the Korean Ministry of Maritime Affairs and Fisheries have shown that collectively, mullets account for 8% of the total consumption and cultivation in Korea (Ministry of Maritime Affairs and Fisheries, 2020). Recently, the high mortality of the redlip mullet in Korea was attributed to *Lactococcus garvieae*, a gram-positive bacterium

that causes green liver syndrome in these fish (Han et al., 2015). However, the knowledge of the host immune defense mechanism of redlip mullet is not known in adequate detail to control the outbreak of this disease. Therefore, identification and functional characterization of immune-related genes in redlip mullet are essential for their successful breeding.

In this study, we performed functional characterization and transcriptional profiling to investigate the potential effect of cDNA sequence of galectin-1 (LhGal-1) on the immune system of redlip mullet. Galectin-1 was identified from the redlip mullet (*L. haematocheilia*) cDNA database, and its spatial and temporal mRNA expression patterns were analyzed in tissues following immunogenic challenge with lipopolysaccharide (LPS), polyinosinic:polycytidylic acid (poly I:C) and *L. garvieae*. Additionally, the carbohydrate-binding ability of recombinant LhGal-1 was examined using lactose, galactose, and glucose. Furthermore, microbial binding assay and agglutination assay were performed for the functional characterization of LhGal-1 as a PRR.

Materials and Methods

Fish rearing, immune challenge, and tissue isolation

All animal experiments were performed with the approval of the Institutional Animal Care and Use Committee at Jeju National University (approval no. 2018-0003). Healthy redlip mullets were obtained from the Sangdeok fishery in Hadong, South Korea, with average body weight and length of 146.5 ± 29.3 and 23.5 ± 1.7 , respectively. Selected fish were reared at 20°C in 300 L tanks for seven days before the experiment.

Following anesthetization with tricaine mesylate (MS-222, 40 mg/L), five healthy redlip mullets were anatomized to collect the kidneys, spleen, gill, intestine, stomach, heart, blood, liver, muscle, skin, brain, and head kidney. A blood sample was drawn from their caudal vein using sterile syringes coated with heparin sodium salt (USB, Thermo Scientific, Waltham, MA, USA) and peripheral blood cells were collected by centrifugation at $3,000\times g$, 4°C for 10 min. All tissues were carefully isolated, frozen immediately in liquid nitrogen, and stored at -80°C until RNA extraction.

For the immune stimulation experiment, healthy mullets were divided into three immune stimulant groups and one control group; LPS ($1.25 \mu\text{g/g}$, from *Escherichia coli* 055:B5; Sigma-Aldrich, St. Louis, MO, USA), poly I:C ($1.5 \mu\text{g/g}$, Sigma-Aldrich), *L. garvieae* (strain: JJJN1, accession no.: CP026502.1, 1

$\times 10^3$ CFU/ μ L) were prepared in $1 \times$ phosphate-buffered saline (PBS). In different experimental groups, each fish was intraperitoneally injected with 100 μ L of each stimulant, whereas an equivalent volume of PBS was injected to the fish of the control group. Subsequently, five fish were selected from each group and dissected 0, 6, 24, 48, 72 h post injection to isolate tissues, as described above.

Total RNA extraction and cDNA synthesis

Total RNA was extracted from the collected tissues ($n = 5$) to study the tissue-specific distribution of LhGal-1 in healthy fish, as well as upon immune challenge. RNAiso plus kit (TaKaRa, Kusatsu, Japan) was used to isolate RNA which was subsequently purified with RNeasy spin column (Qiagen, Germantown, MD, USA) according to the manufacturer's instructions. The concentration and quality of purified RNA were determined with μ Drop Plate (Thermo Scientific) at 260 nm and via 1.5% agarose gel electrophoresis, respectively. Thereafter, cDNA was synthesized using the PrimeScript™ II 1st strand cDNA synthesis kit (TaKaRa) in the final reaction volume of 20 μ L containing 2.5 μ g of RNA. The synthesized cDNA samples were diluted 40 folds and stored at -80°C .

Quantitative real-time PCR (qPCR)

For analyzing the tissue distribution and temporal distribution of *LhGal-1* mRNA expression in different tissues following immune challenge, quantitative real-time PCR (qPCR) (TaKaRa Thermal Cycler Dice: TP850 Real-Time System, TaKaRa) was performed using cDNA as a template and gene-specific primers (Table 1). The gene encoding mullet elongation factor 1 α (*LhEF1 α* ; accession no: MH017208) served as the internal control (Table 1). The 10 μ L reaction mixture contained 3 μ L of cDNA, 5 μ L of 2 \times SYBR® Premix Ex Taq™, 0.4 μ L of each primer (10 pmol/ μ L), and 1.2 μ L of nuclease-free water. Each sample

of cDNA was tested in triplicate. The following reaction conditions were used: initial denaturation at 95°C for 10 sec; 45 cycles of 95°C for 5 sec, 58°C for 10 sec, and 72°C for 20 sec; a final cycle of 95°C for 15 sec, 60°C for 30 sec, and 95°C for 15 sec.

Relative *LhGal-1* mRNA expression levels were determined using the Livak ($2^{-\Delta\Delta\text{CT}}$) method (Livak & Schmittgen, 2001). The Ct values of all the samples were normalized to those of the *LhEF1 α* gene. The Ct values in the tissues with the lowest expression of *LhGal-1* were used for normalization when estimating the tissue-specific distribution. The data for the immune-challenged groups are shown as the relative fold-change compared to the PBS-injected group and calculated as the mean \pm SD at a significance level of $p < 0.05$.

Identification and *in silico* analysis of *LhGal-1* sequences

The redlip mullet cDNA database was constructed using the PacBio platform. All samples were sequenced using the Iso-Seq method (full-length mRNA sequencing technology) after determining the concentration and quality of RNA. For the library preparation, total RNA was isolated from blood, spleen, and head kidney and reverse transcribed into full-length first-strand cDNA using the Clontech SMARTer PCR cDNA Synthesis Kit (TaKaRa). The amplified cDNA was fractionated by size and converted into SMRTbell templates for sequencing. Finally, the full-length transcript isoforms were annotated using Blast-2Go software (Liyanage et al., 2018). The cDNA sequence of *LhGal-1* was identified using the Basic Local Alignment Search Tool (Altschul et al., 1990) available at the National Center for Biotechnology Information (NCBI) website and compared with other known orthologous genes sequences (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). The open reading frames (ORFs) and translated amino acid sequences were determined using the online software ORF finder from NCBI (<https://www.ncbi.nlm.nih.gov/orffinder/>). Then, the NCBI Conserved Domain

Table 1. Summary of primers used in this study

Primer sequence 5'-3'	Description	Amplicon size	Tm ($^\circ\text{C}$)
AGGCTGGTATCTCCAAGAA	Elongation factor 1 α (LhEF1 α), qPCR Forward	130 bp	60
TAACGGGACTGGCTGTAA	Elongation factor 1 α (LhEF1 α), qPCR Reverse	130 bp	60
GTCGCTAAACCTGACGCTTCCAAC	LhGal-1, qPCR, Forward	132 bp	60
ATTGTGTGCAACTCTACCAGGGA	LhGal-1, qPCR, Reverse	132 bp	60
GAGAGAgatatacATGATGAAAGATCTGATCATAAAGAACATGTCCTTTAAGGT	LhGal-1, pMAL-c5X, EcoRV, Forward	408 bp	60.3
GAGAGAggatccTTACTTGATCTCAAAGCTTCTGATGCGAGC	LhGal-1, pMAL-c5X, BamHI, Reverse	408 bp	60.3

qPCR, quantitative real-time PCR; Tm, melting temperature.

Search (<https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi>), InterPro-EMBL-EBI (<https://www.ebi.ac.uk/interpro/>), and ExPASy-PROSITE (<https://prosite.expasy.org/>) were used to analyze the extrapolated protein domain sequences of LhGal-1. Besides, SignalP-4.1 server (<https://services.healthtech.dtu.dk/service.php?SignalP-4.1>) was used to ascertain the location and potential of the signaling peptide in the LhGal-1 sequence. Based on the sequence, the primers for cloning and qPCR were designed by Integrated DNA Technologies (<https://sg.idtdna.com/pages>), following MIQE guidelines (Bustin et al., 2009). Multiple sequence alignment was performed using CLC Main Workbench software version 8.0.1 (<https://digitalinsights.qiagen.com>) to determine the conserved domains and residues. Moreover, the Molecular Evolutionary Genetics Analysis (MEGA) software version 10.2 was used to perform phylogenetic analysis after using the neighbor-joining method with 5,000 bootstrap replicates to determine speciation and evolutionary relationships based on the similarities and differences within the molecular characteristics of different species (Kumar et al., 2018). Pairwise sequence alignment was performed using EMBOSS Needle pairwise sequence alignment software to determine the similarities and identities of the orthologs (https://www.ebi.ac.uk/Tools/psa/emboss_needle/). The 3D structural model of LhGal-1 was inferred from the amino acid sequence using Iterative Threading ASSEmblY Refinement (I-TASSER), which provides protein structure and function predictions based on the sequence-to-structure-to-function model (Roy et al., 2010).

Cloning and purification of recombinant LhGal-1

The pMAL™ Protein Fusion & Purification System (New England Biolabs, Ipswich, MA, USA) was used to obtain a purified recombinant LhGal-1 protein (rLhGal-1). The ORF of LhGal-1 was cloned into the pMAL vector to append it to a sequence encoding the maltose-binding protein.

To clone the ORF of LhGal-1, target sequences were amplified via PCR with primers designed to include the corresponding restriction sites (Table 1). The 50 µL reaction mixture contained 5 µL of 10X ExTaq buffer, 4 µL of 2.5 mM dNTP, 10 pmol of each primer, 5 µL cDNA template, 0.2 µL ExTaq™ DNA polymerase (TaKaRa), and 33.8 µL nuclease-free water. The PCR was run on a thermal cycler (TaKaRa) under the following conditions: initial denaturation at 94°C for 5 min, 35 cycles of denaturation at 94°C for 30 sec, annealing at 59°C for 30 sec, and extension at 72°C for 40 sec, and final extension at 72°C for 7 min. Following PCR product purification, the pMAL-

c5X vector (New England Biolabs) and inserts were digested using *EcoRI* and *EcoRV* followed by ligation using the DNA Ligation Kit (Mighty Mix) (TaKaRa). For the mass production of recombinant plasmid DNA, the *E. coli* DH5α competent cells were transformed and grown. The recombinant plasmid was extracted from the grown cells using the AccuPrep® Plasmid Mini Extraction Kit (Bioneer, Daejeon, Korea), and the samples were sent to Macrogen, Korea for sequencing.

The recombinant protein was overexpressed using isopropyl-β-thiogalactopyranoside (IPTG) induction. Briefly, transformed *E. coli* ER2523 cells were cultured in 500 mL Luria-Bertani (LB) broth containing 500 µL of ampicillin (100 µg/mL) and 0.2% glucose at 37°C. When the optical density at 600 nm (OD₆₀₀) reached 0.6, IPTG (final concentration of 1 mM) was added to induce the expression of rLhGal-1 fusion proteins. The culture was further grown for 8 h at 25°C and 200 rpm. Next, the induced cells were harvested by centrifugation (3,500×g for 15 min at 4°C), washed twice with column buffer (20 mM Tris-HCL, pH 7.4 and 200 mM NaCl), and stored overnight at -20°C. For cell lysis, the cells were thawed in cold water and sonicated on ice. The lysate was centrifuged at 17,000×g for 30 min at 4°C. rLhGal-1 was purified using the pMAL™ Protein Fusion & Purification System (New England Biolabs) per the manufacturer's instructions. Finally, the concentration of purified protein was measured using the Bradford assay and sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was performed with 12% polyacrylamide gel to confirm the size and purity of the target fusion proteins (Bradford, 1976). The eluted protein was stored at -80°C until further analysis.

Sugar-binding assay

The sugar-binding ability of rLhGal-1 was examined by performing an enzyme-linked immunosorbent assay (ELISA) and using three different carbohydrates as follows: α-lactose, D-galactose, and D-glucose (47287-U, G0750, G8270, Sigma-Aldrich). In brief, lactose, galactose, and glucose were each dissolved in carbonate-bicarbonate buffer (50 mmol/L, pH 9.6) to a final concentration of 100 mM. The 96-well microtiter plate was coated with the carbohydrate solutions and incubated at 4°C overnight. After washing three times with Tris-buffered saline with 0.1% Tween 20 (TBS-T), the wells were blocked with 5% skim milk in TBS-T. The plate was washed five times with TBS-T and treated with 50 µL of serially diluted rLhGal-1 or maltose-binding protein (MBP). After incubating for 1 h at 37°C and 10 rpm, the wells were washed three times with

TBS-T, following which, 100 μ L of mouse anti-MBP antibody (1:5,000, New England Biolabs) was added as the primary antibody. The plate was incubated at 37°C for 2 h and washed again with TBS-T. It was then incubated with the secondary antibody (100 μ L goat anti-mouse IgG horseradish peroxidase conjugate; 1:3,000; YOUNG IN Frontier, Seoul, Korea), under the same conditions used for the primary antibody. Finally, the plate was washed five times with TBS-T and incubated with tetramethylbenzidine solution at room temperature for 5 to 10 min in the dark. For terminating the reaction, 50 μ L of stop solution (1 M H₂SO₄) was added to each well and the absorbance was measured at 450 nm using a Multiskan Sky Microplate Spectrophotometer (Thermo Fisher Scientific).

Bacterial binding assay

To determine the microbial binding affinity of rLhGal-1, ELISA was performed with seven bacterial strains as follows: three gram-positive (G+) species (*L. garvieae*, *Streptococcus iniae*, and *Streptococcus parauberis*) and four gram-negative (G-) species (*E. coli*, *Edwardsiella tarda*, *Vibrio anguillarum*, and *Vibrio harveyi*). Briefly, the seven bacteria were cultured in appropriate media overnight and harvested by centrifugation; LB broth was used for growing *E. coli* and *V. anguillarum*, and brain heart infusion broth with 1.5% NaCl was used for growing *L. garvieae*, *S. iniae*, *S. parauberis*, *E. tarda*, and *V. harveyi*. Harvested cells were washed twice and resuspended in PBS until an OD₆₀₀ of 1 was achieved and diluted with carbonate-bicarbonate buffer (50 mmol/L, pH 9.6) to achieve the bacterial count of 1×10^8 cells/mL. The 96-well plate was coated with 100 μ L of bacterial cell suspension diluted in coating buffer (1×10^7 cells/well) and incubated overnight at 4°C. ELISA was performed as described for the sugar-binding assay.

Agglutination assay of rLhGal-1

To examine the agglutination ability of rLhGal-1, the agglutination assay was performed using *E. coli*. Briefly, *E. coli* was cultured in LB broth at 37°C overnight and harvested by centrifugation. Harvested cells were washed twice with TBS (20 mM Tris, 137 mM NaCl, pH 7.8) and 50 μ L of cell suspension was plated into the individual wells of a 96-well plate (5×10^6 CFU/mL). Each well was treated with 10 μ g of rLhGal-1 and incubated at 25°C for 1 h. The control group was treated with an equal amount of MBP. Bacterial cells were observed under a light microscope to determine agglutination. All assays were performed in triplicate.

Statistical analysis

All experiments were performed in triplicates and the data are presented as the mean \pm SD. The Student's *t*-test was used to compare the mean values of control and challenge groups and determine the statistical significance of the results. One-way analysis of variance was used to examine the significant differences within the groups. Results with $p < 0.05$ were considered statistically significant.

Results and Discussion

Sequence characterization

LhGal-1 was identified from the constructed redlip mullet transcriptome database and characterized by sequencing and using several bioinformatics tools. The length of *LhGal-1* (GenBank accession No: MW147017) ORF was 408 bp and encoded a protein of 135 amino acids with a predicted molecular weight of 15.31 kDa, and a theoretical isoelectric points of 4.89 (Fig. 1). The signal 4.1 server predicted a lack of signal peptides in the whole amino acid sequence, suggesting that these molecules might be secreted via a non-classical secretory pathway similar to other galectins (Boulianne et al., 2000).

The amino acid sequence of LhGal-1 harbors a single GLECT CRD (Letunic & Bork, 2018). *LhGal-1* sequence can be considered as the proto-type of galectins (Vasta, 2012). According to our InterPro analysis, GLECT domains of LhGal-1 contained eight conserved residues of sugar-binding pockets for β -galactoside (45H, 47N, 49R, 60V, 62N, 69W, 72E, and 74R) (Fig. 2). Although LhGal-1 has four cysteine residues (61C, 70C, 76C, and 77C), a disulfide bond was not identified by the ScanProsite software (Fig. 2). The predicted 3D model of LhGal-1 revealed that it is folded as a β sandwich comprising two anti-parallel β -sheet bundles (Fig. 2). Furthermore, five strands of each β sheet bundle were slightly bent and shaped to generate a concave and a convex side. All the main residues for carbohydrate binding were positioned on the concave side of the structure. This groove-like formation may allow the holding of linear tetra-saccharide long enough to form a sugar-binding pocket (Leffler et al., 2002).

Multiple sequence alignment showed that LhGal-1 harbored highly conserved motifs within the CRD (H-NPR, -N- and -W-E-R) that are crucial and typical residues of the galectin family, related to carbohydrate-binding affinity (Fig. 3). Moreover, the main residues required for carbohydrate-binding of LhGal-1 were highly conserved with other galectins from fishes,

1	ATG ATG AAA GAT CTG ATC ATA AAG AAC ATG TCC TTT AAG GTC GGA	45
1	M M K D L I I K N M S F K V G	15
46	CAA ACA ATG ACC ATT GTT GGA GTC GCT AAA CCT GAC GCT TCC AAC	90
16	Q T M T I V G V A K P D A S N	30
91	TTT GCA TTC AAC ATC GGC CCC AGT GAG GAG GAG ATT ACA ATG CAC	135
31	F A F N I G P S E E E I T M H	45
136	ATC AAT CCT CGT TTT AAT GCC CAT GGA GAC GAG AAT ACG ATT GTG	180
46	I N P R F N A H G D E N T I V	60
181	TGC AAC TCC TAC CAG GGA GGC AGC TGG TGT GAG GAG CAG CGG GAG	225
61	C N S Y Q G G S W C E E Q R E	75
226	TGC TGC TTC CCT TTC CAG CAG GGG GAG GAG TTC AAG ATT GTC ATC	270
76	C C F P F Q Q G E E F K I V I	90
271	GAA TTC ACC CCG TCA GAG TTC GTG GTG ACT TTA TCG GAC GGC TCT	315
91	E F T P S E F V V T L S D G S	105
316	GCC ATC CAC TTC CCC AAC CGC ATG GGC GCT GAG AAA TAC CAG TTC	360
106	A I H F P N R M G A E K Y Q F	120
361	ATC AGC TTC GAC GGG GAC GCT CGC ATC AGA AGC TTT GAG ATC AAG TAA	
121	I S F D G D A R I R S F E I K	*

Fig. 1. The nucleotide and deduced amino acid sequences of LhGal-1 from redlip mullet. The start codon (ATG) and stop codon (TAA) are indicated by bold letters. The GLECT carbohydrate recognition domain is indicated as a grey box. The residues of sugar-binding pocket and putative alternate dimerization interface are marked by red and green letters, respectively.




Fig. 2. The predicted 3D structure of redlip mullet LhGal-1.^{A-E} The yellow letters indicate the carbohydrate-binding sites and the main residues are indicated in pink color.




Fig. 3. Multiple sequence alignment of LhGal-1 with other known orthologs. The residues of sugar-binding pocket and putative alternate dimerization interface are indicated by ★ and ● symbols, respectively.

mammals, birds, reptiles, and amphibians.

To comprehend the evolutionary relationship of redlip mullet LhGal-1, phylogenetic analysis was performed using the amino acid sequences for galectin-1 orthologs available in NCBI database (Fig. 4). The constructed phylogenetic tree was based on the neighbor-joining method with 5,000 bootstrap replicates. LhGal-1 clustered together with the fish species (Fig. 4). The output data of pairwise comparison and phylogenetic analysis suggested that LhGal-1 was closely related to galectin-1 from rock bream (*Oplegnathus fasciatus*) (Table 2).

Tissue-specific distribution of LhGal-1

The mRNA analysis revealed that *LhGal-1* was expressed in all the tested tissues. The stomach (333.20 folds) showed the highest mRNA expression level, followed by the heart (171.09 folds), muscle (152.25 folds), and brain (111.58 folds) (Fig. 5). The fundic, mucin, and epithelial cell surface glycoalyces are recognized by galectin-1 expressed in the gastrointestinal tract, which might be the reason for the highest expression of LhGal-1 in the stomach (Wasano & Hirakawa, 1997). Galectins are widely abundant in muscle tissues, some neurons, the thymus, and epithelial tissues. As per previous studies, *Ctenopharyngodon idella* and *Epinephelus coioides* galectin-1 are highly expressed in the muscle




Fig. 4. Phylogenetic tree of LhGal-1 was constructed using MEGA version X with 5,000 bootstrap repeats and neighbor-joining method.

Table 2. LhGal-1 identity and similarity with homologs from several other species

No	Species	Taxon	GenBank accession No	Amino acids	Identity (%)	Similarity (%)
1	<i>Dicentrarchus labrax</i>	Bony fishes	ACF77003.1	135 aa	83.7	89.6
2	<i>Oplegnathus fasciatus</i>	Bony fishes	ADV35589.1	135 aa	81.5	89.6
3	<i>Amphiprion ocellaris</i>	Bony fishes	XP_023123473.1	135 aa	81.5	88.9
4	<i>Xiphophorus maculatus</i>	Bony fishes	XP_005806370.1	135 aa	77.8	91.1
5	<i>Fundulus heteroclitus</i>	Bony fishes	XP_012737057.1	135 aa	76.3	90.4
6	<i>Danio rerio</i>	Bony fishes	AAR84190.1	134 aa	63	80
7	<i>Homo sapiens</i>	Primates	NP_002296.1	135 aa	41.6	57.7
8	<i>Mus musculus</i>	Rodents	NP_032521.1	135 aa	38.7	56.9
9	<i>Manacus vitellinus</i>	Birds	XP_008929802.2	134 aa	36.3	51.9
10	<i>Chelonia mydas</i>	Turtles	XP_027680905.1	135 aa	37.8	55.6
11	<i>Xenopus laevis</i>	Frogs	AAK11514.1	134 aa	38.6	50




Fig. 5. Relative expression levels of LhGal-1 in different tissues. The fold change of target gene in each tissues normalized to *LhEF1a* and relative to the expression at the blood that has the lowest average Ct value, was calculated for each sample using the formula provided in Livak ($2^{-\Delta\Delta CT}$) method. BL, blood; LV, liver; KD, kidney; SK, skin; GL, gill; HK, head kidney; SP, spleen; IT, intestine; BR, brain; MS, muscle; HT, heart; ST, stomach.

and heart (Chen et al., 2016; Zhu et al., 2019). Galectin-1 is involved in several biological processes as it interacts with multifarious receptors expressed in different cell types (Elola et al., 2005). Besides, galectin-1 binds to polylactosamine chains of laminin and promotes cell detachment and attachment in the extracellular matrix, which influences muscle development (Chen et al., 2016).

LhGal-1 expression analysis after immune challenge

Time-dependent transcriptional regulation of *LhGal-1* expression

was examined in the gill and blood to estimate its dynamic expression profile in response to different immunogenic stimulants (Fig. 6). Fish gills are the main sites of pathogen entry and are a part of the first line of defense in aquatic organisms living in a pathogen-rich environment. Furthermore, blood is the pipeline for the migration of immune cells to the inflammation site. As immunomodulators of innate immunity, galectins are expressed in almost all immune cells and are involved in controlling related biological processes (Cerliani et al., 2011). Accordingly, gill and blood were selected to understand their complex functions following exposure to immune stimulants (LPS, typical cell wall components of gram-negative bacteria; poly I:C, structural analog of viral double-stranded RNA, *L. garvieae*, gram-positive bacteria). Following the injection of LPS and *L. garvieae*, *LhGal-1* mRNA expression was significantly upregulated after 48 h in the gill (Fig. 6A). Consistently, the mRNA expression of galectin-1 in rock bream gill was reported to be significantly upregulated 48 h post injection of *E. tarda* (G-) and *S. iniae* (G+) (Thulasitha et al., 2016). The poly I:C-treated group showed significant upregulation between 48 and 72 h, but not between 6 and 24 h, in the gill (Fig. 6A). Similar early phase downregulation was reported in olive flounder (*Paralichthys olivaceus*) after injection of poly I:C (Liu et al., 2013). Moreover, expression of quadruple domain-containing galectin was downregulated in disk abalone until 12 h after injection of poly I:C and viral hemorrhagic septicemia virus (Gayashani Sandamalika & Lee, 2020). However, the reasons for early downregulation and possible underlying mechanisms remain unclear. It is possible that early phase downregulation of *LhGal-1* in poly I:C-treated group helps in curbing the increase in the viral infection.




Fig. 6. The mRNA expression of (A) *LhGal-1* in the gills post-challenge with LPS, poly I:C, and *Lactococcus garvieae*. The mRNA expression of (B) *LhGal-1* in blood cells after challenge with LPS, poly I:C, and *L. garvieae*. The fold change of *LhGal-1* normalized to *LhEF1a* and relative to the expression in phosphate-buffered saline-injected mullet, was calculated for each sample using the formula provided in Livak ($2^{-\Delta\Delta CT}$) method. Error bar represents the SD ($n = 3$). LPS, lipopolysaccharide; poly I:C, polyinosinic:polycytidylic acid.

On the contrary, the relative *LhGal-1* mRNA expression in blood was significantly upregulated 6 h post injection of LPS, poly I:C, and *L. garvieae*, and 24 h post injection of LPS and poly I:C (Fig. 6B). Consistently, early phase upregulation of galectin-1 was observed in mollusk (*Solen grandis*) hemocytes upon PAMP stimulation (LPS, peptidoglycans [PGN], and glucan) (Wei et al., 2012). Galectins expressed by immune cells not only recognize some pathogens as PRRs (PAMPs) but also contribute to innate immune cell activation and restore homeostasis against inflammation (damage-associated molecular patterns) (Sato et al., 2009). Taken together, these findings along with the previous studies suggested that in redlip mullet, *LhGal-1* might act as a regulator of immune cells and modulate the activity of the encountered pathogen.

Protein purification

The coding sequence of *LhGal-1* was cloned into a pMAL-c5X expression vector, and these plasmids were introduced into the *E. coli* strain, ER2523. The recombinant MBP (rMBP) fusion protein was obtained from IPTG-induced cells and purified using amylose affinity chromatography. SDS-PAGE was performed to visualize rLhGal-1 MBP fusion proteins and samples were obtained from each purification step; uninduced lysates, induced lysates, and eluted proteins. SDS-PAGE indicated no protein expression in the pellet and supernatant of the uninduced lysates,

while specific bands were observed for the rLhGal-1-MBP fusion protein, at approximately 58 kDa (predicted molecular weights = rLhGal-1:15.31 kDa and MBP 42.5 kDa) (Fig. 7).

Sugar-binding ability of rLhGal-1

To investigate the carbohydrate specificity and binding ability of rLhGal-1, a sugar-binding assay was performed with three different carbohydrates as follows: α -lactose, D-galactose, and D-glucose. Although all carbohydrates bound to rLhGal-1, the disaccharide α -lactose showed a higher specificity than the monosaccharides galactose and glucose (Fig. 8). MBP-treated groups showed no significant activity with carbohydrates, which indicated that MBP might not have affected the binding properties of rLhGal-1 (Fig. 8). Significant differences noted in the binding ability of rLhGal-1 to lactose and galactose were possibly due to their structural features. *LhGal-1* CRD forms a beta-sandwich with two skewed sheets and shapes the carbohydrate-binding site. Six strands in the beta-sheet form a groove that is long enough to capture linearized polysaccharides (Leffler et al., 2002). Although both galactose and lactose commonly possess beta-galactoside, rLhGal-1 has a higher affinity for lactose, which is longer than galactose. These results suggested that the binding ability of galectins is dependent on their affinity to the source as well as the length of their ligands.




Fig. 7. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis analysis of maltose-binding protein fusion proteins rLhGal-1 (57.81 kDa). Lane 1: protein marker, Lane 2: lysate of uninduced *Escherichia coli* ER2523 (pellet), Lane 3: lysate of uninduced *E. coli* ER2523 cells (supernatant), Lane 4: lysate of IPTG-induced *E. coli* ER2523 cells (pellet), Lane 5: lysate of IPTG-induced *E. coli* ER2523 cells (supernatant), Lane 6: purified recombinant protein after elution.




Fig. 8. Sugar-binding activity of rLhGal-1 for α -lactose, D-galactose and D-glucose. rMBP, recombinant maltose-binding protein.

Microbial binding activity of rLhGal-1

The microbial binding ability of rLhGal-1 was evaluated using ELISA and revealed that rLhGal-1 could bind to all the tested G (+) and G (-) bacteria, whereas no binding activity was detected

with the elution buffer and MBP controls (Fig. 9). A previous study on the binding activity of galectin from *Eriocheir sinensis* suggested that the interactions between rEsGal and PAMPs (LPS, PGN, and glycan) occur in a dose-dependent manner. Also,




Fig. 9. The microbial-binding activity of rLhGal-1 against several gram-negative and gram-positive bacteria: *Escherichia coli*, *Edwardsiella tarda*, *Vibrio anguillarum*, *Vibrio harveyi*, *Streptococcus iniae*, *Streptococcus parauberis* and *Lactococcus garvieae*. rMBP, recombinant maltose-binding protein; OD, optical density.

the highest binding activity was observed from 200 nmol/L of galectin toward LPS from *E. coli* (Wang et al., 2016). Among the PAMPs, LPS and PGN are major components of G (+) and G (-) bacterial cell walls, respectively. Thus, these results suggested that rLhGal-1 can recognize various pathogens and might act as a PRR involved in the innate immune response of the redlip mullet.

Microbial agglutination activity of rLhGal-1

The microbial agglutination activity of rLhGal-1 was examined using *E. coli*. rLhGal-1 aggregated the tested bacteria while no agglutination was observed with rMBP (Fig. 10). The bacterial agglutination activity of rLhGal-1 further affirmed that *LhGal-1* could be a PRR involved in the immune defense against bacteria. It might also serve as a facilitator of opsonization and phagocytosis by macrophages as shown for galectins in other systems (Cerliani et al., 2011).

Conclusion

We identified galectin-1 homolog from redlip mullet, named LhGal-1, and characterized its function. The *in silico* analysis was




Fig. 10. Agglutination of *Escherichia coli* by rLhGal-1. (A) The cells at 5×10^6 CFU/mL were treated with 20 μ g of rLhGal-1. (B) Recombinant maltose-binding protein as control. The results were observed at 20 \times magnification under light microscope.

performed using several bioinformatics tools. To comprehend the potential role of *LhGal-1* in the host immune system, the spatial and temporal mRNA expressions of *LhGal-1* were estimated via qPCR. As a result, *LhGal-1* was expressed in all tested tissues and upregulation was shown in gill and blood upon being challenged with immune stimulants. The carbohydrate-binding assay revealed that rLhGal-1 has a higher affinity to α -lactose than D-galactose and D-glucose. In addition, rLhGal-1 could bind to all tested bacteria and aggregate *E. coli*. Therefore, this study indicates that *LhGal-1* may be an essential gene for recognizing invading pathogens and modulating host defense mechanisms.

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was supported by the 2022 education, research and student guidance grant funded by Jeju National University.

Acknowledgements

Not applicable.

Availability of data and materials

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Ethics approval and consent to participate

All animal experiments in this study were approved by Institutional Animal Care and Use Committee Jeju National University (approval no. 2018-0003).

ORCID

Chaehyeon Lim <https://orcid.org/0000-0002-9977-561X>
 Hyukjae Kwon <https://orcid.org/0000-0003-4025-4622>
 Jehhee Lee <https://orcid.org/0000-0001-9144-3648>

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol.* 1990;215:403-10.
 Boulianne RP, Liu Y, Aebi M, Lu BC, Kües U. Fruiting body development in *Coprinus cinereus*: regulated expression of two galectins secreted by a non-classical pathway. *Microbiology.* 2000;146:1841-53.

Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976;72:248-54.
 Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, et al. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem.* 2009;55:611-22.
 Cerliani JP, Stowell SR, Mascanfroni ID, Arthur CM, Cummings RD, Rabinovich GA. Expanding the universe of cytokines and pattern recognition receptors: galectins and glycans in innate immunity. *J Clin Immunol.* 2011;31:10-21.
 Chen HY, Weng IC, Hong MH, Liu FT. Galectins as bacterial sensors in the host innate response. *Curr Opin Microbiol.* 2014;17:75-81.
 Chen X, Wei J, Xu M, Yang M, Li P, Wei S, et al. Molecular cloning and characterization of a galectin-1 homolog in orange-spotted grouper, *Epinephelus coioides*. *Fish Shellfish Immunol.* 2016;54:333-41.
 Elola MT, Chiesa ME, Alberti AF, Mordoh J, Fink NE. Galectin-1 receptors in different cell types. *J Biomed Sci.* 2005;12:13-29.
 Gayashani Sandamalika WM, Lee J. Quadruple domain-containing galectin from marine invertebrate disk abalone (*Haliotis discus discus*): molecular perspectives in early development, immune expression, and potent antiviral responses. *Fish Shellfish Immunol.* 2020;106:920-9.
 Han HJ, Lee NS, Kim MS, Jung SH. An outbreak of *Lactococcus garvieae* infection in cage-cultured red lip mullet *Chelon haematocheilus* with green liver syndrome. *Fish Aquat Sci.* 2015;18:333-9.
 Hirabayashi J, Kasai K. The family of metazoan metal-independent β -galactoside-binding lectins: structure, function and molecular evolution. *Glycobiology.* 1993;3:297-304.
 Kumagai Y, Takeuchi O, Akira S. Pathogen recognition by innate receptors. *J Infect Chemother.* 2008;14:86-92.
 Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol.* 2018;35:1547-9.
 Leffler H, Carlsson S, Hedlund M, Qian Y, Poirier F. Introduction to galectins. *Glycoconj J.* 2002;19:433-40.
 Letunic I, Bork P. 20 years of the SMART protein domain annotation resource. *Nucleic Acids Res.* 2018;46:D493-6.
 Liu S, Hu G, Sun C, Zhang S. Anti-viral activity of galectin-1 from flounder *Paralichthys olivaceus*. *Fish Shellfish Immunol.* 2013;34:1463-9.
 Livak KJ, Schmittgen TD. Analysis of relative gene expression

- data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods*. 2001;25:402-8.
- Liyana DS, Omeka WKM, Godahewa GI, Lee S, Nam BH, Lee J. Membrane attack complex-associated molecules from redlip mullet (*Liza haematocheila*): molecular characterization and transcriptional evidence of C6, C7, C8 β , and C9 in innate immunity. *Fish Shellfish Immunol*. 2018;81:1-9.
- Ministry of Maritime Affairs and Fisheries. 2020 Ministry of Oceans and Fisheries: statistical year book of maritime affairs and fisheries. Sejong: Ministry of Maritime Affairs and Fisheries; 2020.
- Minos G, Imsiridou A, Economidis PS. *Liza haematocheilus* (Pisces, Mugilidae) in the northern Aegean Sea. In: Golani D, Appelbaum-Golani B, editors. Fish invasions mediterranean sea: change renewal. Sofia: Pensoft; 2009. p.313-32.
- Roy A, Kucukural A, Zhang Y. I-TASSER: a unified platform for automated protein structure and function prediction. *Nat Protoc*. 2010;5:725-38.
- Sato S, St-Pierre C, Bhaumik P, Nieminen J. Galectins in innate immunity: dual functions of host soluble β -galactoside-binding lectins as damage-associated molecular patterns (DAMPs) and as receptors for pathogen-associated molecular patterns (PAMPs). *Immunol Rev*. 2009;230:172-87.
- Shi W, Xue C, Su X, Lu F. The roles of galectins in parasitic infections. *Acta Trop*. 2018;177:97-104.
- Thulasitha WS, Umasuthan N, Whang I, Nam BH, Lee J. Anti-microbial response of galectin-1 from rock bream *Oplegnathus fasciatus*: molecular, transcriptional, and biological characterization. *Fish Shellfish Immunol*. 2016;50:66-78.
- Vasta GR. Galectins as pattern recognition receptors: structure, function, and evolution. In: Lambris JD, Hajishengallis G, editors. Current topics in innate immunity II. New York, NY: Springer; 2012.
- Vasta GR, Nita-Lazar M, Giomarelli B, Ahmed H, Du S, Cammarata M, et al. Structural and functional diversity of the lectin repertoire in teleost fish: relevance to innate and adaptive immunity. *Dev Comp Immunol*. 2011;35:1388-99.
- Wang M, Wang L, Huang M, Yi Q, Guo Y, Gai Y, et al. A galectin from *Eriocheir sinensis* functions as pattern recognition receptor enhancing microbe agglutination and haemocytes encapsulation. *Fish Shellfish Immunol*. 2016;55:10-20.
- Wang WH, Lin CY, Chang MR, Urbina AN, Assavalapsakul W, Thitithayanont A, et al. The role of galectins in virus infection - a systemic literature review. *J Microbiol Immunol Infect*. 2020;53:925-35.
- Wasano K, Hirakawa Y. Recombinant galectin-1 recognizes mucin and epithelial cell surface glycocalyxes of gastrointestinal tract. *J Histochem Cytochem*. 1997;45:275-83.
- Wei X, Yang J, Liu X, Yang D, Xu J, Fang J, et al. Identification and transcriptional analysis of two types of lectins (SgCTL-1 and SgGal-1) from mollusk *Solen grandis*. *Fish Shellfish Immunol*. 2012;33:204-12.
- Zhu D, Fu P, Huang R, Xiong L, Wang Y, He L, et al. Molecular characterization, tissue distribution and functional analysis of galectin 1-like 2 in grass carp (*Ctenopharyngodon idella*). *Fish Shellfish Immunol*. 2019;94:455-63.