

RESEARCH ARTICLE

Mycorrhization of *Quercus acutissima* with *Tuber borchii* and *Tuber melanosporum*

Jung-Min Lee, and Ahn-Heum Eom*


Department of Biology Education, Korea National University of Education, Cheongju 28173, Korea

*Corresponding author: eomah@knue.ac.kr

ABSTRACT

Truffles are ectomycorrhizal fungi that belong to the genus *Tuber*. They exhibit symbiotic relationships, particularly with oak (*Quercus* spp.) and hazel (*Corylus* spp.) trees. We performed an inoculation using a spore suspension to synthesize mycorrhizae between European truffles, *Tuber borchii* and *Tuber melanosporum*, and an indigenous oak species, *Quercus acutissima*. This resulted in the formation of mycorrhizae within 2 months after inoculation. Despite having the same host plant, differences in features were observed between *Tuber* species, including color and mantle type. These results indicate that *Q. acutissima* is a suitable host plant for truffle cultivation in Korea and provide a better understanding of the mycorrhization of *T. borchii* and *T. melanosporum*.

Keywords: Ectomycorrhiza, *Quercus acutissima*, Symbiosis, *Tuber borchii*, *Tuber melanosporum*



OPEN ACCESS

pISSN : 0253-651X
eISSN : 2383-5249

Kor. J. Mycol. 2022 December, 50(4): 275-280
<https://doi.org/10.4489/KJM.20220029>

Received: August 26, 2022

Revised: December 13, 2022

Accepted: December 09, 2022

© 2022 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Truffles are fungi belonging to the genus *Tuber* that produce hypogeous fruiting bodies with a unique taste and aroma. They have ectomycorrhizal (ECM) association with several host plants, especially oak and hazel trees, by colonizing the cortex cells of their roots to form a Hartig net and creating a thick mycelium outside the roots [1]. *Tuber* spp. have high host specificity [2]. For example, *Tuber melanosporum* shows a higher host preference for Northern Hemisphere angiosperms (e.g., *Quercus* spp., *Corylus* spp.) but a lower host preference for *Pinus* spp. In contrast, *Tuber borchii* has a wider host range, including angiosperms (e.g., *Quercus* spp., *Populus* spp.) and gymnosperms (e.g., *Pinus* spp., *Picea* spp., *Cedrus* spp.) [3]. The selection of host plants affects mycorrhiza formation [4], and the rate of mycorrhizal root colonization varies depending on the host plant [5]. The *Tuber* ECM, such as the ECM system or parenchymatous mantle structure, has general characteristics [6]. However, there are distinctive features of the ECM according to the host plant and *Tuber* spp. For example, the *Tuber indicum* ECM shows morphological differences depending on the host plant [7,8]. In contrast, the mycorrhizal characteristics of Chinese white truffles on the same host plant can be different [9].

In this study, we inoculated a spore suspension of European truffles, *T. borchii* and *T. melanosporum*, onto an indigenous host plant, *Quercus acutissima*. The Périgord black truffle (*T. melanosporum*) and *T. borchii* are European truffles of high economic and commercial value. Furthermore, these are the most successfully cultivated

species in many country [10-12]. However, there has been no record of successful mycorrhization between those truffles with host plants in Korea, although *T. himalayense* and *T. huidongense* collected in Korea were found to produce mycorrhiza with two oak trees in Korea [5]. Moreover, *Q. acutissima* is a widely distributed oak species in Korea. Therefore, the objectives of this study were (1) to determine whether European truffle species can form ECM with an indigenous host plant and (2) to identify the morphological and anatomical characteristics of truffle mycorrhizae.

MATERIALS AND METHODS

Preparation of seedlings

Q. acutissima seeds were collected from a natural site in Cheongju, Korea. The shells of these seeds were then removed. The seeds were washed with tap water and surface sterilized with 10% sodium hypochlorite (NaOCl) for 30 min. For germination, seeds were placed in plastic pots (280 mL) with a mixture of sterile vermiculite and perlite (1:1 ratio) and maintained in a growth room for 6 months (8 h photoperiod per day, $55 \pm 5\%$ relative humidity, and $24 \pm 1^\circ\text{C}$ temperature).

Inoculation onto seedlings and mycorrhizal synthesis

Ascocarps of *T. borchii* and *T. melanosporum* were purchased from Chiko Corporation, Korea. The ascocarps were rinsed with tap water, sterilized with 70% ethanol, and then ground in sterile water. Six-month-seedlings of *Q. acutissima* were inoculated with 1 mL of a spore suspension containing 1.4×10^6 spores/mL, which were counted using a hemocytometer (Paul Marienfeld GmbH & CO., KG, Lauda-Königshofen, Germany). The inoculated seedlings were placed in plastic pots (280 mL) with a mixture of sterile vermiculite and perlite (1:1 ratio). They were cultured in a growth room for 6 months and watered weekly (8 h photoperiod per day, $55 \pm 5\%$ relative humidity, and $24 \pm 1^\circ\text{C}$ temperature).

Molecular identification of ECM

The mycorrhizal root tips of inoculated seedlings were sampled to confirm mycorrhiza formation after 2 months of inoculation. Genomic DNA was extracted using the HiGene™ Genomic DNA Prep Kit (BIOFACT, Daejeon, Korea). The rDNA region, including the internal transcribed spacer (ITS) region, was amplified by PCR using ITS1F and ITS4 primers [13]. Nucleotide sequences were analyzed at Solgent (SolGent Co., Ltd., Daejeon, Korea). Sequence similarity was determined using BLAST (<https://www.ncbi.nlm.nih.gov/>).

Six months after inoculation, the morphological characteristics of *T. borchii* and *T. melanosporum* ECM were observed. First, the ECM of inoculated seedlings was sampled. Then, the morphological characteristics of the ECM were observed using a dissection microscope (Olympus SZX7, Tokyo, Japan). Cross-and longitudinal sections were obtained using a cryostat (CM1850, Leica, Wetzlar, Germany). Subsequently, anatomical characteristics were observed via a light microscope (Axio Imager A1, Carl ZEISS, Oberkochen, Germany).

RESULTS

Mycorrhization and molecular identification of ECM

Mycorrhization of *T. borchii* and *T. melanosporum* on *Q. acutissima* was confirmed 2 months after inoculation. The ITS sequences amplified from the mycorrhizal root tips of *T. borchii* (NCBI accession No. OP975718) and *T. melanosporum* (NCBI accession No. OP978675) showed more than 99% similarity with those of sporocarps inoculated to seedlings using BLAST.

Morphological and anatomical characterization of ECM

Six months after inoculation, 20 seedlings that formed mycorrhiza were selected per *Tuber* species. Morphological and anatomical features of the ECM of *T. borchii* and *T. melanosporum* were observed, measured, and analyzed (Table 1). For the *T. borchii* ECM, the ectomycorrhizal system was simple, with a monopodial pinnate pattern. The unramified ends were straight, cylindrical, or club-shaped with rounded ends. Their texture was short-spiny, and their color was whitish light brown to brown and rarely dark brown (Fig. 1). The mantle type had a non-interlocking pattern, with an angular pattern arranged in 2-5 hyphal layers. The outer mantle was yellowish and reddish brown. The emanating hyphae were straight, simple, bristle-like, awl-shaped, and septate.

Table 1. Dimensional measurements of *Tuber borchii* and *Tuber melanosporum* mycorrhizae (mean±standard error).

Dimensions	<i>T. borchii</i>	<i>T. melanosporum</i>
Ectomycorrhizal system (mm)	1.97±0.14	1.7±0.14
Length of unramified ends (µm)	716±38.7	415±38.2
Diameter of unramified ends (µm)	220±7.77	185±9.87
Thick of mantle (µm)	15.8±0.97	18.5±0.65
Length of cystidia (µm)	87.4±5.10	101.9±5.15
Diameter of cystidia (µm)	2.61±0.21	2.57±0.16




Fig. 1. Macro-morphological and anatomical characteristics of ectomycorrhizal roots of *Quercus acutissima* inoculated with *Tuber borchii*. Mycorrhizal root tips (A, B); cross-section of mycorrhizal root tips (C, D); outer mantle surface structure (E); and separate hyphae emanating from outer mantle layer (F, G) (scale bars: A, B=500 µm; C-E=20 µm; F, G=50 µm).

ECM of *T. melanosporum* was simple and rarely followed a monopodial pinnate pattern (Fig. 2). The shape of the unramified ends was straight, bent, cylindrical, or mostly club-shaped with rounded ends and often long-spiny. Their colors were ochre, brown, dark brown, and mostly black. The mantle type showed an irregular interlocking pattern with 2-5 angularly arranged hyphal layers. The outer mantle was reddish-brown and brown. The emanating hyphae were bent, simple, bristle-like, awl-shaped, and septate and had right-angle ramifications.




Fig. 2. Macro-morphological and anatomical characteristics of ectomycorrhizal roots of *Quercus acutissima* inoculated with *Tuber melanosporum*. Mycorrhizal root tips (A, B); cross-section of mycorrhizal root tips (C, D); outer mantle surface structure (E); and separate hyphae emanating from outer mantle layer (F, G) (scale bars: A, B=500 μ m; C-E=20 μ m; F, G=50 μ m).

DISCUSSION

The time required for mycorrhiza formation differs depending on the host species [5]. *T. borchii* and *T. melanosporum* ECM on *Q. acutissima* exhibited mycorrhization 2 months after inoculation. In previous studies, the mycorrhization of *T. borchii* with *Arbutus unedo* was found to occur 2 months after inoculation [14] and that of *T. melanosporum* with *Quercus* spp. was observed 6 months after inoculation [10]. In *Tuber himalayense* and *T. huidongense*, mycorrhization with an indigenous *Quercus* spp. requires 2.5 months, and it was confirmed that mycorrhization occurs more quickly compared to that with other host plants [5]. These results indicated that *Tuber* spp. have a higher preference for native oak trees in Korea. Moreover, *Q. acutissima* was found to be a suitable host plant for *Tuber* spp.

Compared to those reported in previous studies, most ECM features of *T. borchii* and *T. melanosporum* have been described similarly. However, distinct features were also observed. The *T. melanosporum* ECM with *Quercus* spp., *A. unedo*, and *Carya illinoensis* is yellow-ocher or brown [10,15,16], but that with *Q. acutissima* is mainly dark. In terms of the mantle structure, the *T. borchii* ECM with *A. unedo*, *Tilia platyphyllos*, *C. illinoensis*, and *Quercus robur* has an irregular interlocking pattern [14,17-19], whereas that with *Q. acutissima* shows a non-interlocking irregular pattern.

Despite the host plant being the same, the characteristics of mycorrhizae vary depending on the *Tuber* species. In this study, the ECM and mantle color of *T. borchii* were brighter than those of *T. melanosporum*. In the mantle type, *T. borchii* showed a non-interlocking irregular pattern, whereas *T. melanosporum* showed an irregular interlocking pattern. Whereas *T. borchii* had straight cystidia, *T. melanosporum* had bent cystidia. The mycorrhizal characteristics of the two *Tuber* spp. were thus clearly different. It is reasonable to combine morphological and anatomical characteristics to distinguish the ECM of different *Tuber* species [6]. However, these characteristics alone are not sufficient to determine which *Tuber* species form mycorrhizae. This is because phylogenetically similar *Tuber* species also have similar mycorrhizal characteristics; therefore, molecular analysis is needed for accurate identification [8].

We successfully induced mycorrhization between a European truffle species and an indigenous oak species in Korea. The inoculation method using a spore suspension could further be applied to the mycorrhization of European truffle species on indigenous oak species in Korea. This suggests that indigenous oak can be an appropriate host plant for commercial truffle cultivation. In addition, mycorrhiza formation by *Tuber* spp. with other indigenous oak species needs to be confirmed. This is the first report of the mycorrhization of *T. borchii* and *T. melanosporum* with *Q. acutissima*. Thus, the range of host plants for *T. borchii* and *T. melanosporum* has expanded. This study provides the morphological and anatomical characteristics of the ECM depending on the *Tuber* spp. However, further studies are needed to determine whether the mycorrhizae of *Tuber* spp. are maintained in a natural environment after transplant.

CONFLICT OF INTERESTS

No conflict of interest was reported by the author(s).

ACKNOWLEDGMENT

This work was supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (IPET) funded by the Ministry of Agriculture, Food and Rural Affairs (MAFRA) (IPET319106052HD050).

REFERENCES

1. Fischer C, Oliach D, Bonet Lledos JA, Colinas C. Best practices for cultivation of truffles. Solsona: Centre Tecnològic Forestal de Catalunya; 2017.
2. Molina R, Trappe JM. Patterns of ectomycorrhizal host specificity and potential among Pacific Northwest conifers and fungi. *For Sci* 1982;28:423-58.
3. Benucci GMN, Bonito G, Falini LB, Bencivenga M, Donnini D. Truffles, timber, food, and fuel: sustainable approaches for multi-cropping truffles and economically important plants. Berlin: Springer; 2012. p. 265-80.

4. Bonito G, Trappe JM, Rawlinson P, Vilgalys R. Improved resolution of major clades within *Tuber* and taxonomy of species within the *Tuber gibbosum* complex. *Mycologia* 2010;102:1042-57.
5. Gwon JH, Park H, Eom AH. Mycorrhization of *Quercus* spp. with *Tuber huidongense* and *T. himalayense* Collected in Korea. *Mycobiology* 2022;50:104-9.
6. Cairney JW, Chambers SM. Ectomycorrhizal fungi: key genera in profile. Berlin: Springer Science & Business Media; 2013.
7. Geng LY, Wang XH, Yu FQ, Deng XJ, Tian XF, Shi XF, Xie XD, Liu PG, Shen YY. Mycorrhizal synthesis of *Tuber indicum* with two indigenous hosts, *Castanea mollissima* and *Pinus armandii*. *Mycorrhiza* 2009;19:461-7.
8. Huang LL, Guerin-Laguette A, Wang R, Yu FQ. Characterization of *Tuber indicum* (Pezizales, Tuberales) mycorrhizae synthesized with four host trees exotic to China. *Symbiosis* 2020;82:215-24.
9. Wan SP, Yu FQ, Tang L, Wang R, Wang Y, Liu PG, Wang XH, Zheng Y. Ectomycorrhizae of *Tuber huidongense* and *T. liyuanum* with *Castanea mollissima* and *Pinus armandii*. *Mycorrhiza* 2016;26:249-56.
10. Wang R, Guerin-Laguette A, Butler R, Huang LL, Yu FQ. The European delicacy *Tuber melanosporum* forms mycorrhizae with some indigenous Chinese *Quercus* species and promotes growth of the oak seedlings. *Mycorrhiza* 2019;29:649-61.
11. Zambonelli A, Iotti M, Puliga F, Hall IR. Enhancing white truffle (*Tuber magnatum* Picco and *T. borchii* Vittad.) cultivation through biotechnology innovation. Berlin: Springer; 2021. p. 505-32.
12. Donnini D, Gargano M, Perini C, Savino E, Murat C, Di Piazza S, Altobelli E, Salerni E, Rubini A, Rana G. Wild and cultivated mushrooms as a model of sustainable development. *Plant Biosys* 2013;147:226-36.
13. White TJ, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. New York: Academic Press, Inc.; 1990.
14. Lancellotti E, Iotti M, Zambonelli A, Franceschini A. Characterization of *Tuber borchii* and *Arbutus unedo* mycorrhizas. *Mycorrhiza* 2014;24:481-6.
15. Marozzi G, Sánchez S, Benucci GMN, Bonito G, Falini LB, Albertini E, Donnini D. Mycorrhization of pecan (*Carya illinoensis*) with black truffles: *Tuber melanosporum* and *Tuber brumale*. *Mycorrhiza* 2017;27:303-9.
16. Ori F, Leonardi M, Faccio A, Sillo F, Iotti M, Pacioni G, Balestrini R. Synthesis and ultrastructural observation of arbutoid mycorrhizae of black truffles (*Tuber melanosporum* and *T. aestivum*). *Mycorrhiza* 2020;30:715-23.
17. Giomaro G, Zambonelli A, Sisti D, Cecchini M, Evangelista V, Stocchi V. Anatomical and morphological characterization of mycorrhizas of five strains of *Tuber borchii* Vittad. *Mycorrhiza* 2000;10:107-14.
18. Benucci GMN, Bonito G, Falini LB, Bencivenga M. Mycorrhization of pecan trees (*Carya illinoensis*) with commercial truffle species: *Tuber aestivum* Vittad. and *Tuber borchii* Vittad. *Mycorrhiza* 2012;22:383-92.
19. Leonardi P, Iotti M, Zeppa SD, Lancellotti E, Amicucci A, Zambonelli A. Morphological and functional changes in mycelium and mycorrhizas of *Tuber borchii* due to heat stress. *Fungal Ecology* 2017;29:20-9.