



ARTICLE

## Preparation and Antioxidant Activities of High Fischer's Ratio Oligopeptides from Goat Whey

Yusi Qin<sup>1</sup>, Ming Cheng<sup>2</sup>, Xiaoxue Fan<sup>1</sup>, Xiaoqing Shao<sup>1</sup>, Cunfang Wang<sup>1,\*</sup>, Hua Jiang<sup>1,\*</sup>, and Xiaoning Zhang<sup>1</sup>

<sup>1</sup>College of Food Science and Engineering, Qilu University of Technology (Shandong Academy of Sciences), Jinan 250353, China

<sup>2</sup>Qingdao Research Institute of Husbandry and Veterinary, Qingdao 266100, China



Received April 14, 2022  
Revised June 11, 2022  
Accepted June 22, 2022

\*Corresponding author :

Cunfang Wang  
College of Food Science and Engineering,  
Qilu University of Technology (Shandong  
Academy of Sciences), Jinan 250353, China  
Tel: +86-183-5310-8782  
E-mail: [cunfangwang@163.com](mailto:cunfangwang@163.com)

Hua Jiang  
College of Food Science and Engineering,  
Qilu University of Technology (Shandong  
Academy of Sciences), Jinan 250353, China  
Tel: +86-135-8910-1572  
E-mail: [jhmengyou@163.com](mailto:jhmengyou@163.com)

\*ORCID

Yusi Qin  
<https://orcid.org/0000-0002-8468-2824>  
Ming Cheng  
<https://orcid.org/0000-0003-0183-1597>  
Xiaoxue Fan  
<https://orcid.org/0000-0002-4048-7875>  
Xiaoqing Shao  
<https://orcid.org/0000-0001-7736-0707>  
Cunfang Wang  
<https://orcid.org/0000-0001-7992-6524>  
Hua Jiang  
<https://orcid.org/0000-0003-0013-7173>  
Xiaoning Zhang  
<https://orcid.org/0000-0002-5914-5437>

**Abstract** This study aimed to obtain high Fischer's ratio oligopeptides from goat whey (HFO) and investigate antioxidant property of it. Hydrolysis of goat whey was done with the approach of sequential digestion of pepsin and flavourzyme. With the adsorption of aromatic amino acids by activated carbon, HFO with a Fischer's ratio of 27.070 and a molecular weight of 200–1,000 Da were obtained, and the branched-chain amino acids accounted for 22.87%. Then the antioxidant activity of HFO was evaluated. At the concentrations of 2.0 mg/mL and 0.50 mg/mL, HFO scavenged 77.27% and 99.63% of 1,1-diphenyl-2-picrylhydrazyl and 3-ethylbenzthiazoline-6-sulphonate free radicals respectively. The scavenging rate of HFO against hydroxyl radicals reached 92.31% at the concentration of 0.25 mg/mL. Animal experiments demonstrated that HFO could moderate the changes of malondialdehyde, superoxide dismutase and glutathione peroxidase caused by CCl<sub>4</sub>-induced oxidative stress *in vivo*. This study indicated that HFO from goat whey was capable of oxidation resistance both *in vivo* and *in vitro*, which provided a scientific basis for the high-value processing and application of goat milk whey.

**Keywords** goat whey, enzymatic hydrolysis, high Fischer's ratio oligopeptides, antioxidant, free radical

### Introduction

Whey is a by-product of the cheese making process, containing a lot of whey proteins such as  $\alpha$ -lactalbumin,  $\beta$ -lactoglobulin, immunoglobulin and serum albumin (Chen et al., 2019; Xia et al., 2021). Whey also has the characteristics of high protein efficiency ratio, high biological value and high utilization rate (Da Silva et al., 2017). Whey protein contains higher content of branched-chain amino acids (BCAA) than other protein sources, which is crucial for tissue growth and repair (Kerasiotti et al., 2019). Goat whey has remarkable antioxidant property at both the molecular and tissue levels because of its ability to neutralize numerous free radicals (Kerasiotti et al., 2014). Compared with the milk from cow, buffalo, sheep and camel, the whey protein from

goat milk contains the highest levels of proline ( $53 \pm 1.30$  mg/g) and eight essential amino acids (Rafiq et al., 2015). However, 47% of the whey is directly discharged around the world, and approximately 1.3 million tons of goat whey is discarded or used as animal feeds every year, causing serious environmental problems (Garay et al., 2021; Papademas and Kotsaki, 2019). Therefore, the high-value reuse of whey has become an urgent problem to be solved. The value of whey can be enhanced by hydrolysis, a process that can mitigate the sensitivity of whey protein by reducing the size of peptides (Bordenave et al., 2000). In addition, BCAA in whey protein hydrolytes have been shown to be able to protect skin cells from inflammation, and low molecular weight peptides usually show strong antioxidant activity after protein hydrolysis (Da Silva et al., 2017; Ranathunga et al., 2006). Goat whey peptides possess biological properties such as immunomodulation, antimicrobial and antihypertension (Osman et al., 2016). Based on the above, high Fischer's ratio (F ratio) oligopeptides can be prepared from goat whey to increase the value of goat whey. High F ratio oligopeptides refers to the active peptides with the BCAA to aromatic amino acid (AAA) ratio of over 20, and molecular weight (Mw) of 200–1,000 Da, physiological property of it in relieving phenylketonuria and anti-fatigue have been reported (Li et al., 2019). Up to now, the raw materials for the preparation of high F ratio oligopeptides are mostly plant-derived proteins, such as soy and flaxseed protein (Salma et al., 2022; Yan et al., 2021). For animal-derived proteins, only cow casein was reported in 2004 (Pedroche et al., 2004). As an ideal source both in terms of BCAA content and the activity of goat whey peptides, using whey protein for the preparation of high F ratio oligopeptides has not been reported.

In this study, goat whey was hydrolyzed to obtain high F ratio oligopeptides through two-step hydrolysis of pepsin and flavourzyme. The optimal conditions for the double-enzyme stepwise hydrolysis of whey protein and the conditions for activated carbon to adsorb into AAA were discussed. The Mw distribution of high Fischer's ratio oligopeptides from goat whey (HFO) was defined, and the antioxidant effects of HFO were evaluated by performing various antioxidant assays after the pre-treatment of HFO both *in vivo* and *in vitro*. This study not only expanded the high-value processing and utilization of goat whey, but also was of great significance for the diversification and advanced development as well as application of HFO products.

## Materials and Methods

### Collection and preparation of goat whey samples

Goat milk samples were collected from healthy adult dairy goats raised in AoTe Goat Farm Qingdao Shandong Province, China, and transported to the laboratory by cold chain. After two 10-minute centrifugation processes at  $4^{\circ}\text{C}$ ,  $2,149 \times g$ , the pH value of goat milk was adjusted to 4.1 with 10% acetic acid. Then milk samples were stored at  $4^{\circ}\text{C}$  for 30 min until casein precipitated, and then centrifuged at  $2,149 \times g$  for 10 min at  $4^{\circ}\text{C}$  to obtain the supernatant, namely the goat whey. After that, the goat whey was filtered with a dialysis bag of 3,500 Da to remove the small molecules, such as salt and lactose. Then the milk whey was freeze-dried to obtain goat whey powder for the enzymatic hydrolysis in the next step.

### Optimization of enzymatic hydrolysis conditions and the degree of hydrolysis (DH)

The two-step enzymatic reaction was performed in a beaker containing a mixture of 100 mL of distilled water and 5 g of goat whey powder. Pepsin hydrolyses were carried out under different conditions where five values for each indicator of time, temperature, pH, and pepsin were chosen. Then the resulting hydrolysate was further hydrolyzed with flavourzyme under the same varied conditions as pepsin. The details of the conditions were shown in Table 1. An orthogonal L9 ( $3^4$ ) test design was

**Table 1. Enzymatic hydrolysis conditions**

Optimal conditions	Variable	Quantitative
Time (h)	2–6	Add 1.25% (w/w) pepsin, pH=2, 45°C
Temperature (°C)	40–60	Add 1.25% (w/w) pepsin, pH=2, 2 h
pH	1–3	Add 1.25% (w/w) pepsin, 45°C, 2 h
The amount of pepsin added [% (w/w)]	0.75–1.75	pH=2, 45°C, 2 h
Time (h)	2–6	Add 1% (w/w) flavourzyme, pH=6.5, 45°C
Temperature (°C)	40–60	Add 0.75% (w/w) flavourzyme, pH=7, 3 h
pH	6–8	Add 1% (w/w) flavourzyme, 45°C, 3 h
The amount of flavourzyme added [% (w/w)]	0.25–1.25	pH=7, 3 h, 45°C

used to investigate the optimal hydrolysis condition of each enzyme. Four controllable variables, including time (A), temperature (B), pH (C), and the amount of pepsin added (D) were selected for optimization. And three independent trials were conducted at the optimum of enzymatic treatments. The DH of the hydrolysate was determined with the pH-stat method, which was calculated according to the following Eq. (1) (Mat et al., 2020).

$$DH = 100 \times \frac{V \times N}{M \times h_{tot}} \times \frac{1}{1 - \alpha_{COOH}} \quad (1)$$

where  $V$  is the volume of added HCl (mL),  $N$  is the normality of the acid,  $M$  is the mass of proteins (g),  $h_{tot}$  is the total number of peptide bonds in whey proteins (8.8 meqv/g), and  $\alpha_{COOH}$  is the mean degree of dissociation of the carboxylic groups produced, which was calculated using the following Eq. (2).

$$\alpha_{COOH} = \frac{10^{PH-PK_A(COOH)}}{1 + 10^{PH-PK_A(COOH)}} \quad (2)$$

### Optimization of conditions for aromatic amino acid (AAA) adsorption with activated carbon

The activated carbon (200 mesh, powder) was used to adsorb AAA in the hydrolysate solution of goat milk whey (Cermakova et al., 2017). The optimum adsorption condition was explored by evaluating the adsorption of the hydrolysate under varying time, pH, and carbon to liquid ratio conditions, the details of which are shown in Table 2. The F ratio of the hydrolysate was measured according to the method of Li et al. (2019). An orthogonal L9 (3<sup>3</sup>) test design was used to investigate the optimal adsorption condition of active carbon. Three controllable variables, including time (A), pH (B) and carbon to liquid ratio (C), were selected for optimization. And three independent trials were conducted at the optimum of adsorption. The absorbance of the hydrolysate was measured at 280 nm and 220 nm using the ultraviolet spectrophotometer.

**Table 2. Adsorption conditions**

Optimal conditions	Variable	Quantitative
Time (h)	0.5, 1, 2, 3, 4	pH=3, carbon to liquid ratio=1:5
pH	1, 2, 3, 4, 5	Carbon to liquid ratio=1:10, 3 h
Carbon to liquid ratio	1:3, 1:4, 1:5, 1:6, 1:7	pH=3, 3 h

F ratio was defined according to the following Eq. (3), where OD<sub>280</sub> and OD<sub>220</sub> refer to the absorbance of the hydrolysate at 280 nm and 220 nm, respectively.

$$F \text{ ratio} = \frac{OD_{220}}{OD_{280}} \quad (3)$$

### Determination of amino acids and molecular weight (Mw)

The amount of amino acids was determined with a method modified slightly according to the study of Song et al. (2020). The oligopeptide solution was thoroughly mixed with 6 M HCl, where nitrogen was blown for three times, then the obtained substance was hydrolyzed with acid for 22–24 h at 110°C. After that, the hydrolysate was filtered with a double filter paper and diluted with ultrapure water. The filtrate was then spin-evaporated at 50°C to remove HCl and filtered with a 0.22 µm microfiltration membrane filter. The amino acid content in the oligopeptide was determined with an automatic amino acid analyzer (Hitachi High-Tech Science Corporation, Tokyo, Japan).

The Mw distribution of the oligopeptide was determined by gel permeation chromatography (PL\_GPC50, Polymer Laboratories, Church Stretton, UK) with two PL aqua gel-OH mixed 8 µm (Agilent, Santa Clara, CA, USA) columns at 30°C, and R1 detector (Waters Corporation, Milford, MA, USA). The mobile phase was 0.1 M sodium nitrate and 500 ppm sodium azide aqueous solution. PEG/PEO was prepared in the mobile phase as a reference standard for determining Mw distribution of the oligopeptide. The oligopeptide solutions were filtered with a 0.45 µm filter before sampling and analysis. The relative Mw and Mw distribution were calculated with the software according to the peak time of the oligopeptide solution.

### 1,1-Diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity (RSA)

The DPPH scavenging activity of the HFO was determined according to the method described by Duan et al. (2014). HFO solution was freeze-dried afterwards to obtain HFO powder. 2 mL solutions with different HFO concentrations were added respectively to 2.0 mL ethanol with DPPH (0.1 mM). The mixture was left in dark for 30 min. The absorbances by the HFO solutions were measured using the spectrophotometer (UV756, Shanghai Yoke Instrument, Shanghai, China) at 517 nm. The DPPH RSA was calculated using the following Eq. (4).

$$\text{DPPH RSA (\%)} = (A_0 - A) / A_0 \times 100\% \quad (4)$$

In the formula, A<sub>0</sub> is the DPPH absorbance by distilled water, and A is the DPPH absorbance by the HFO solution. Analyses of all samples were run in triplicate and averaged.

### 3-Ethylbenzthiazoline-6-sulphonate (ABTS) radical scavenging activity (RSA)

The ABTS free radicals assay was conducted according to the method reported by Li et al. (2012). 0.3 mL solutions with different HFO concentrations were added to 1.2 mL of ABTS<sup>•+</sup> solution. After incubation for 30 min, the absorbance levels at 734 nm were recorded using the spectrophotometer (UV756, Shanghai Yoke Instrument). An equivalent volume with ascorbic acid concentration was used as the control. The initial absorbance at 734 nm was controlled to 0.70±0.02. The ABTS radical scavenging effect was calculated using the following Eq. (5).

$$\text{ABTS RSA (\%)} = (A_0 - A) / A_0 \times 100\% \quad (5)$$

In the formula,  $A_0$  is the absorbance at 734 nm with ascorbic acid solution, while  $A$  is the absorbance at 734 nm by the HFO solution. Analyses of all samples were run in triplicate and averaged.

### Hydroxyl radical scavenging activity (RSA)

The hydroxyl RSA was determined according to the method described by Tang and Huang (2018). 1 mL solutions with different HFO concentrations were put into a test tube and then 1 mL  $9 \times 10^{-3}$  M  $\text{FeSO}_4$  solution, 1 mL  $9 \times 10^{-3}$  M salicylic acid ethanol (70%) solution and 1 mL  $9 \times 10^{-3}$  M  $\text{H}_2\text{O}_2$  were added. After reaction in a water bath for 30 minutes at  $37^\circ\text{C}$ , the ultraviolet absorbance of the mixture was measured at 510 nm. Ascorbic acid was used as a control. The scavenging effect of hydroxyl radical was calculated using the following Eq. (6).

$$\text{RSA (\%)} = (A_0 - A) / A_0 \times 100\% \quad (6)$$

In the formula,  $A_0$  is the absorbance of the mixture by the ascorbic acid solution, and  $A$  is the absorbance of the mixture by the HFO solution. Analyses of all samples were run in triplicate and averaged.

### Animals and treatment

Six-week-old male Kunming mice ( $20 \pm 2$  g) from Jinan Pengyue Laboratory Animal Breeding (Jinan, China) were kept in polypropylene cages and were maintained at  $25 \pm 2^\circ\text{C}$  and  $55 \pm 10\%$  relative humidity with equal 12-h light/dark cycles. All animals were on a standard laboratory feed and water *ad libitum*. All procedures were approved by the Animal Welfare Ethics Committee of the Shandong Academy of Sciences.

After adaptive feeding for 7 days, the mice were randomly divided into 5 groups. The grouping and administration are as follows ( $n=6$ ): normal control (NC; Physiological Saline), model group (MG; Physiological Saline), positive control (PC; 10 mg/mL Silymarin) and two groups treated with HFO at a dosage of 300 mg/mL (HDG, high dose group) and 75 mg/mL (LDG, low dose group), respectively. All groups were administered with 0.2 mL of Physiological Saline, Silymarin or HFO with gastrointestinal administration for 10 consecutive days. Except for the NC group, all other groups were injected intraperitoneally (i.p.) with  $\text{CCl}_4$  (7 mL/kg BW, 0.2% in peanut oil) 2 hours after the last gastrointestinal administration, and mice in the NC group were injected with an equivalent volume of peanut oil (i.p.). All the mice were sacrificed after fasting and continuous water for 24 h. Blood was taken from the eyeballs and the serum was separated.

### Biochemistry experiments

The blood was collected and centrifuged at 3,500 r/min for 10 min, then the supernatant was taken. Malondialdehyde (MDA), superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) levels were measured on an automatic biochemical analyzer (7500, Hitachi, Tokyo, Japan) for the antioxidant activity *in vitro* on an automatic biochemical analyzer (7500, Hitachi), according to the instructions of assay kits (Nanjing Jiancheng Bioengineering Institute, Nanjing, China).

### Data processing and analysis


IBM SPSS Statistics 23 software is used for the analysis of the experimental data by using analysis of variance (ANOVA)

and the difference analysis of the significance at the 0.05 level between means were determined by Duncan's multiple range test. The analysis of K and R (the result of extreme analysis) in orthogonal analysis and all the charts were processed using Office Excel 2019.


## Results and Discussion

### Optimization of enzymolysis conditions

Taking F ratio and DH as indicators, the hydrolysis conditions such as reaction time, the amount of enzyme, temperature and pH were optimized as illustrated in Fig. 1 (pepsin) and Fig. 2 (flavourzyme). When the pH, temperature and amount of enzyme were constant, DH showed a gradual increase. In contrast, F ratio showed an initial increase followed by a decline with the prolongation of hydrolysis time (Figs. 1A and 2A). However, changes to both F ratio and DH were insignificant. This is because the enzyme is more active at the beginning of hydrolysis, and with the progress of hydrolysis, the amount of enzyme and its combination with substrate diminished. The increase of AAA in hydrolyzed liquid resulted in the drop of F ratio. The DH value increased and reached saturation finally due to the increase in the number of active sites of enzymes in the hydrolysis system, and the F ratio reached its maximum at the addition of 1% (w/w) pepsin (Fig. 1B) and 0.5% (w/w) flavourzyme (Fig. 2B). As seen from Figs. 1C and 2C, both DH and F ratio showed a tendency of increasing first and then decreasing with the rise of temperature. Pepsin had the highest DH and F ratio at 50°C, while flavourzyme showed maximum



**Fig. 1.** Effects of different factors on enzymolysis by pepsin. <sup>A-E, a-e</sup> Different letters indicate a significant difference (p<0.05) among the different enzymatic hydrolysis conditions and the error bars represent the SD. F ratio, Fischer's ratio; DH, degree of hydrolysis.



**Fig. 2.** Effects of different factors on enzymolysis by flavourzyme. <sup>A-D, a-d</sup> Different letters indicate a significant difference ( $p < 0.05$ ) among the different enzymatic hydrolysis conditions and the error bars represent the SD. F ratio, Fischer's ratio; DH, degree of hydrolysis.

values at 45°C. When the range of pH change is not large, the activity of enzymes is affected by the dissociation of the relevant chemical groups at the active site (Li et al., 2019). Both the DH and F ratio of pepsin hydrolysis occur at pH 2 (Fig. 1D) and the maximum ratio of flavourzyme DH and F ratio at pH 7 (Fig. 2D). In view of orthogonal analysis (Tables 3 and 4), the range analysis revealed that the influential extent of the four factors to DH for pepsin and flavourzyme were  $C > B > D > A$ , and the optimal conditions were A2B2C2D2 and A2B1C2D1, respectively. The DH of pepsin obtained in three parallel experiments under optimal conditions were 15.93%, 15.52%, and 15.77%, with an average of 15.74%, and that of flavourzyme were 46.64%, 49.86%, and 47.53%, with an average of 48.01%. Based on the above results, the enzymatic conditions in this study were selected as follows. Pepsin (1%, w/w) was used to hydrolyze the goat whey at pH 2 and 50°C for 2 h. Afterwards, flavourzyme (0.5%, w/w) was used to hydrolyze the former hydrolysates at pH 7 and 45°C for 3 h.

### Optimization of aromatic amino acid (AAA) adsorption with activated carbon

Activated carbon is preferred for AAA adsorption because the benzene ring in AAA is mediated in combination with the graphite microcrystal structure of the activated carbon surface (Majidi and Karami, 2015; Rahmani et al., 2016). As can be seen from Fig. 3A, with the extension of adsorption time, the F ratio of the solution showed the tendency of rises followed by drops, and the F ratio peaked at 2 h of adsorption. The adsorption with activated carbon is better under the acidic conditions, for hydrolytes have a lower solubility at acidic pH, which can facilitate the efficient adsorption of AAA onto activated carbon (Jiang et al., 2017; Li et al., 2019). The results in this study showed that the maximum F ratio was reached at pH 2 (Fig. 3B).

**Table 3.** The range analysis of pepsin on DH obtained from the L9 (3<sup>4</sup>) orthogonal experiment

No.	(A)	(B)	(C)	(D)	DH (%)
1	1	1	1	1	8.80
2	1	2	2	2	15.64
3	1	3	3	3	6.39
4	2	1	2	3	11.75
5	2	2	3	1	10.93
6	2	3	1	2	12.09
7	3	1	3	2	8.30
8	3	2	1	3	11.90
9	3	3	2	1	9.75
K1	10.28	9.62	10.93	9.83	
K2	11.59	12.82	12.38	12.01	
K3	9.98	9.41	8.54	10.01	
Best level	A2	B2	C2	D2	
R	1.61	3.41	3.84	2.18	
R order	C>B>D>A				


Four controllable variables, including time (A), temperature (B), pH (C), and the amount of pepsin added (D) were selected for optimization. DH, degree of hydrolysis.

**Table 4.** The range analysis of flavourzyme on DH obtained from the L9 (3<sup>4</sup>) orthogonal experiment

No.	(A)	(B)	(C)	(D)	DH (%)
1	1	1	1	1	42.64
2	1	2	2	2	40.98
3	1	3	3	3	32.47
4	2	1	2	3	46.76
5	2	2	3	1	34.65
6	2	3	1	2	35.34
7	3	1	3	2	35.96
8	3	2	1	3	33.56
9	3	3	2	1	43.45
K1	38.70	41.79	37.18	40.25	
K2	38.92	36.4	43.73	37.43	
K3	37.66	37.09	34.36	37.60	
Best level	A2	B1	C2	D1	
R	1.26	5.39	9.37	2.65	
R order	C>B>D>A				

Four controllable variables, including time (A), temperature (B), pH (C), and the amount of flavourzyme added (D) were selected for optimization. DH, degree of hydrolysis.





**Fig. 3.** Effects of different factors on adsorption of AAA by activated carbon. <sup>a-d</sup> Different letters indicate a significant difference ( $p < 0.05$ ) among different conditions and the error bars represent the SD. F ratio, Fischer's ratio; AAA, aromatic amino acid.

The addition of active carbon also affected the F ratio, according to Fig. 3C. The carbon to liquid ratio of 1:5 was shown to be the most ideal. Under the condition of constant pH and duration, the adsorption of active carbons onto AAA may be inadequate when the amount of active carbon is not enough. As the activation of excessive carbon can cause BCAA to be adsorbed, both cases would result in a decrease in F ratio. The results of the orthogonal analysis are shown in Table 5, the range analysis revealed that the influential extent of the three factors to F ratio for active carbon was  $B > A > C$  and the optimal conditions were A2B2C2. And the F ratio of the solution obtained from three parallel experiments under optimal conditions were 24.43, 27.35, and 27.18, with an average of 26.32. To sum up, the optimized condition is pH 2 and carbon to liquid ratio of 1:5 for 2 h.

**Table 5.** The range analysis of active carbon on F ratio obtained from the L9 ( $3^3$ ) orthogonal experiment

No.	(A)	(B)	(C)	F ratio
1	1	1	1	12.09
2	1	2	2	23.48
3	1	3	3	14.23
4	2	1	2	21.50
5	2	2	3	25.79
6	2	3	1	18.98
7	3	1	3	13.89
8	3	2	1	19.63
9	3	3	2	16.90
K1	16.60	15.83	16.90	
K2	22.09	22.97	20.63	
K3	16.81	16.70	17.97	
Best level	A2	B2	C2	
R	5.49	7.14	3.73	
R order	B>A>C			

Three controllable variables, including time (A), temperature (B), and pH (C) were selected for optimization. F ratio, Fischer's ratio.

### Amino acid and molecular weight distribution

The amino acid profile of the HFO mixture was listed in Fig. 4A and Table 6. The BCAA content in the solution was 101.568 ng/mL and the phenylalanine content was only 3.752 ng/mL. BCAA accounted for 23.32% of total amino acids while AAA accounted for only 0.86%. The F ratio was calculated at 27.070, which is very close to the value of 26.32 calculated with absorbance values in 2.2.2. The content of Lys was 50.201 ng/mL, which can make up for the deficiency of lysine content in edible grain food. His, Tyr, Met, and Cys have been reported to have antioxidant activity (Xie et al., 2008). The results showed that when the amino acids have a high BCAA content and low AAA content, HFO mixture may have such functions as anti-fatigue and liver protection (Kawaguchi and Torimura, 2019; Mikulski et al., 2015; Park et al., 2020). HFO might be a good active ingredient to be added to the diet of patients with Phenylketonuria to reduce the intake of phenylalanine (Lichter-Konecki and Vockley, 2019). In addition, a total of 171.736 ng/mL of human essential amino acids were identified in this mixture. It might be reasonable to infer that HFO mixture obtained in this study was a hydrolysate possessing many physiological functions and nutritional value.

As illustrated in Fig. 4B, the hydrolysates showed three peaks: Peak 1, from 17.38 to 18.83 min, with the average Mw of 790, including about 4 to 7 amino acid residues of the medium peptide segment, accounting for 16.73%; Peak 2, from




Fig. 4. The results of amino acid analysis and the molecular weight (Mw) distribution.

**Table 6. Amino acid composition of HFO**


Amino acid	Concentration (ng/mL)	Type of amino acid
Aspartic acid (Asp)	15.520	
Threonine (Thr)	14.760	
Serine (Ser)	6.521	
Glutamate (Glu)	38.199	
Glycine (Gly)	25.330	
Alanine (Ala)	123.091	
Cysteine (Cys)	27.063	
Valine (Val)	48.423	BCAA
Methionine (Met)	5.207	
Isoleucine (Ile)	30.114	BCAA
Leucine (Leu)	23.031	BCAA
Tyrosine (Tyr)	3.752	AAA
Phenylalanine (Phe)	0.000	AAA
Lysine (Lys)	50.201	
Histidine (His)	6.542	
Tryptophan (Trp)	0.000	AAA
Arginine (Arg)	17.757	
BCAA	101.568	
AAA	3.752	
F ratio	27.070	

HFO, high Fischer's ratio oligopeptides from goat whey; F ratio, Fischer ratio; BCAA, branched-chain amino acids; AAA, aromatic amino acid.

18.83 to 20.51 min, with the average Mw of 236, dipeptides and tripeptides accounting for 60.98% of the total amino acids; Peak 3, from 20.51 to 22.47 min, with the average Mw of 29, which may be residual salts, accounting for 22.29%. Based on the result of amino acid composition, it can be concluded that HFO were obtained from goat whey after hydrolysis and adsorption. It is easier for small-molecular peptides such as HFO to be absorbed by human body than proteins and free amino acids. In addition, previous studies have reported that  $\alpha$ -leaf protein with a Mw of below 1,000 Da has antioxidant properties (Xie et al., 2008), the hydrolysate with the strongest hydroxyl RSA in soybean protein components is mainly the peptides of 0.7–0.8 kDa (Moure et al., 2006). The Mw of HFO obtained in this study was similar to that of the peptides reported previously. Therefore, it was inferred that HFO might show good oxidation resistance due to its small Mw.

### Free radical scavenging activity (RSA) *in vitro*

DPPH free radical is characterized by stability at room temperature, which is different from most other free radicals that are highly active but short-lived. The ability of HFO to scavenge DPPH was verified using ascorbic acid as the PC, as represented in Fig. 5A. There was an increase in the scavenging rate of DPPH with the increase of HFO concentration. A similar result has been reported for  $\alpha$ -leaf protein hydrolysate (Xie et al., 2008). When the concentration reached 1.6 mg/mL, the scavenging rate tended to stabilize, and remained at about 77% ( $p > 0.05$ ), which indicated that the maximum activity of



**Fig. 5. Scavenging activity of HFO on DPPH, ABTS and hydroxyl free radicals.** Means were performed for three samples. <sup>a-e</sup> Different letters indicate a significant difference ( $p < 0.05$ ) among different concentration and the error bars represent the SD. HFO, high Fischer's ratio oligopeptides from goat whey; DPPH, 1,1-diphenyl-2-picrylhydrazyl; ABTS, 3-ethylbenzthiazoline-6-sulphonate.


HFO in scavenging DPPH was observed at the concentration of 1.6 mg/mL. At the same concentration, the ascorbic acid showed a DPPH scavenging rate of over 95% ( $p > 0.05$ ).

Antioxidants inhibit lipid peroxidation through blocking free radical chain reactions and chelating metal ions that promote lipid oxidation (Zheng et al., 2016). As is shown in Fig. 5B, the RSA of ABTS free radicals increased with the rise of HFO concentration. At the concentration of 0.5 mg/mL, the RSA of ABTS free radicals reached 99.63%, which was the same as the scavenging rate of ascorbic acid. It indicated that HFO had a strong ability of scavenging ABTS free radicals, namely strong antioxidant activity. The ABTS scavenging rate of HFO was much higher than that of the feather keratin antioxidant peptides reported previously (Fontoura et al., 2019). In the amino acid sequencing results obtained in this study, the concentration of Cys was 27.063 ng/mL, consistent with the finding that Cys plays a major role in the scavenging of ABTS (Huo et al., 2020).

Hydroxyl free radicals are the most reactive in comparison with other oxygen free radicals and have been proven to be highly damaging species in free radical pathology. It can react with almost all substances in cells and cause serious damage (Zhang et al., 2012). Therefore, the effective removal of hydroxyl free radicals may be a valid defense for the living body against various diseases. As is shown in Fig. 5C, the scavenging ability of HFO solutions against hydroxyl free radicals varied with the HFO concentration ( $p < 0.05$ ). When the HFO concentration was at 0.25 mg/mL, high hydroxyl scavenging rate was observed (92.31%), which was slightly lower than that of the same ascorbic acid concentration (98.53%). This result was much higher than the scavenging rate of soybean protein hydrolysate (69.75%). Soybean protein hydrolytes with the strongest ability of scavenging hydroxyl free radicals mainly consist of peptides from 0.7 to 0.8 kDa, which also indicated that the antioxidant activity of HFO was correlated with the low molecular weight. The high hydroxyl radical scavenging ability of HFO also may be due to their susceptibility to oxidation by extremely reactive hydroxyl radicals, enhancing its antioxidant activity and protecting the body's ability of preventing hydroxyl free radical damage (Moure et al., 2006). The results of HFO about the hydroxyl RSA can provide useful information for antioxidant activity (Kitts, 2005).

### Antioxidant activity of high Fischer's ratio oligopeptides from goat whey (HFO) *in vivo*

Reactive oxygen species (ROS) are usually produced upon liver injury induced by  $\text{CCl}_4$ , resulting in lipid peroxidation and oxidative stress (Yuan et al., 2005; Zelko et al., 2002). So in this study,  $\text{CCl}_4$  was selected to induce oxidative stress in mice (Fig. 6). Compared with the NC group, there was significant increase in MDA level in the MG ( $p < 0.05$ ; Fig. 6A), which indicated that oxidative stress was generated in mice after  $\text{CCl}_4$  induction and this model can be used to evaluate the



**Fig. 6.** Effect of HFO on the MDA SOD and GSH-Px level in mice. <sup>A, B, a, b</sup> Different letters indicate a significant difference ( $p < 0.05$ ) among different group and the error bars represent the SD. HFO, high Fischer's ratio oligopeptides from goat whey; MDA, malondialdehyde; NC, normal control; MG, model group; PC, positive control; HDG, high dose group; LDG, low dose group; SOD, superoxide dismutase; GSH-Px, glutathione peroxidase.

antioxidant capacity of HFO *in vivo*. Compared with the MG group, HDG and LDG group showed significant declines in the level of MDA ( $p < 0.05$ ). As a product of membrane reaction between free radicals and unsaturated fatty acids, MDA reflects the sensitivity of lipid peroxidation and the degree of cell damage (Cemek et al., 2010; Meng et al., 2019). The results of this study showed that the intervention of HFO could delay the accumulation of lipid peroxides *in vivo*. Oxidative damage induced by  $\text{CCl}_4$  can not only cause changes in MDA, but also detect changes in oxidative stress parameters such as SOD and GSH-Px (Hamid et al., 2017; Lin et al., 2008). Antioxidant enzymes (SOD and GSH-Px) can convert peroxides into small or harmless substances (Zhang et al., 2003). The antioxidant enzymes are considered the main defense agent against oxidative damage of biological macromolecules, inhibiting the chain reaction of lipid peroxidation by effectively removing the production of ROS (Liu et al., 2018). As is shown in Fig. 6B, a decrease in SOD and GSH-Px was observed in MG group, indicating that excessive ROS may inhibit the development of antioxidant enzymes. Compared with MG group, there were significant increases in SOD and GSH-Px for HDG group and SOD for LDG ( $p < 0.05$ ), but GSH-Px of LDG showed no significant differences from that of MG. This observation may be due to the scavenging impact on free radicals by antioxidant enzymes. The results showed that HFO could alleviate oxidative stress induced by  $\text{CCl}_4$ . This may be related to the amino acid composition of HFO and the molecular weight distribution. Leu and Ala could release protons easily to electron-deficient free radicals, exhibiting excellent free radical scavenging ability and metal ion chelation ability. Moreover, oligopeptides can be easily absorbed by the intestine to exert an excellent antioxidant effect in the body (Jiang et al., 2020; Li et al., 2008; Wu et al., 2013). A research has shown that oral administration of small peptides can improve the antioxidant activity in ultraviolet-induced photoaging mice, which is consistent with the results of this study.

## Conclusion

In this study, goat whey was used to prepare high F ratio oligopeptides with a F ratio of 27.070, and the optimized enzymatic hydrolysis and adsorption condition was determined. The HFO have a high BCAA concentration and low molecular weight of less than 1,000 Da. Free radical scavenging results showed that HFO has the ability of scavenging DPPH, ABTS and hydroxyl free radicals, and the characteristic of concentration-dependence. Animal experiment shows that HFO could reduce the MDA produced by oxidative stress in mice and elevate the level of SOD and GSH-Px *in vivo*. This work can not only offer an important basis for the high-value processing of goat whey, but also for the diversification of HFO sources, expanding the source

of antioxidant peptides, while providing a theoretical basis for the development of HFO for special medical foods.

## Conflicts of Interest

The authors declare no potential conflicts of interest.

## Acknowledgements

This work was funded by the Key Technology Research and Development Program of Shandong province (2019YYSP025); Natural Science Foundation of Shandong Province (ZR2020MC210); the Open Project Program of State Key Laboratory of Dairy Biotechnology (SKLDB2021-003); Shandong Major Agricultural Technology Innovation Project (SD2019ZZ006).

## Author Contributions

Conceptualization: Wang C, Jiang H. Data curation: Qin Y, Cheng M, Zhang X. Formal analysis: Qin Y, Cheng M, Wang C. Methodology: Qin Y, Fan X, Shao X. Software: Fan X, Shao X, Zhang X. Validation: Fan X, Shao X, Zhang X. Investigation: Cheng M, Wang C, Jiang H. Writing - original draft: Qin Y. Writing - review & editing: Qin Y, Cheng M, Fan X, Shao X, Wang C, Jiang H, Zhang X.

## Ethics Approval

This research was approved by the Animal Welfare Ethics Committee of Shandong Academy of Sciences (IACUC Issue No. SWS20220227).

## References

- Bordenave S, Sannier F, Ricart G, Piot JM. 2000. Characterization of a goat whey peptic hydrolysate produced by an ultrafiltration membrane enzymic reactor. *J Dairy Res* 67:551-559.
- Cemek M, Aymelek F, Büyükkuroğlu ME, Karaca T, Büyükben A, Yilmaz F. 2010. Protective potential of royal jelly against carbon tetrachloride induced-toxicity and changes in the serum sialic acid levels. *Food Chem Toxicol* 48:2827-2832.
- Cermakova L, Kopecka I, Pivokonsky M, Pivokonska L, Janda V. 2017. Removal of cyanobacterial amino acids in water treatment by activated carbon adsorption. *Sep Purif Technol* 173:330-338.
- Chen D, Li XY, Zhao X, Qin YS, Zhang XX, Li J, Wang JM, Wang CF. 2019. Proteomics and microstructure profiling of goat milk protein after homogenization. *J Dairy Sci* 102:3839-3850.
- Da Silva MS, Bigo C, Barbier O, Rudkowska I. 2017. Whey protein hydrolysate and branched-chain amino acids downregulate inflammation-related genes in vascular endothelial cells. *Nutr Res* 38:43-51.
- Duan X, Ocen D, Wu F, Li M, Yang N, Xu J, Chen H, Huang L, Jin Z, Xu X. 2014. Purification and characterization of a natural antioxidant peptide from fertilized eggs. *Food Res Int* 56:18-24.
- Fontoura R, Daroit DJ, Corrêa APF, Moresco KS, Santi L, Beys-da-Silva WO, Yates JR, Moreira JCF, Brandelli A. 2019. Characterization of a novel antioxidant peptide from feather keratin hydrolysates. *New Biotechnol* 49:71-76.
- Garay PA, Villalva FJ, Paz NF, de Oliveira EG, Ibarguren C, Alcocer JC, Curti CA, Ramón AN. 2021. Formulation of a

- protein fortified drink based on goat milk whey for athletes. *Small Rumin Res* 201:106418.
- Hamid M, Liu D, Abdulrahim Y, Liu Y, Qian G, Khan A, Gan F, Huang K. 2017. Amelioration of CCl<sub>4</sub>-induced liver injury in rats by selenizing astragalus polysaccharides: Role of proinflammatory cytokines, oxidative stress and hepatic stellate cells. *Res Vet Sci* 114:202-211.
- Huo J, Luo X, Huang M, Wu J, Zhang J, Liu X, Li H, Sun X. 2020. Identification and antioxidant activity of a novel peptide from Baijiu. *Int J Pept Res Ther* 26:1199-1210.
- Jiang W, Yang Y, Tao R, Jiang X, Zhou X, Zhou Z. 2017. Preparation of a novel chitosan-based biosorbent cross-linked with phenethylamine for adsorption of aromatic amino acids. *Carbohydr Polym* 176:236-245.
- Jiang X, Cui Z, Wang L, Xu H, Zhang Y. 2020. Production of bioactive peptides from corn gluten meal by solid-state fermentation with *Bacillus subtilis* MTCC5480 and evaluation of its antioxidant capacity *in vivo*. *LWT-Food Sci Technol* 131:109767.
- Kawaguchi T, Torimura T. 2019. Branched chain amino acids: A factor for zone 3 steatosis in non-alcoholic fatty liver disease. *Hepatol Res* 49:841-843.
- Kerasiote E, Stagos D, Priftis A, Aivazidis S, Tsatsakis AM, Hayes AW, Kouretas D. 2014. Antioxidant effects of whey protein on muscle c2c12 cells. *Food Chem* 155:271-278.
- Kerasiote E, Veskoukis A, Virgiliou C, Theodoridis G, Taitzoglou I, Kouretas D. 2019. The strong antioxidant sheep/goat whey protein protects against mtor overactivation in rats: A mode of action mimicking fasting. *Antioxidants* 8:71.
- Kitts DD. 2005. Antioxidant properties of casein-phosphopeptides. *Trends Food Sci Technol* 16:549-554.
- Li T, Tian Y, Sun F, Wang Z, Zhou N. 2019. Preparation of high Fischer's ratio corn oligopeptides using directed enzymatic hydrolysis combined with adsorption of aromatic amino acids for efficient liver injury repair. *Process Biochem* 84:60-72.
- Li X, Lin J, Gao Y, Han W, Chen D. 2012. Antioxidant activity and mechanism of *Rhizoma cimicifugae*. *Chem Cent J* 6:140.
- Li XX, Han LJ, Chen LJ. 2008. *In vitro* antioxidant activity of protein hydrolysates prepared from corn gluten meal. *J Sci Food Agric* 88:1660-1666.
- Lichter-Konecki U, Vockley J. 2019. Phenylketonuria: Current treatments and future developments. *Drugs* 79:495-500.
- Lin HM, Tseng HC, Wang CJ, Lin JJ, Lo CW, Chou FP. 2008. Hepatoprotective effects of *Solanum nigrum* Linn extract against CCl<sub>4</sub>-induced oxidative damage in rats. *Chem Biol Interact* 171:283-293.
- Liu Y, Liu Q, Hesketh J, Huang D, Gan F, Hao S, Tang S, Guo Y, Huang K. 2018. Protective effects of selenium-glutathione-enriched probiotics on CCl<sub>4</sub>-induced liver fibrosis. *J Nutr Biochem* 58:138-149.
- Majidi R, Karami AR. 2015. Aromatic amino acids adsorption on graphyne: A density functional theory study. *Struct Chem* 26:5-10.
- Mat DJL, Souchon I, Michon C, Feunteun SL. 2020. Gastro-intestinal *in vitro* digestions of protein emulsions monitored by pH-stat: Influence of structural properties and interplay between proteolysis and lipolysis. *Food Chem* 311:125946.
- Meng X, Wang Z, Liang S, Tang Z, Liu J, Xin Y, Kuang H, Wang Q. 2019. Hepatoprotective effect of a polysaccharide from radix *Cyathulae officinalis* Kuan against CCl<sub>4</sub>-induced acute liver injury in rat. *Int J Biol Macromol* 132:1057-1067.
- Mikulski T, Dabrowski J, Hilgier W, Ziemia A, Krzeminski K. 2015. Effects of supplementation with branched chain amino acids and ornithine aspartate on plasma ammonia and central fatigue during exercise in healthy men. *Folia Neuropathol* 53:377-386.
- Moure A, Domínguez H, Parajó JC. 2006. Antioxidant properties of ultrafiltration-recovered soy protein fractions from industrial effluents and their hydrolysates. *Process Biochem* 41:447-456.
- Osman A, Goda HA, Abdel-Hamid M, Badran SM, Otte J. 2016. Antibacterial peptides generated by alcalase hydrolysis of

- goat whey. *LWT-Food Sci Technol* 65:480-486.
- Papademas P, Kotsaki P. 2019. Technological utilization of whey towards sustainable exploitation. *J Adv Dairy Res* 7:231.
- Park JG, Tak WY, Park SY, Kweon YO, Chung WJ, Jang BK, Bae SH, Lee HJ, Jang JY, Suk KT, Oh MJ, Heo J, Woo HY, Jang SY, Lee YR, Lee JS, Kim DY, Kim SH, Suh JI, Kim IH, Kang MK, Lee WK. 2020. Effects of branched-chain amino acid (BCAA) supplementation on the progression of advanced liver disease: A Korean nationwide, multicenter, prospective, observational, cohort study. *Nutrients* 12:1429.
- Pedroche J, Yust MM, Lqari H, Girón-Calle J, Vioque J, Alaiz M, Millán F. 2004. Production and characterization of casein hydrolysates with a high amino acid Fischer's ratio using immobilized proteases. *Int Dairy J* 14:527-533.
- Rafiq S, Huma N, Pasha I, Sameen A, Mukhtar O, Khan MI. 2015. Chemical composition, nitrogen fractions and amino acids profile of milk from different animal species. *Asian-Australas J Anim Sci* 29:1022-1028.
- Rahmani F, Nouranian S, Mahdavi M, Al-Ostaz A. 2016. Molecular simulation insights on the *in vacuo* adsorption of amino acids on graphene oxide surfaces with varying surface oxygen densities. *J Nanopart Res* 18:320.
- Ranathunga S, Rajapakse N, Kim SK. 2006. Purification and characterization of antioxidative peptide derived from muscle of conger eel (*Conger myriaster*). *Eur Food Res Technol* 222:310-315.
- Salma NU, Govindaraju K, Kumar BSG, Muthukumar SP, Lakshmi AJ. 2022. Ameliorative effect of enhanced Fischer ratio flaxseed protein hydrolysate in combination with antioxidant micronutrients on ethanol-induced hepatic damage in a rat model. *Br J Nutr* 127:696-710.
- Song W, Kong X, Hua Y, Li X, Zhang C, Chen Y. 2020. Antioxidant and antibacterial activity and *in vitro* digestion stability of cottonseed protein hydrolysates. *LWT-Food Sci Technol* 118:108724.
- Tang Q, Huang G. 2018. Preparation and antioxidant activities of cucurbit polysaccharide. *Int J Biol Macromol* 117:362-365.
- Wu Y, Tian Q, Li L, Khan MN, Yang X, Zhang Z, Hu X, Chen S. 2013. Inhibitory effect of antioxidant peptides derived from *Pinctada fucata* protein on ultraviolet-induced photoaging in mice. *J Funct Foods* 5:527-538.
- Xia J, He J, Xu J, Liu X, Qiu Z, Xu N, Su L. 2021. Direct conversion of cheese whey to poly malic acid by mixed culture of *Aureobasidium pullulans* and permeabilized *Kluyveromyces marxianus*. *Bioresour Technol* 337:125443.
- Xie Z, Huang J, Xu X, Jin Z. 2008. Antioxidant activity of peptides isolated from alfalfa leaf protein hydrolysate. *Food Chem* 111:370-376.
- Yan L, Shi T, Li Y, Liu X. 2021. Optimization of enzymatic hydrolysis and activated carbon adsorption for producing high Fischer ratio peptides from soy protein. *Int J Pept Res Ther* 27:1363-1372.
- Yuan YV, Bone DE, Carrington MF. 2005. Antioxidant activity of dulce (*Palmaria palmata*) extract evaluated *in vitro*. *Food Chem* 91:485-494.
- Zelko IN, Mariani TJ, Folz RJ. 2002. Superoxide dismutase multigene family: A comparison of the cuzn-sod (sod1), mn-sod (sod2), and ec-sod (sod3) gene structures, evolution, and expression. *Free Radic Biol Med* 33:337-349.
- Zhang Q, Li N, Zhou G, Lu X, Xu Z, Li Z. 2003. *In vivo* antioxidant activity of polysaccharide fraction from *Porphyra haitanensis* (Rhodophyta) in aging mice. *Pharmacol Res* 48:151-155.
- Zhang Y, Duan X, Zhuang Y. 2012. Purification and characterization of novel antioxidant peptides from enzymatic hydrolysates of tilapia (*Oreochromis niloticus*) skin gelatin. *Peptides* 38:13-21.
- Zheng L, Zhao Y, Dong H, Su G, Zhao M. 2016. Structure–activity relationship of antioxidant dipeptides: Dominant role of tyr, trp, cys and met residues. *J Funct Foods* 21:485-496.