



# Phytochemical constituents of *Lactuca serriola* leaves and their content analysis by HPLC-UV

Juree Kim<sup>1</sup> · Hak-Dong Lee<sup>1</sup> · Jungwon Choi<sup>1</sup> · Sanghyun Lee<sup>1,2,3</sup>

Received: 17 June 2022 / Accepted: 19 July 2022 / Published Online: 30 September 2022  
© The Korean Society for Applied Biological Chemistry 2022

**Abstract** This study aimed to identify the phytochemical constituents of *Lactuca serriola* leaves and perform quantitative analysis of the methanol (MeOH) extract of *L. serriola*, *L. indica*, *L. raddeana*, *L. sativa*, and *L. triangulata*. Six compounds were isolated from the MeOH extracts of *L. serriola* using open column chromatography and identified as protocatechuic acid (**1**), caffeic acid (**2**), cynaroside (**3**), apigenin 7-glucuronide (**4**), luteolin (**5**), and apigenin (**6**) using <sup>1</sup>H-, <sup>13</sup>C-nuclear magnetic resonance, and mass spectrometry. Quantitative analysis of the six compounds was performed on the MeOH extract of *Lactuca* species using high-performance liquid chromatography (HPLC) and an ultraviolet (UV). A reverse-phased column was used, and the UV absorbance was set to 280 nm. The contents of compounds **2** and **3** were the highest (1.58 and 2.64 mg/g ext., respectively) in *L. serriola* extracts. However, compounds **4** and **6** were higher (1.46 and 0.40 mg/g ext., respectively) in *L. triangulata*. These results provide quantitative data for the application of *Lactuca* species in the pharmaceutical and functional food industries.

**Keywords** High-performance liquid chromatography-ultraviolet · *Lactuca serriola* · Quantitative analysis

Sanghyun Lee (✉)  
E-mail: slee@cau.ac.kr

<sup>1</sup>Department of Plant Science and Technology, Chung-Ang University, Anseong 17546, Republic of Korea

<sup>2</sup>BET Research Institute, Chung-Ang University, Anseong 17546, Republic of Korea

<sup>3</sup>Natural Product Institute of Science and Technology, Anseong 17546, Republic of Korea

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Introduction

*Lactuca serriola* L. (Asteraceae) is an annual or biennial plant known as prickly lettuce [1]. *Lactuca serriola* is the closest wild relative of *L. sativa*, and is native to Asia, Europe, and North Africa [2]. It is a naturalized plant and is designated as an ecosystem-disturbing species in Korea. Unlike *L. sativa*, which is the common lettuce, the leaves of *L. serriola* are oblong and lanceolate, with fine spines along the veins and leaf edges. It has traditionally been used for multiple purposes, including as a sedative, purgative, diuretic, vasorelaxant, and demulcent, and it is used to treat bronchitis, gastrointestinal injury, asthma, and pertussis. In addition, it was recently found that the aerial parts of *L. serriola* have remarkable  $\alpha$ -glucosidase inhibitory [3] and antivenom activities [4]. *Lactuca serriola* contains lactucone, lactucin, lactic acids [5], alkaloids, oxalic acid, lactucopicrin [1], vitamins, beta-carotene, iron [6], sesquiterpene esters [7], flavonoids [8], and phenolic acids [4].

*Lactuca serriola* has been designated an ecosystem-disturbing species by the Ministry of Environment in Korea [9]. Ecosystem-disturbing species can be artificially or naturally introduced from foreign countries and causes disturbances in the balance of the ecosystem. Genetically modified organisms can be created through genetic modification with these species. In addition, *Rana catesbeiana* [10], *Ambrosia artemisiaefolia* var. *elatior* [11], and *Sicyos angulatus* [12] have also been identified as ecosystem-disturbing species in Korea. Ecosystem-disturbing species have rapid reproductive power and vigorous growth, which have a serious impact on the ecosystem and cause economic damage when they enter pastures and arable land. However, in recent years, they have been studied for use as resources [13,14]. Scientific research that supports their use is still lacking. Studies on *L. serriola* have focused on its photosynthetic characteristics [15,16], genetic differences [17], genetic transformation [18], and morphological variability [19]. Studies are needed to identify the active compounds or activities of *L. serriola*.

The objectives of this study were to identify the phytochemical constituents of *L. serriola* and to quantify the methanol (MeOH) extracts of *L. serriola*, *L. indica*, *L. raddeana*, *L. sativa*, and *L. triangulata* using high-performance liquid chromatography-ultraviolet (HPLC-UV).

## Materials and Methods

### Plant materials

Leaves of *L. serriola* (June, 2017) were collected from Jinwicheon, Pyeongtaek, Republic of Korea.

### Chemicals and apparatus

Ethyl acetate (EtOAc), *n*-hexane, chloroform (CHCl<sub>3</sub>), *n*-butanol (*n*-BuOH), and MeOH were purchased from Samchun Pure Chemicals (Pyeongtaek, Republic of Korea). Silica gel (200–400 mesh) (Merck Co., Darmstadt, Germany) and Sephadex LH-20 (Sigma-Aldrich Co., St. Louis, MO, USA) were used as the stationary phases for open column chromatography. Silica gel 60 F<sub>254</sub>-precoated thin-layer chromatography (TLC) (Merck KGaA, Darmstadt, Germany) was used to monitor the chromatographic separation. An Avance 500 MHz spectrometer (Bruker, Rheinstetten, Germany) was used for nuclear magnetic resonance (NMR) assignment and a JEOL-MS spectrometer (Jeol, Tokyo, Japan) was used for mass spectrometry data. Chromatographic analysis was performed using an HPLC system (Waters Alliance 2695 Separations Module, Milford, MA, USA) equipped with a pump, auto-sampler, and photodiode-array detection (Waters 996 Photodiode Array Detector). HPLC-grade solvents, such as water and acetonitrile, were purchased from J. T. Baker (Avantor, Radnor, PA, USA). Acetic acid (glacial, 100%) was purchased from Supelco® (Merck KGaA).

### Extraction, fractionation, and isolation

Dried leaves of *L. serriola* (1.7 kg) were extracted with MeOH under reflux at 88 °C for 3 h, three times in succession. After extraction, the solvents were filtered using filter paper and evaporated using a rotary evaporator (Sunileyela Co., Ltd., Seongnam, Republic of Korea) at 55 °C to obtain MeOH extract (300 g). The extract (187 g) was suspended in distilled water (H<sub>2</sub>O) and partitioned into polar fractions sequentially using *n*-hexane, CHCl<sub>3</sub>, EtOAc, and *n*-BuOH. Each partitioned layer of the solvent was evaporated to produce *n*-hexane (22.7 g), CHCl<sub>3</sub> (24.5 g), EtOAc (7.2 g), and *n*-BuOH (13.4 g) fractions.

Each fraction was separated using open column chromatography and silica gel was used as the stationary phase. Each fraction was eluted using a solvent mixture of a stepwise gradient system (100:0 v/v to 0:100 v/v). All small fractions on the TLC plates were visually confirmed by spraying them with 10% H<sub>2</sub>SO<sub>4</sub> in MeOH and heating them on a hot plate. The EtOAc fraction (5.2 g) was eluted with CHCl<sub>3</sub>/MeOH (100:0 v/v to 0:100 v/v)

using a stepwise gradient system to obtain 146 small fractions. Fractions were grouped into four sub-fractions (LSE1-4), and LSE1-3 were re-chromatographed over a Sephadex LH-20 column with a solvent system of H<sub>2</sub>O/MeOH (5:1 v/v to 1:5 v/v). LSE1 yielded 55 sub-fractions; sub-fractions (53-54) were recrystallized using MeOH to obtain compound 6. LSE2 yielded 79 sub-fractions; sub-fractions (50-60) were identified as compound 5. LSE3 yielded 94 sub-fractions; sub-fractions (31-33) were identified as compound 2. The *n*-BuOH (10 g) fraction was also eluted with CHCl<sub>3</sub>/MeOH to obtain 169 small fractions, which were then grouped into three sub-fractions (LSB1-3). LSB2 was re-chromatographed over a Sephadex LH-20 column with a solvent system H<sub>2</sub>O/MeOH (5:1 v/v to 1:5 v/v) to obtain 86 sub-fractions. Sub-fractions (52-55) were obtained using a solvent system of H<sub>2</sub>O/MeOH (1:1 v/v) and identified as compound 1. Sub-fractions (64-66) were recrystallized using MeOH to obtain compound 4. Sub-fractions (83-85) were identified as compound 3.

Compound 1: Light brown solid; <sup>1</sup>H-NMR (CD<sub>3</sub>OD): δ 6.76 (H-5, d, 8.0), 7.39 (H-6, dd, 2.0, 8.0), 7.42 (H-2, d, 2.0); <sup>13</sup>C-NMR (CD<sub>3</sub>OD): δ 115.8 (C-5), 117.7 (C-2), 123.6 (C-1), 125.4 (C-6), 146.1 (C-3), 150.7 (C-4), 171.9 (C-7).

Compound 2: Yellow amorphous solid; <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 6.15 (H-8, d, 15.5), 6.75 (H-5, d, 8.0), 6.94 (H-6, dd, 2.0, 8.0), 7.01 (H-2, d, 2.0), 7.37 (H-7, d, 15.5).

Compound 3: Yellow powder; <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 5.08 (H-1", d, 7.5), 6.43 (H-6, d, 2.0), 6.73 (H-3, s), 6.78 (H-8, d, 2.0), 6.86 (H-5', d, 8.5), 7.40 (H-2', d, 2.0), 7.44 (H-6', dd, 2.5, 8.5), 13.04 (5-OH, s); <sup>13</sup>C-NMR (DMSO-*d*<sub>6</sub>): δ 60.6 (C-6"), 69.5 (C-4"), 73.1 (C-2"), 76.4 (C-3"), 77.1 (C-5"), 94.7 (C-8), 99.5 (C-6), 99.8 (C-1"), 102.8 (C-3), 105.3 (C-10), 113.2 (C-2'), 115.9 (C-5'), 119.3 (C-6'), 121.5 (C-1'), 145.9 (C-3'), 150.1 (C-4'), 156.9 (C-9), 161.1 (C-5), 162.9 (C-7), 164.6 (C-2), 181.8 (C-4).

Compound 4: Yellow powder; <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 5.02 (H-1', d, 7.5), 6.41 (H-6, d, 1.5), 6.81 (H-8, d, 1.5), 6.83 (H-3, s), 6.89 (H-3',5', d, 8.5), 7.92 (H-2',6', d, 8.5), 13.00 (5-OH, s); <sup>13</sup>C-NMR (DMSO-*d*<sub>6</sub>): δ 72.0 (C-4"), 72.9 (C-2"), 73.6 (C-3"), 76.6 (C-5"), 94.6 (C-8), 99.6 (C-6), 99.6 (C-1"), 102.8 (C-3), 105.2 (C-10), 116.1 (C-3',5'), 121.0 (C-1'), 128.5 (C-2',6'), 156.9 (C-9), 161.0 (C-4'), 162.0 (C-5), 163.1 (C-7), 164.3 (C-2), 171.7 (C-6"), 181.9 (C-4).

Compound 5: Yellow powder; <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 6.17 (H-6, d, 2.0), 6.42 (H-8, d, 2.5), 6.66 (H-3, s), 6.87 (H-5', d, 8.5), 7.38 (H-2', d, 2.5), 7.41 (H-6', dd, 2.5, 8.5), 12.99 (5-OH, s).

Compound 6: Yellow powder; <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 6.01 (H-6, s), 6.28 (H-8, s), 6.66 (H-3, s), 6.90 (H-3',5', d, 9.0), 7.87 (H-2',6', d, 9.0), 12.93 (5-OH, s).

### Preparation of samples and stock solutions for HPLC

The MeOH extracts of *L. indica*, *L. raddeana*, *L. sativa*, and *L. triangulate* were purchased from Korea Research Institute of Bioscience and Biotechnology, Daejeon, Republic of Korea.

Sample stock solutions were prepared by dissolving each extract (30 mg/mL) in MeOH, sonicating for 20 min, and filtering through a 0.45  $\mu\text{m}$  polyvinylidene difluoride membrane. Stock solutions of the six standards (1 mg/mL) were prepared using the same procedure as above.

### HPLC conditions

The six compounds were quantified using HPLC-UV using a reverse-phase YMC-Pack Pro C18 column (4.6 $\times$ 250 mm, 5  $\mu\text{m}$ ) (YMC Korea Co. Ltd., Seongnam, Republic of Korea) maintained at 30  $^{\circ}\text{C}$ . All the samples were detected at a wavelength of 280 nm. The flow rate was 1 mL/min and the injection volume was 10  $\mu\text{L}$ . The mobile phases were 0.5% acetic acid in water (A) and acetonitrile (B). Samples and standards were eluted using a gradient program as follows: 0 min, 95% A; 22 min, 30% B; 29 min, 30% B; 32 min, 40% B; 42 min, 100% B; 45 min, 100% B; 47 min, 95% A; and 55 min, 95% A. All samples were injected in triplicate.


### Calibration curves

The working solutions were prepared by serially diluting the standard stock solution of each standard to the specified concentrations and were used to construct the calibration curve. The calibration functions of the standards were calculated from the peak area (Y) and concentration (X,  $\mu\text{g/mL}$ ), and the values are presented as means  $\pm$  standard deviation (n=3).

## Results and Discussion

The chemical structures of the six isolated compounds were identified using NMR and MS and compared with previous literature. Two phenolic acids and four flavonoids were isolated and identified from the MeOH extracts of *L. serriola* (Fig. 1).

The NMR spectra of compounds 1 and 2 indicated phenolic acids [protocatechuic acid (1) and caffeic acid (2)], and their spectral data were compared with those of previous studies [20–23]. Signals of the ABX splitting type in the aromatic ring were




Compound	R <sub>1</sub>	R <sub>2</sub>
3	Glc	OH
4	Gln	H
5	OH	OH
6	OH	H


**Fig. 1** Chemical structures of compounds 1-6: protocatechuic acid (1), caffeic acid (2), cynaroside (3), apigenin 7-glucuronide (4), luteolin (5), and apigenin (6)

observed in both compounds. The spectrum of compound 1 showed aromatic proton signals at  $\delta$  6.76 (d, 8.0), 7.39 (dd, 2.0, 8.0), and 7.42 (d, 2.0) and carbonyl carbon signals at  $\delta$  171.9. Furthermore, the  $^1\text{H-NMR}$  spectrum of compound 2 showed trans-ene double bond signals at  $\delta$  6.15 (d, 15.5) and 7.37 (d, 15.5).

Compounds 3-6 were identified as flavonoids and their derivatives [cynaroside (3), apigenin 7-glucuronide (4), luteolin (5), and apigenin (6)] by comparing their spectral data with those of previous studies [24–29]. “Flavonoids” belong to a class of secondary metabolites of polyphenols found in plants, and originate from the Latin word “*flavus*”, meaning yellow. Flavonoids have a basic skeleton of 15-carbons and are composed of two phenyl rings and one heterocyclic ring containing an embedded oxygen [30]. The spectra of compounds 3-6 indicated a 5-OH signal of the flavonoid A-ring and a carbonyl (C=O) signal. Compound 3 had a glucose attached to compound 5, and



**Fig. 2** HPLC chromatograms of compounds 1-6: protocatechuic acid (1), caffeic acid (2), cynaroside (3), apigenin 7-glucuronide (4), luteolin (5), and apigenin (6)



**Fig. 3** HPLC chromatograms of the methanol extracts [protocatechuic acid (1), caffeic acid (2), cynaroside (3), apigenin 7-glucuronide (4), luteolin (5), and apigenin (6)] of (A) *Lactuca serriola*, (B) *L. indica*, (C) *L. raddeana*, (D) *L. sativa*, and (E) *L. triangulata*

**Table 1** Calibration curves of methanol extracts of *Lactuca serriola*: protocatechuic acid (1), caffeic acid (2), cynaroside (3), apigenin 7-glucuronide (4), luteolin (5), and apigenin (6)

Compound	t <sub>R</sub> <sup>a</sup>	Calibration equation <sup>b</sup>	Correlation factor, r <sup>2c</sup>
1	9.5	Y = 14486X - 2266.9	0.9999
2	14.5	Y = 26598X - 10400	0.9994
3	19.7	Y = 15814X + 16652	0.9992
4	26.5	Y = 16062X - 29206	0.9991
5	28.8	Y = 20652X + 3588.9	0.9992
6	34.9	Y = 25529X + 3059.4	0.9994

<sup>a</sup> t<sub>R</sub> = retention time<sup>b</sup> Y = peak area, X = concentration of the standard (µg/mL)<sup>c</sup> r<sup>2</sup> = correlation coefficient for five data points in the calibration curve**Table 2** Content of protocatechuic acid (1), caffeic acid (2), cynaroside (3), apigenin 7-glucuronide (4), luteolin (5), and apigenin (6) in methanol extracts of *Lactuca* species

Sample	Content (mg/g ext.)					
	1	2	3	4	5	6
<i>L. serriola</i>	0.29±0.05	1.58±0.36	2.64±0.45	0.21±0.02	0.30±0.02	0.01±0.00
<i>L. indica</i>	0.35±0.00	0.29±0.01	0.46±0.02	0.51±0.01	0.01±0.01	tr
<i>L. raddeana</i>	0.10±0.01	ND	0.04±0.01	0.23±0.04	0.01±0.00	tr
<i>L. sativa</i>	0.09±0.00	0.09±0.00	0.22±0.01	ND	ND	ND
<i>L. triangulata</i>	0.32±0.01	ND	0.13±0.01	1.46±0.08	0.22±0.03	0.40±0.02

ND, not detected; tr, trace

an anomeric proton signal was observed at  $\delta$  5.08 (d, 7.5). The <sup>13</sup>C-NMR signals of glucose and anomeric carbon were indicated at  $\delta$  60.6, 69.5, 73.1, 76.4, 77.1, and 99.8. Compound 4 had glucuronic acid combined with compound 6, and the carbon signal at  $\delta$  171.7 (C-6'') indicated a glucuronic acid moiety. Compound 5 showed proton signals at  $\delta$  6.87 (d, 8.5), 7.38 (d, 2.5), and 7.41 (dd, 2.5, 8.5), indicating the ABX splitting type of the flavonoid B-ring. Compound 6 showed proton signals at  $\delta$  6.90 (d, 9.0) and 7.87 (d, 9.0), indicating A2B2 splitting of the B-ring.

Six compounds were quantified in the extract of *L. serriola*, and extracts of four plants of the same genus were also quantitatively analyzed and their contents were compared. The six compounds were well separated by HPLC, and the chromatograms of the standards and extracts are shown in Fig. 2 and Fig. 3, respectively. The correlation coefficients (r<sup>2</sup>) of the standards ranged from 0.9991 to 0.9999 (Table 1). The contents of each compound in the extracts of *Lactuca* species are shown in Table 2, and all six compounds were detected in the extract of *L. serriola*. There was a high content of compound 3 (2.64 mg/g ext.) in the *L. serriola* extract, which was also higher than the other extracts; compound 2 was high. The other compounds were present in relatively small amounts in *L. serriola* extracts. Unlike *L. serriola*, the content of compound 4 was the highest in *L. triangulata* compared to other extracts (1.46 mg/g ext.). Compound 1 was detected in all extracts, and the contents were similar. In contrast to the other extracts, only compounds 1-3 were detected in *L. sativa*.

In other studies, several flavonoids and phenolic acids, including chlorogenic acid, *p*-coumaric acid, luteolin, caffeic acid, and cynaroside, have been isolated from *L. serriola* and their antioxidant activity has been reported [8,14,31]. Wang et al. isolated and identified compounds 1-3 from *L. indica* and evaluated the free radical-scavenging activity in the extract [32]. Nugroho et al. quantified phenolic compounds, including luteolin and cynaroside, from the extract of *L. raddeana*, suggesting its anti-obesity activity [33]. *Lactuca sativa* contains vitamins, carotenoids, and phenolic compounds [34], and polyphenols are mainly composed of caffeic acid derivatives and flavonoids [35]. Kim et al. conducted an HPLC analysis from the seventeen extracts of the Compositae plants; among them, *L. triangulata* showed the highest total phenolic content and peroxynitrite-scavenging activity [36]. Although there are several studies on *L. serriola*, their photosynthetic characteristics and genetic and morphological variability have been studied, and studies on the phytochemical constituents of *L. serriola* are still lacking. In addition, *L. serriola* is an ecosystem-disturbing species with high research value and potential as a usable resource.

In conclusion, we isolated six phytochemical constituents from *L. serriola* and performed quantitative analysis of the MeOH extracts of *L. serriola*, *L. indica*, *L. raddeana*, *L. sativa*, and *L. triangulata* using HPLC-UV. These results can be used as key data for economic and industrial utilization of *L. serriola* and *Lactuca* species.

**Acknowledgments** This research was supported by a grant from Natural Product Institute of Science and Technology, Anseong 17546, Republic of Korea. We gratefully acknowledge Mr. Dong-gun Woo for the collection of the plant material and fundamental side experiments.

## References

- Baquer SR (1989) Medicinal and poisonous plants of Pakistan. Karachi, Pakistan
- Mucina L (1978) Ruderal communities with dominant species *Lactuca serriola*. *Biologia (Bratislava)* 33: 809–819
- Hussein N, Amen Y, Abdel Bar F, Halim A, Saad H-E (2020) Antioxidants and  $\alpha$ -glucosidase inhibitors from *Lactuca serriola* L. *Rec Nat Prod* 14: 410–415
- Bouimeja B, Yetongnon K, Touloun O, Berrougui H, Laaradia M, Ouanaimi F, Chait A, Boumezzough A (2019) Studies on antivenom activity of *Lactuca serriola* methanolic extract against *Buthus atlantis* scorpion venom by in vivo methods. *S Afr J Bot* 125: 270–279. doi: 10.1016/j.sajb.2019.07.044
- Dymock W, Warden CJH, Hooper D (1983) *Pharmacographia Indica: A history of the principal drugs of vegetable origin, met with in British India*. K. Paul, Trench, Trübner & Company
- Usmanhani K, Saeed A, Alam MT (1997) *Indusynic medicine : traditional medicine of herbal animal and mineral origin in Pakistan*. Karachi: Dept. of Pharmacognosy, Faculty of Pharmacy, University of Karachi
- Marco JA, Sanz JF, Albiach R (1992) A sesquiterpene ester from *Lactuca-serriola*. *Phytochemistry* 31: 2539–2540. doi: 10.1016/0031-9422(92)83321-O
- Kim DK (2001) Antioxidative components from the aerial parts of *Lactuca scariola* L. *Arch Pharm Res* 24: 427–430. doi: 10.1007/BF02975189
- Ryu T-B, Lim J-C, Lee C-H, Kim E-J, Choi B-K (2017) Distribution of invasive species in metropolitan Busan, South Korea. *J Life Sci* 27: 408–416. doi: 10.5352/JLS.2017.27.4.408
- Kim G (2020) Alien species, american bullfrog (*Lithobates catesbeianus*) and making invasive species in Korea. Dissertation, Seoul National University
- Lee I-Y, Kim S-H, Hong S-H (2021) Occurrence characteristics and management of invasive weeds, *Ambrosia artemisiifolia*, *Ambrosia trifida* and *Humulus japonicus*. *Weed Turf Sci* 10: 227–242
- Lim DO, Hwang IC, Chekar E, Choi DH (2016) A management of invasive alien species and naturalized plant at Jeonju city. *Proceeding of the Korean J Environ Ecol Conference* 2016: 93
- Awan AF, Akhtar MS, Anjum I, Mushtaq MN, Fatima A, Mannan A, Ali I (2020) Anti-oxidant and hepatoprotective effects of *Lactuca serriola* and its phytochemical screening by HPLC and FTIR analysis. *Pak J Pharm Sci* 33: 2823–2830
- Moon S-I, Kim S-W, Huh W, Kim S-Y, Kim J-S, Lee K-J (2009) Isolation and characterization of bio-active materials from prickly lettuce (*Lactuca serriola*). *J Life Sci* 19: 206–212. doi: 10.5352/JLS.2009.19.2.206
- Werk KS, Ehleringer J (1985) Photosynthetic characteristics of *Lactuca serriola* L. *Plant Cell Environ* 8: 345–350. doi: 10.1111/j.1365-3040.1985.tb01409.x
- Guo S, Fang F, Ni L, Chen W, Shi L (2006) Photosynthetic characteristics and coenological survey of *Lactuca serriola* in its invaded area. *Chin J Appl Ecol* 17: 2316–2320
- Alexander JM (2010) Genetic differences in the elevational limits of native and introduced *Lactuca serriola* populations. *J Biogeogr* 37: 1951–1961. doi: 10.1111/j.1365-2699.2010.02335.x
- El-Esawi MA, Elkelish A, Elansary HO, Ali HM, Elshikh M, Witzak J, Ahmad M (2017) Genetic transformation and hairy root induction enhance the antioxidant potential of *Lactuca serriola* L. *Oxid Med Cell* 2017: 5604746. doi: 10.1155/2017/5604746
- Novotna A, Dolezalova I, Lebeda A, Krskova M, Berka T (2011) Morphological variability of achenes of some European populations of *Lactuca serriola* L. *Flora* 206: 473–483. doi: 10.1016/j.flora.2010.09.012
- An L, Guan S, Shi G, Bao Y, Duan Y, Jiang B (2006) Protocatechuic acid from *Alpinia oxyphylla* against MPP<sup>+</sup>-induced neurotoxicity in PC12 cells. *Food Chem Toxicol* 44: 436–443. doi: 10.1016/j.fct.2005.08.017
- Jeong C-H, Jeong HR, Choi GN, Kim D-O, Lee U, Heo HJ (2011) Neuroprotective and anti-oxidant effects of caffeic acid isolated from *Erigeron annuus* leaf. *Chin Med* 6: 1–9. doi: 10.1186/1749-8546-6-25
- Kim MJ, Wang HS, Lee MW (2020) Anti-inflammatory effects of fermented bark of *Acanthopanax sessiliflorus* and its isolated compounds on lipopolysaccharide-treated RAW 264.7 macrophage cells. *Evid-based Complement Altern* 2020. doi: 10.1155/2020/6749425
- Liu W, Nisar MF, Wan C (2020) Characterization of phenolic constituents from *Prunus cerasifera* Ldb leaves. *J Chem* 2020. doi: 10.1155/2020/5976090
- Eshbakova K, Toshmatov Z, Yili A, Aisa H, Abdullaev N (2013) Flavonoid galacturonides and glucuronide from the aerial part of *Scutellaria schachristanica*. *Chem Nat Compd* 49: 103–105. doi: 10.1007/s10600-013-0519-y
- Fajriah S, Megawati M, Darmawan A (2016) Apigenin, an anticancer isolated from *Macaranga gigantifolia* leaves. *J Trop Life Sci* 6: 7–9
- Hu W, Wang X, Wu L, Shen T, Ji L, Zhao X, Si C-L, Jiang Y, Wang G (2016) Apigenin-7-O- $\beta$ -D-glucuronide inhibits LPS-induced inflammation through the inactivation of AP-1 and MAPK signaling pathways in RAW 264.7 macrophages and protects mice against endotoxin shock. *Food Funct* 7: 1002–1013. doi: 10.1039/c5fo01212k
- Lin L-C, Pai Y-F, Tsai T-H (2015) Isolation of luteolin and luteolin-7-O-glucoside from *Dendranthema morifolium* Ramat Tzvel and their pharmacokinetics in rats. *J Agric Food Chem* 63: 7700–7706. doi: 10.1021/jf505848z
- Park C-H, Hur J-M, Song K-S, Park J-C (2007) Phenolic compounds from the leaves of *Nelumbo nucifera* showing DPPH radical scavenging effect. *Korean J Pharmacogn* 38: 263–269
- Van Loo P, De Bruyn A, Buděšínský M (1986) Reinvestigation of the structural assignment of signals in the <sup>1</sup>H and <sup>13</sup>C NMR spectra of the flavone apigenin. *Magn Reson Chem* 24: 879–882. doi: 10.1002/mrc.1260241007
- Delage B (2015) *Flavonoids*. Linus Pauling Institute, Oregon State University, Corvallis, Oregon
- Sedlářová M, Lebeda A (2001) Histochemical detection and role of phenolic compounds in the defense response of *Lactuca* spp. to lettuce downy mildew (*Bremia lactucae*). *J Phytopathol* 149: 693–697. doi: 10.1046/j.1439-0434.2001.00698.x
- Wang SY, Chang HN, Lin KT, Lo CP, Yang NS, Shyur LF (2003) Antioxidant properties and phytochemical characteristics of extracts from *Lactuca indica*. *J Agric Food Chem* 51: 1506–1512. doi: 10.1021/jf0259415
- Nugroho A, Choi JS, An H-J, Park H-J (2015) Phytochemical analysis of the phenolic fat-suppressing substances in the leaves of *Lactuca raddeana* in 3T3-L1 adipocytes. *Nat Prod Sci* 21: 42–48
- Kim MJ, Moon Y, Tou JC, Mou B, Waterland NL (2016) Nutritional value, bioactive compounds and health benefits of lettuce (*Lactuca sativa* L.). *J Food Compos Anal* 49: 19–34. doi: 10.1016/j.jfca.2016.03.004
- Degl'Innocenti E, Pardossi A, Tattini M, Guidi L (2008) Phenolic compounds and antioxidant power in minimally processed salad. *J Food Biochem* 32: 642–653. doi: 10.1111/j.1745-4514.2008.00188.x
- Kim M-H, Nugroho A, Lim S-C, Moon H-E, Choi J-S, Park H-J (2011) Phytochemical analysis of phenolic compounds in 30% ethanolic extracts from the Compositae plants and peroxynitrite-scavenging effect. *Korean J Pharmacogn* 42: 149–154