

# A New Record of *Hadennia nakatanii* (Lepidoptera, Erebidae) from Korea

Bora Shin<sup>1</sup>, Sung-Soo Kim<sup>2</sup>, Sei-Woong Choi<sup>3,\*</sup>

<sup>1</sup>Department of Biology, Mokpo National University, Muan 58554, Korea

<sup>2</sup>Research Institute for East Asian Environment and Biology, Seoul 05264, Korea

<sup>3</sup>Department of Environmental Education, Mokpo National University, Muan 58554, Korea

## ABSTRACT

One species of the Herminiinae, *Hadennia nakatanii* Owada is newly recorded from Korea. *Hadennia nakatanii* can be distinguished by the light blackish forewing with the relatively thick, transverse medial line, a large light yellowish discal dot, the thick blackish slanted band from the apex to the middle part of the dorsum, and the light blackish hindwing, basally suffused with black and medially broadly suffused with black. The female genitalia can be diagnosed by the simple antrum, the long strongly sclerotized ductus bursae, and the large rectangular corpus bursae with a small uplifted signa patch with minute spicules. We provided the figures of adult and the genitalia with short notes on distribution.

**Keywords:** Erebidae, Herminiinae, new record, Korea

## INTRODUCTION

The Herminiinae, a subfamily of the Erebidae, are defined by several morphological characters: the prespiracular counter-tympanal hood, the divided M4 muscle in the male genitalia, the lack of orbicular stigma of the forewing, the presence of male sexual organs throughout the body such as antennae, labial palps, forelegs, and wings, and three SV setae on the larval segments (Owada, 1987; Kitching and Rawlins, 1999; Holloway, 2008). However, Zahiri et al. (2012) found a close relationship between Aganainae and Herminiinae within Erebidae using one mitochondrial and seven nuclear genes, and the monophyly of the Herminiinae was in doubt.

The adults of the Herminiinae are cryptic and the larvae largely feed on dead or withered foliage within the plant (Holloway, 2008). In Korea, about 63 species in 20 genera of the group are recorded (Kim et al., 2016; Lee et al., 2021). The purpose of the present study is to report one species of the Herminiinae, *Hadennia nakatanii* Owada for the first time in Korea.

## MATERIALS AND METHODS

Adult moth was collected at night using a 22-watt UV-light

bucket trap with a 12 V battery (BioQuip, USA). The collected adult was preserved in a freezer and mounted for examination. Species identification was mainly based on the external morphology of adults including the genitalia. For genitalia slide preparation, the specimen was prepared by boiling the abdomen in 10% KOH for approximately 15–20 min. The scales and tissues were removed, stained with Chlorazol black, and mounted on slides in Euparal mountant.

The terminology of the adult characteristics, including the female genitalia, refers to Owada (1987). The examined material has been deposited in the National Institute of Biological Research, Incheon, South Korea. Abbreviations are as follows: TS, type species; TL, type locality; GB, Gyeong-sangbuk-do.

## SYSTEMATIC ACCOUNTS

Order Lepidoptera Linnaeus, 1758

Family Erebidae Leach, 1810

Subfamily Herminiinae Leach, 1815

Genus *Hadennia* Moore, [1885]

Type species: *Bocana hypenalis* Walker, [1859]

= *Wilkara* Swinhoe, 1918 (TS: *Wilkara nigerrima* Swin-

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

**\*To whom correspondence should be addressed**

Tel: 82-61-450-2783, Fax: 82-61-450-2789

E-mail: choisw@mokpo.ac.kr



**Fig. 1.** Adult of *Hadennia nakatanii* in Korea. Scale bar= 10 mm.

hoe, 1918)

<sup>1</sup>\**Hadennia nakatanii* Owada, 1979 (Figs. 1, 2)

*Hadennia nakatanii* Owada, 1979: 131. TL: Japan, Wakayama Pref., Higashi-Muro, Mt. Oto-san, Osugi-dani.

**Material examined.** Korea: 1 female, GB: Mungyung-si, Nongam-myon, 28 Sep 2020, Kim SS.

**Diagnosis.** Wingspan 27 mm. *Hadennia nakatanii* can be distinguished by the long upturned grayish labial palpi, the light blackish forewing with the relatively thick, transverse medial line, a large light yellowish discal dot, and the thick blackish slanted band from the apex to the middle part of the dorsum, and the light blackish hindwing, basally suffused with black and medially broadly suffused with black. *Hadennia nakatanii* is externally similar to *H. incongruens* but can be easily distinguished by the large light yellowish discal dot of the forewing. The female genitalia can be diagnosed by the almost equal length of anterior and posterior apophyses, the simple antrum, the strongly sclerotized ductus bursae with anteriorly narrowed, and the large rectangular corpus bursae, almost twice the length of ductus bursae, with a small uplifted signa patch with minute spicules. The female genitalia of *H. nakatanii* are similar to those of *H. incongruens* but can be distinguished by the long ductus bursae and the medially located small uplifted signa patch of the corpus bursae.

**Distribution.** Korea, Japan, and China (Taiwan).

**Remarks.** The genus is characterized by the wing pattern elements that show the dark brown or black with grayish violet or purple bands on wings, the distally dentate post-medial band and the pale yellow discal dot of the forewing, and the simple valva, the short saccula process with apically rounded, and the finely scobinate vesica of aedeagus (Hol-



**Fig. 2.** Female genitalia of *Hadennia nakatanii* in Korea. Scale bar= 1 mm.

loway, 2008). About 20 species are known worldwide, predominantly occurring in Southeast Asia (Holloway, 2008). In Korea, three species of *Hadennia* are known, including *H. nakatanii*.

## ORCID

Bora Shin: <https://orcid.org/0000-0002-0081-0711>

Sung-Soo Kim: <https://orcid.org/0000-0001-5693-4142>

Sei-Woong Choi: <https://orcid.org/0000-0001-6326-399X>

## CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

## ACKNOWLEDGMENTS

We are grateful to anonymous reviewers for their appreciation of our work. This work was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea

Korean name: <sup>1</sup>\*노란점검은피수염나방 (신칭)

(NIBR202203201, NIBR202231206).

## REFERENCES

- Holloway JD, 2008. The moths of Borneo: Family Noctuidae, subfamilies Rivulinae, Phytometrinae, Herminiinae, Hypeninae and Hypenodinae. *Malayan Nature Journal* 60:1-268.
- Kim SS, Choi SW, Kononenko V, Schintlmeister A, Sohn JC, 2016. Revised list of the Korean Noctuoidea based on latest classification. *Entomological Research Bulletin*, 32:138-160.
- Kitching IJ, Rawlins JE, 1999. The Noctuoidea. In: *Lepidoptera, moths and butterflies* (Ed., Kristensen NP). *Handbuch der Zoologie, Band IV*. Walter de Gruyter, Berlin, pp. 355-401.
- Lee KW, Koo JM, Kim JD, Han HL, Park KT, 2021. A new species of *Zanclognatha* Lederer and three new records of the families Erebiidae and Noctuidae (Lepidoptera, Noctuoidea) from Korea, with notes on larval feeding habits. *Journal of Forestry Research*, 32:1235-1240. <https://doi.org/10.1007/s11676-020-01170-2>
- Owada M, 1979. The Herminiine moths of the genus *Hadennia* (Noctuidae) from Japan, with reference to their distribution in the Kii Peninsula. *Memoirs of the National Science Museum, Tokyo*, 12:123-137.
- Owada M, 1987. A taxonomic study on the subfamily Herminiinae of Japan (Lepidoptera, Noctuidae). *National Science Museum, Tokyo*, pp. 1-208.
- Zahiri R, Holloway JD, Kitching IJ, Lafontaine JD, Mutanen M, Wahlberg N, 2012. Molecular phylogenetics of Erebiidae (Lepidoptera, Noctuoidea). *Systematic Entomology*, 37:102-124. <https://doi.org/10.1111/j.1365-3113.2011.00607.x>

Received September 26, 2022  
Revised October 28, 2022  
Accepted October 28, 2022