

A Pumilio Activity Sensor Reveals Bag-of-Marbles Inhibition of Pum Activity in the *Drosophila* Ovary

Wijeong Jang and [†]Changsoo Kim

School of Biological Sciences and Technology, Chonnam National University, Gwangju 61186, Korea



Received: January 8, 2023

Revised: February 23, 2023

Accepted: March 17, 2023

[†]Corresponding author

Changsoo Kim
School of Biological Sciences and
Technology, Chonnam National University,
Gwangju 61186, Korea
Tel: +82-62-530-5201
E-mail: changgk2001@hanmail.net

Copyright © 2023 The Korean Society of
Developmental Biology.

This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License
([http://creativecommons.org/licenses/
by-nc/4.0/](http://creativecommons.org/licenses/by-nc/4.0/)) which permits unrestricted
non-commercial use, distribution, and
reproduction in any medium, provided
the original work is properly cited.

ORCID

Wijeong Jang
<https://orcid.org/0000-0002-3056-4566>
Changsoo Kim
<https://orcid.org/0000-0002-2852-9649>

Conflict of interests

The authors declare no potential conflict
of interest.

Acknowledgements

This research was supported by
grants from the National University
Development Project 2021-
4065/202233410001 (Changsoo Kim) and
NRF-2020R111A1A01074292 (Wijeong
Jang). We thank Dr. Cohen for the GFP
translational reporter vector, which was
used to make the Pum activity sensor.
We are grateful to Dr. McKearin and Dr.
Wharton for the flies.

Authors' contributions

Conceptualization: Jang W, Kim C.
Data curation: Jang W.
Methodology: Jang W.
Writing-original draft: Jang W.
Writing-review & editing: Jang W, Kim C.

Abstract

Pumilio (Pum) is an RNA-binding protein and translational repressor important to diverse biological processes. In the *Drosophila* ovary, Pum is expressed in female germline stem cells (GSCs), wherein it acts as an intrinsic stem cell maintenance factor via repressing target mRNAs that are as yet mostly unknown. Pum recognizes the Pum binding sequence (PBS) in the mRNA 3'UTR through its C-terminus Puf domain. Translational repression is mediated through its N-terminal domain, but the molecular mechanism remains largely unknown. We previously showed that Bag-of-marbles, a critical differentiation-promoting factor of female GSCs, physically interacts with the N-terminus of Pum. We further showed that this interaction is critical to Bam inhibition of Pum repressive action in cultured cells, but the physiological relevance was not addressed. Here we design an *in vivo* GFP translational reporter bearing the PBS in its 3'UTR and show that GFP expression is reduced in cells wherein Pum is known to be active. Furthermore, we demonstrate in *pum* mutant ovary that this GFP repression requires Pum, and also that the sensor faithfully monitors Pum activity. Finally, we show that forced expression of Bam inhibits Pum-mediated repression, validating that Bam inhibits Pum activity *in vivo*.

Keywords: Pumilio, Bam, Stem cells, Translation repression, *Drosophila melanogaster*

INTRODUCTION

Stem cells possess the unique ability to self-renew in addition to generating differentiated progeny. The *Drosophila* ovary offers an incisive genetic system that allows for dissection of the molecular mechanism of stem cell self-renewal (Xie & Spradling, 2000). Two female germline stem cells (GSCs) reside at the apical tip of the germarium, a structure located at the anterior tip of the ovariole, and are physically linked through adherent junctions to niche cells, called cap cells (Xie & Spradling, 1998). During self-renewal stem cell division, one stem cell daughter attached to the cap cells remains as a stem cell while the other, which is one cell distant from the cap cells, starts to differentiate due to reduction of niche signals (Xie & Spradling, 2000). It becomes a cystoblast (CB), which undergoes incomplete mitotic cell division four times to produce a cyst composed of 16 germline cells. Ultimately, 15 of those cells differentiate to nurse cells and one becomes an oocyte (Lin & Spradling, 1993).

In addition to signals from the cap cells, maintaining the stemness of GSCs also requires stem-cell-specific intrinsic factors (Parisi & Lin, 1999; Wang & Lin, 2004). The RNA binding and translational

Ethics approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

repressor Pum is expressed in GSCs and is required for stemness (Parisi & Lin, 1999; Harris et al., 2011). In *pum* mutant ovary, GSCs lose their stemness and undergo differentiation, depleting the germarium (Lin & Spradling, 1997; Parisi & Lin, 1999). This suggests that Pumilio (Pum)'s role is to block differentiation of stem cells by inhibiting intrinsic differentiation-promoting factors, though the specific factors it targets remain mostly unknown (Szakmary et al., 2005). The C-terminal Puf motif of Pum recognizes the Pum binding sequence (PBS) in the 3'UTR of target mRNAs (Wharton et al., 1998; Edwards et al., 2001; Weidmann et al., 2016; Qiu et al., 2019), while its N-terminal domain mediates translational repression through molecular mechanisms that are as yet poorly understood (Kim et al., 2010; Weidmann & Goldstrohm, 2012; Arvola et al., 2020).

Bag-of-marbles (Bam) is not expressed in GSCs, but is expressed in the 2-, 4-, and 8-cell CBs, wherein Bam promotes differentiation (McKearin & Spradling, 1990; Lavoie et al., 1999). In the *bam* mutant ovary, GSC daughters fail to undergo differentiation, culminating in population of the germaria with stem-cell-like cells (McKearin & Spradling, 1990). Forced ectopic expression of Bam in GSCs promotes their differentiation (Ohlstein & McKearin, 1997; Ohlstein et al., 2000), raising the hypothesis that Bam might inhibit Pum activities. We previously tested this hypothesis *in vitro* and found that Bam physically interacts with the N-terminus of Pum (Kim et al., 2010). We further showed that this physical interaction is essential to inhibition of Pum translational repression activity in cultured cells (Kim et al., 2010). In this work, we design a GFP-based translational reporter bearing the PBS in its 3'UTR and show that this translational reporter is a Pum activity sensor that faithfully monitors Pum activity in the germarium. We furthermore use this sensor to show that Bam inhibits Pum activity *in vivo* in the *Drosophila* ovary.

MATERIALS AND METHODS

1. *Drosophila* stocks

All strains were grown at 25°C on standard yeast, cornmeal agar medium. *pum*¹/TM6 and *pum*^{Msc}/TM6 flies were obtained from Dr. Wharton. *hs-bam* flies were a gift from Dr. McKearin.

2. Pumilio activity sensor

The Pum activity sensor was constructed by introducing the PBS into the p2035 vector (Brennecke & Cohen, 2003). The PBS, which is present in the nanos response element (NRE) (+97 to +148) of the hunchback 3'UTR, was amplified by PCR (using primers ctagaattattttgtgtccaaaattgtacataagccgaattcc, tcgaggaattcggttatgtacattttggacaacaaaataatt) and inserted between the *Xba*I and *Xho*I restriction sites of the p2035 vector. The vector construct was then injected into *w*¹¹¹⁸ embryos to produce the Pum activity sensor transgene.

3. Immunohistochemistry

Ovaries were dissected, fixed, and stained as described previously (Ohlstein & McKearin, 1997). In brief, ovaries were treated for 30 min with fixing solution (4% paraformaldehyde, 0.1% Tween 20). After several washes using PBS with 0.1% Tween 20 (PBST), the samples were preabsorbed for one hour with PBST containing 10% normal goat serum, then incubated with the primary antibody overnight at 4°C. After washing with PBST, the ovaries were incubated with the secondary antibody for two hours at room temperature. After again washing with PBST, the samples were mounted in Vectashield (Vector Laboratories, Road Burlingame, CA, USA). Confocal images were taken and processed by a Zeiss LSM510 Microscope. The GFP level in the area at the apical tip of each germarium was quantified by ImageJ (National Institutes of Health,

Bethesda, MD, USA) and the data were graphed using Sigma Plot.

4. Antibodies

Mouse monoclonal anti-1B1 antibody (1:20) was purchased from Developmental Studies Hybridoma Bank (Iowa City, IA, USA). Rabbit polyclonal anti-GFP (1:2,000) was obtained from Invitrogen (Carlsbad, CA, USA). Goat anti-mouse and goat anti-rabbit IgG conjugated to Alexa 488 or Alexa 555 (1:200) were purchased from Molecular Probe (Eugene, OR, USA).

RESULTS AND DISCUSSION

1. Pumilio activity sensor

We constructed a translational reporter to monitor translational repressive activity of Pum *in vivo* (Fig. 1A), henceforth termed the Pum activity sensor. This reporter consists of the tubulin promoter upstream of the GFP coding sequence which has the PBS in its 3'UTR. The PBS (UGUAXAUA) was taken from the NRE of the *bunchback* 3'UTR (Wharton & Struhl, 1991; Murata & Wharton, 1995; Zamore et al., 1997; Kim et al., 2010). We expected that Pum would bind the PBS in the reporter and repress translation of GFP. If so, GFP would not be expressed in cells with active Pum, but would be expressed in cells lacking Pum activity, thereby enabling the monitoring of Pum activity.

2. The Pumilio activity sensor faithfully monitors Pumilio activity in the germaria

Two cells located at the apical tip in the germaria of the *Drosophila* ovary are GSCs that possess a spectrosome (or round fusome) recognizable by 1B1 antibody (Fig. 1B) (Deng & Lin, 1997). Mitotic cell division of the GSCs produces differentiating germline cysts (consisting of two, four, eight, and 16 cells successively) that are recognized by the presence of a branched spectrosome or fusome, which is also detectable with 1B1 antibody (Lin & Spradling, 1995; Deng & Lin, 1997).

We examined whether the Pum activity sensor faithfully monitors Pum activity in the germaria of the ovary. First, transgenic flies harboring the Pum activity sensor were generated. Ovaries from those transgenic flies were dissected and co-stained with GFP and 1B1 antibodies. In wild-type ovary, GFP expression was severely reduced in the GSCs and in immediate daughter cells derived from their mitotic cell division, which is consistent with reports that Pum is active in the GSCs and immediate daughter cells (Fig. 1C). Meanwhile, GFP was strongly expressed in the germline cyst cells, consistent with reports that Pum is not active in those cells.

To further validate the Pum activity sensor, we examined whether the observed repression of GFP expression requires Pum. Ovaries from transheterozygous mutant (*pum¹/pum^{Msc}*) flies (Wharton et al., 1998) were stained with GFP and 1B1 antibodies. GFP was found to be expressed in all cells (Fig. 1D), demonstrating that its repression requires Pum activity. Combined with the above, these findings confirm that the Pum activity sensor faithfully reports Pum activities in the ovary.

3. Ectopic expression of bag-of-marbles inhibits Pumilio activity as detected by the Pumilio activity sensor

We previously showed that Bam inhibits Pum activity in cultured cells (Kim et al., 2010). Here we examined the *in vivo* relevance of this finding using the Pum activity sensor. To overexpress Bam in all cells including GSCs, we used the *hs-bam* transgene (Ohlstein & McKearin, 1997), which consists of a heat-shock promoter upstream of the *bam* coding sequence and enables ubiquitous expression of *bam* upon a brief heat shock. *hs-bam* flies also harboring the Pum activity sensor were




Fig. 1. The Pum activity sensor faithfully monitors Pum activity in the germaria. (A) Schematic drawing of the Pum activity sensor, which consists of the tubulin (*tub*) promoter upstream of the GFP coding sequence and the Pum binding sequence (PBS) in the 3'UTR. Pum binds the PBS to mediate translational repression. (B) Schematic drawing of the germarium located at the tip of the *Drosophila* ovary. Two germline stem cells distinguished by the spectrosome (red circle) are located at the apical tip of the germarium. The star denotes an immediate stem cell daughter. Germline cysts containing 2, 4, 8, and 16 germline cells are displayed in green with the fusome (branched spectrosome) shown in red. (C,D) Confocal images of germaria bearing the Pum activity sensor in the wild-type background (C) and *pum* mutant background (D); images were co-stained with GFP antibody (green) and 1B1 antibody (red). *Pum* mutant denotes the *pum* transheterozygous mutant (*pum*¹/*pum*^{Msc}). Four-day-old flies were used. White dashed lines encircle cells at the apical tips of germaria. Pum, Pumilio; GFP, green fluorescence protein.

subjected to a brief heat shock (37°C, two hours with one-hour interval) and their ovaries were dissected and co-stained with GFP and 1B1 antibodies at 1, 2, 6, 15, 24, 30, 42, and 48 h post-heat shock (PHS).

Control (no heat shock) flies exhibited no GFP in the GSCs (Fig. 2A). At 2 h PHS, a slight increase of GFP occurred in GSCs at the anterior tip (Fig. 2B, E). At 24 h PHS, peak GFP expression was observed (Fig. 2C, E, and Table 1). At 48 h PHS, GFP expression was severely reduced (Fig. 2D, E, and Table 1). In short, induction of Bam by a brief heat shock induced gradual increase of GFP in the GSCs, followed by a decrease (Fig. 2E and Table 1). This suggests that a gradual increase of Bam in turn gradually inhibits Pum activity in a temporal manner. It is of note that the gradual increase of GFP, which reflects a gradual decrease of Pum activity, is accompanied by a gradual decrease in the number of GSCs at the apical tip of the germaria, which is consistent with Pum activity being required for stem cell maintenance.

4. Conclusions

Here we examine the physiological relevance of previous findings that Bam inhibits Pum activity in cultured cells. We generated an *in vivo* Pum activity sensor consisting of a translational




Fig. 2. Brief expression of Bam inhibits Pum activity as monitored by the Pum activity sensor. (A–D) Confocal images of germlaria bearing the Pum activity sensor in conjunction with either the *hs-gal4* (A) or the *hs-bam* (B–D) transgene were co-stained with GFP (green) and 1B1 (red) antibodies. Germlaria were from three-day-old flies. *hs-gal4* denotes heat-shock promoter-*gal4* coding sequence. *hs-bam* denotes heat-shock promoter-*bam* coding sequence. (A) Germlaria with no heat-shock treatment showed severely reduced GFP level in the GSCs. (B–D) Germlaria treated with a brief heat-shock showed slight increase of GFP at 2 h post-heat-shock (PHS) (B), high expression of GFP at 24 h PHS (C), and less GFP at 48 h PHS (D). (E) Graphical representation of GFP level at the apical tip and GSC number. GFP level at the apical tip of the germlarium (circled areas in Fig. 2A–D) was quantified using ImageJ (NIH); the data are presented in Table 1. GFP, green fluorescence protein; GSCs, germline stem cells.

Table 1. Quantification of GFP level and GSC number

PHS (hour)	GFP level	GSC number	n
0	1.0 (+/- 0.14)	2.5 (+/- 0.5)	13
2	1.1 (+/- 0.14)	1.9 (+/- 0.2)	14
4	1.2 (+/- 0.12)	1.6 (+/- 0.5)	12
6	1.4 (+/- 0.15)	0.8 (+/- 0.7)	15
15	1.7 (+/- 0.3)	0.0 (+/- 0.0)	19
24	2.3 (+/- 0.6)	0.0 (+/- 0.0)	12
30	2.4 (+/- 0.5)	0.0 (+/- 0.0)	13
42	1.7 (+/- 0.3)	1.2 (+/- 0.7)	15
48	1.5 (+/- 0.3)	1.3 (+/- 0.5)	11

GFP levels in the area of the GSCs at the apical tips of germlaria (circled areas in Fig. 2A–D) were quantified using Image J (NIH). GFP level was normalized to the observed in the no heat shock control.

Spectrosome-containing cells at the apical tips of germlaria were counted as GSCs.

PHS denotes post-heat shock. n indicates the number of germlaria examined.

Data are presented as mean (\pm SD).

GFP, green fluorescence protein; GSC, germline stem cell; PHS, post-heat shock.




Fig. 3. Bam inhibition of Pum activity in the *Drosophila* ovary. (A) The N-terminus of Pum mediates translational repression, possibly through interaction with other proteins. (B) Bam physically interacts with the N-terminus of Pum, inhibiting Pum translational repression activity. PBS, Pum binding sequence.

GFP reporter that harbors the PBS in its 3'UTR. The sensor faithfully reported Pum activity in the GSCs of the *Drosophila* ovary; namely, GFP expression was severely reduced in the GSCs, which contain active Pum. In addition, we used the sensor to show that ectopically expressed Bam inhibits Pum activity, which was monitored as increased GFP level. Bam physically interacts with the N-terminus of Pum, which mediates translational repression, possibly through interaction with other proteins (Fig. 3A). Our findings suggest that Bam inhibits Pum activity through blocking such interactions (Fig. 3B).

REFERENCES

- Arvola RM, Chang CT, Buytendorp JP, Levdansky Y, Valkov E, Freddolino PL, Goldstrohm AC (2020) Unique repression domains of Pumilio utilize deadenylation and decapping factors to accelerate destruction of target mRNAs. *Nucleic Acids Res* 48:1843-1871.
- Brennecke J, Cohen SM (2003) Towards a complete description of the microRNA complement of animal genomes. *Genome Biol* 4:228.
- Deng W, Lin H (1997) Spectrosomes and fusomes anchor mitotic spindles during asymmetric

- germ cell divisions and facilitate the formation of a polarized microtubule array for oocyte specification in *Drosophila*. *Dev Biol* 189:79-94.
- Edwards TA, Pyle SE, Wharton RP, Aggarwal AK (2001) Structure of Pumilio reveals similarity between RNA and peptide binding motifs. *Cell* 105:281-289.
- Harris RE, Pargett M, Sutcliffe C, Umulis D, Ashe HL (2011) Brat promotes stem cell differentiation via control of a bistable switch that restricts BMP signaling. *Dev Cell* 20:72-83.
- Kim JY, Lee YC, Kim C (2010) Direct inhibition of Pumilio activity by Bam and Bgcn in *Drosophila* germ line stem cell differentiation. *J Biol Chem* 285:4741-4746.
- Lavoie CA, Ohlstein B, McKearin DM (1999) Localization and function of Bam protein require the benign gonial cell neoplasm gene product. *Dev Biol* 212:405-413.
- Lin H, Spradling AC (1993) Germline stem cell division and egg chamber development in transplanted *Drosophila* germaria. *Dev Biol* 159:140-152.
- Lin H, Spradling AC (1995) Fusome asymmetry and oocyte determination in *Drosophila*. *Dev Genet* 16:6-12.
- Lin H, Spradling AC (1997) A novel group of pumilio mutations affects the asymmetric division of germline stem cells in the *Drosophila* ovary. *Development* 124:2463-2476.
- McKearin DM, Spradling AC (1990) Bag-of-marbles: A *Drosophila* gene required to initiate both male and female gametogenesis. *Genes Dev* 4:2242-2251.
- Murata Y, Wharton RP (1995) Binding of pumilio to maternal hunchback mRNA is required for posterior patterning in *Drosophila* embryos. *Cell* 80:747-756.
- Ohlstein B, Lavoie CA, Vef O, Gateff E, McKearin DM (2000) The *Drosophila* cystoblast differentiation factor, benign gonial cell neoplasm, is related to DExH-box proteins and interacts genetically with bag-of-marbles. *Genetics* 155:1809-1819.
- Ohlstein B, McKearin D (1997) Ectopic expression of the *Drosophila* Bam protein eliminates oogenic germline stem cells. *Development* 124:3651-3662.
- Parisi M, Lin H (1999) The *Drosophila* pumilio gene encodes two functional protein isoforms that play multiple roles in germline development, gonadogenesis, oogenesis and embryogenesis. *Genetics* 153:235-250.
- Qiu C, Dutcher RC, Porter DF, Arava Y, Wickens M, Hall TMT (2019) Distinct RNA-binding modules in a single PUF protein cooperate to determine RNA specificity. *Nucleic Acids Res* 47:8770-8784.
- Szakmary A, Cox DN, Wang Z, Lin H (2005) Regulatory relationship among *piwi*, *pumilio*, and *bag-of-marbles* in *Drosophila* germline stem cell self-renewal and differentiation. *Curr Biol* 15:171-178.
- Wang Z, Lin H (2004) Nanos maintains germline stem cell self-renewal by preventing differentiation. *Science* 303:2016-2019.
- Weidmann CA, Goldstrohm AC (2012) *Drosophila* Pumilio protein contains multiple autonomous repression domains that regulate mRNAs independently of Nanos and brain tumor. *Mol Cell Biol* 32:527-540.
- Weidmann CA, Qiu C, Arvola RM, Lou TF, Killingsworth J, Campbell ZT, Tanaka Hall TM, Goldstrohm AC (2016) *Drosophila* Nanos acts as a molecular clamp that modulates the RNA-binding and repression activities of Pumilio. *eLife* 5:e17096.
- Wharton RP, Sonoda J, Lee T, Patterson M, Murata Y (1998) The Pumilio RNA-binding domain is also a translational regulator. *Mol Cell* 1:863-872.
- Wharton RP, Struhl G (1991) RNA regulatory elements mediate control of *Drosophila* body pattern by the posterior morphogen nanos. *Cell* 67:955-967.
- Xie T, Spradling AC (1998) *Decapentaplegic* is essential for the maintenance and division of

germline stem cells in the *Drosophila* ovary. *Cell* 94:251-260.

Xie T, Spradling AC (2000) A niche maintaining germ line stem cells in the *Drosophila* ovary. *Science* 290:328-330.

Zamore PD, Williamson JR, Lehmann R (1997) The Pumilio protein binds RNA through a conserved domain that defines a new class of RNA-binding proteins. *RNA* 3:1421-1433.