

Nonlinear Functional Analysis and Applications

Vol. 28, No. 2 (2023), pp. 365-382

ISSN: 1229-1595(print), 2466-0973(online)

<https://doi.org/10.22771/nfaa.2023.28.02.03>

<http://nfaa.kyungnam.ac.kr/journal-nfaa>

Copyright © 2023 Kyungnam University Press



EXISTENCE AND STABILITY RESULTS FOR STOCHASTIC FRACTIONAL NEUTRAL DIFFERENTIAL EQUATIONS WITH GAUSSIAN NOISE AND LÉVY NOISE

P. Umamaheswari¹, K. Balachandran², N. Annapoorani³
and Daewook Kim⁴

¹Department of Mathematics, Dr.N.G.P. Arts and Science College,
Coimbatore 641048, India
e-mail: umamaths.tamil@gmail.com

²Department of Mathematics, Bharathiar University,
Coimbatore 641046, India
e-mail: kb.maths.bu@gmail.com

³Department of Mathematics, Bharathiar University,
Coimbatore 641046, India
e-mail: pooranimaths@gmail.com

⁴Department of Mathematics Education, Seowon University,
Cheongju 28674, South Korea
e-mail: kdw@seowon.ac.kr

Abstract. In this paper we prove the existence and uniqueness of solution of stochastic fractional neutral differential equations with Gaussian noise or Lévy noise by using the Picard-Lindelöf successive approximation scheme. Further stability results of nonlinear stochastic fractional dynamical system with Gaussian and Lévy noises are established. Examples are provided to illustrate the theoretical results.

1. INTRODUCTION

Many stochastic systems do not only depend on the present and past states but also involve derivatives with delays. Besides, many practical systems may

⁰Received June 19, 2022. Revised September 4, 2022. Accepted September 7, 2022.

⁰2020 Mathematics Subject Classification: 34A08, 60H10, 34A12, 93D23.

⁰Keywords: Existence, successive approximation, Mittag-Leffler function, stability,

⁰Corresponding author: K. Balachandran(kb.maths.bu@gmail.com).

suffer some jump type stochastic perturbations for example, earthquakes, epidemic and so on. Such problems often arise in the study of two or more simple oscillatory systems with some interconnections between them wherein the connection can be replaced by differential equations involving delays in the highest order derivatives. This becomes the reason to study the neutral differential equations with the addition of Gaussian noise or Lévy noise. Stability of stochastic differential equations is important to avoid possible blow up of solutions. Several authors [4, 5, 6, 8, 11, 12, 14, 19, 20] have studied the existence and stability of neutral stochastic systems.

Recently fractional differential equations are attracted the attention of many researchers due to its applications in many fields [7, 9, 10, 18]. Stochastic fractional differential equations are the natural generalizations of the fractional differential equations. For this equations the delays and in particular neutral delays affect the behavior of the systems. A comprehensive study of fractional neutral differential equations can be found in [1, 15, 16, 17]. The introduction of stochastic nature in these models provide interesting results on the behaviour of the solutions. Boufoussi and Hajji [2] discussed the successive approximation of neutral functional stochastic differential equations with jumps. Chadha and Bora [3] studied the existence, uniqueness and stability of mild solution for stochastic evolution equations with infinite delay and Poisson jumps. Umamaheswari et al. [22] discussed the existence and stability results for fractional stochastic differential equations with Lévy noise.

In this paper, we derive the stability results of stochastic fractional neutral systems driven by Gaussian and Lévy noises. Successive approximation technique is used to prove the existence results. Uniqueness of such equations is also proved under suitable conditions. Examples are provided to support the theory.

2. PRELIMINARIES

In this section we present a few well-known concepts of fractional and stochastic differential equations.

Definition 2.1. ([21]) (Riemann-Liouville fractional integral). The Riemann-Liouville fractional integral operator of order $\alpha > 0$ of a function $f \in L^1(\mathbb{R}^+)$ is defined as

$$I_{0+}^{\alpha} f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s) ds, \quad t > 0, \quad (2.1)$$

where $\Gamma(\cdot)$ is the Euler gamma function.

Definition 2.2. ([21]) (Riemann-Liouville fractional derivative). The Riemann-Liouville fractional derivative of order $\alpha > 0$, $n-1 < \alpha < n$, $n \in \mathbb{N}$, is defined

as

$$D_{0+}^\alpha f(t) = \left(\frac{d}{dt}\right)^n I_{0+}^{n-\alpha} f(t) = \frac{1}{\Gamma(n-\alpha)} \left(\frac{d}{dt}\right)^n \int_0^t (t-s)^{n-\alpha-1} f(s) ds, \quad (2.2)$$

where the function $f(t)$ has absolutely continuous derivatives upto order $(n - 1)$.

Definition 2.3. ([21]) (Caputo fractional derivative). The Caputo fractional derivative of order $\alpha > 0$, $n - 1 < \alpha < n$, $n \in \mathbb{N}$, is defined as

$${}^C D_{0+}^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{n-\alpha-1} f^{(n)}(s) ds, \quad (2.3)$$

where the function $f(t)$ has absolutely continuous derivatives upto order $(n - 1)$.

Definition 2.4. ([13]) (Stochastic Process). A collection $\{X(t) \mid t \geq 0\}$ of random variables is called a stochastic process.

Definition 2.5. ([13]) (Brownian motion). A real-valued stochastic process $W(t)$ is called a Brownian motion or Wiener process if

- (1) $W(0) = 0$ a.s.,
- (2) $W(t) - W(s)$ is $N(0, t - s)$ for all $t \geq s \geq 0$,
- (3) for all times $0 < t_1 < t_2 < \dots < t_n$, the random variables $W(t_1), W(t_2) - W(t_1), \dots, W(t_n) - W(t_{n-1})$ are independent (independent increments).

Definition 2.6. ([22]) (Lévy Process). Let $\{X(t) \mid t \geq 0\}$ be a stochastic process defined on a complete probability space $(\Omega, \mathcal{F}, \mathcal{P})$. Then X is a Lévy process if

- (1) $X(0) = 0$ (a.s),
- (2) X has an independent and stationary increments,
- (3) X is stochastically continuous. That is, for all $a > 0$ and for all $s > 0$ $\lim_{t \rightarrow s} P(|X(t) - X(s)| > a) = 0$.

Lemma 2.7. ([13]) (Chebyshev’s Inequality). If X is a random variable and $1 \leq p < \infty$, then

$$\mathbb{P}(|X| \geq \lambda) \leq \frac{1}{\lambda^p} \mathbb{E}(|X|^p) \text{ for all } \lambda > 0.$$

Lemma 2.8. ([13]) (Borel Cantelli Lemma). If $\{A_k\} \subset \mathcal{F}$ and $\sum_{k=1}^\infty \mathbb{P}(A_k) < \infty$, then

$$\mathbb{P}(\limsup_{k \rightarrow \infty} A_k) = 0.$$

The following theorem is used to prove our main result.

Theorem 2.9. ([13]) (i) *If $\{X_n\}_{n=1}^\infty$ is a submartingale, then*

$$\mathcal{P} \left(\max_{1 \leq k \leq n} X_k \geq \lambda \right) \leq \mathbb{E}(X_n^+),$$

for all $n = 1, 2, \dots$ and $\lambda > 0$.

(ii) *If $\{X_n\}_{n=1}^\infty$ is a martingale and $1 < p < \infty$, then*

$$\mathbb{E} \left(\max_{1 \leq k \leq n} |X_k|^p \right) \leq \left(\frac{p}{p-1} \right)^p \mathbb{E}(|X_n|^p),$$

for all $n = 1, 2, \dots$.

The following lemmas are necessary to obtain the main results. For that, we assume the following hypothesis:

(H): There exist constants $M > 0, a > 0$ such that for $t \geq 0$,

$$|E_{\alpha,\beta}(At^\alpha)| \leq Me^{-at},$$

where $0 < \alpha < 1$ and $\beta = 1, 2, \alpha$ and the Mittag Leffler function of an $n \times n$ matrix A is defined as

$$E_{\alpha,\beta}(At) = \sum_{k=0}^\infty \frac{(At)^k}{\Gamma(\alpha k + \beta)}, \quad (\alpha, \beta > 0).$$

Lemma 2.10. *Assume that the hypothesis (H) holds. Then for any stochastic process $F : [0, \infty) \rightarrow \mathbb{R}^n$ which is strongly measurable with $\int_0^T \mathbb{E}(|F(s)|^2)ds < \infty, 0 < T \leq \infty$, the following inequality holds for $0 < t \leq T$*

$$\mathbb{E} \left| \int_0^t E_{\alpha,\beta}(A(t-s)^\alpha) F(s) ds \right|^2 \leq (M^2/a) \int_0^t \exp(-a(t-s)) \mathbb{E}(|F(s)|^2) ds,$$

where $\alpha \in (1/2, 1)$ and $\beta = 1, 2, \alpha$.

Proof. Assume that the hypothesis (H) holds; that is there exist constants $M > 0, a > 0$ such that for $t \geq 0$

$$|E_{\alpha,\beta}(At^\alpha)| \leq Me^{-at},$$

where $0 < \alpha < 1$ and $\beta = 1, 2, \alpha$. By the Hölder inequality, we obtain, for $0 < t \leq T$,

$$\begin{aligned} & \mathbb{E} \left| \int_0^t E_{\alpha,\beta}(A(t-s)^\alpha) F(s) ds \right|^2 \\ & \leq \mathbb{E} \left(\int_0^t M \exp(-(a/2)(t-s)) \exp(-(a/2)(t-s)) |F(s)| ds \right)^2 \\ & \leq \mathbb{E} \left(\int_0^t M \exp(-(a/2)(t-s)) ds \right)^2 \mathbb{E} \left(\int_0^t \exp(-(a/2)(t-s)) |F(s)| ds \right)^2 \\ & \leq (M^2/a) \int_0^t \exp(-a(t-s)) \mathbb{E}(|F(s)|^2) ds, \end{aligned}$$

which complete the proof of the lemma. □

Lemma 2.11. ([22]) *Assume that the hypothesis (H) holds. Then for any B_t -adapted predictable process $\Phi : [0, \infty) \rightarrow \mathbb{R}^n$ with $\int_0^T \mathbb{E}|\Phi(s)|^2 ds < \infty$, $t \geq 0$, the following inequality holds for $0 < t \leq T$*

$$\mathbb{E} \left| \int_0^t E_{\alpha,\beta}(A(t-s)^\alpha) \Phi(s) dW(s) \right|^2 \leq M^2 \int_0^t \exp(-a(t-s)) \mathbb{E}|\Phi(s)|^2 ds,$$

where $\alpha \in (1/2, 1)$ and $\beta = 1, 2, \alpha$.

3. NONLINEAR EQUATION WITH GAUSSIAN NOISE

Consider the stochastic fractional neutral differential equation with Gaussian noise of the form:

$$\left. \begin{aligned} {}^C D^\alpha(x(t) - p(t, x(t))) &= b(t, x(t)) + \sigma(t, x(t)) \frac{dW(t)}{dt}, \quad t \in J = [0, T], \\ x(0) &= x_0, \end{aligned} \right\} (3.1)$$

where $\alpha \in (1/2, 1)$, $p \in C(J \times \mathbb{R}^n, \mathbb{R}^n)$, $b \in C(J \times \mathbb{R}^n, \mathbb{R}^n)$, $\sigma \in C(J \times \mathbb{R}^n, \mathbb{R}^{n \times m})$ and $W = \{W(t), t \geq 0\}$ is an m -dimensional Brownian motion on a complete probability space $(\Omega, \mathcal{F}, \mathcal{P})$. The integral form of equation (3.1) is given by

$$\begin{aligned} x(t) &= x_0 - p(0, x_0) + p(t, x(t)) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} b(s, x(s)) ds \\ & \quad + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sigma(s, x(s)) dW(s). \end{aligned} \tag{3.2}$$

Assume the following conditions:

(H1) (Linear growth condition) $b : J \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\sigma : J \times \mathbb{R}^n \rightarrow \mathbb{R}^{n \times m}$ and $p : J \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ are continuous and there exists a constant $K > 0$ such that

$$|b(t, x)|^2 + |\sigma(t, x)|^2 + |p(t, x)|^2 \leq K^2(1 + |x|^2)$$

for all $x \in \mathbb{R}^n$.

(H2) (Lipschitz condition) There exists a constant $L > 0$ such that

$$|b(t, x) - b(t, y)|^2 + |\sigma(t, x) - \sigma(t, y)|^2 + |p(t, x) - p(t, y)|^2 \leq L^2(|x - y|^2)$$

for all $x, y \in \mathbb{R}^n$.

Theorem 3.1. *If the hypotheses (H1) – (H2) are satisfied and if x_0 is a random variable defined on $(\Omega, \mathcal{F}, \mathcal{P})$ and independent of the σ -algebra $\mathcal{F}_s^t \subset \mathcal{F}$ generated by $\{W(s), t \geq s \geq 0\}$ such that $\mathbb{E}(|x_0|^2) < \infty$ then the initial value problem (3.1) has a unique solution which is t -continuous with the property that $x(t, \omega)$ is adapted to the filtration $\mathcal{F}_t^{x_0}$ generated by x_0 and $\{W(s)(\cdot), s \leq t\}$ and*

$$\sup_{0 \leq t \leq T} \mathbb{E}[|x(t)|^2] < \infty.$$

Proof. First we prove the existence of solution of the initial value problem. For that, define $x^{(0)}(t) = x_0$ and $x^{(k)}(t) = x^{(k)}(t, \omega)$ inductively by

$$\begin{aligned} x^{(k+1)}(t) = & x_0 - p(0, x_0) + p(t, x^{(k)}(s)) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} b(s, x^{(k)}(s)) ds \\ & + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sigma(s, x^{(k)}(s)) dW(s), \end{aligned} \tag{3.3}$$

for $k = 0, 1, 2, \dots$. If, for fixed $k \geq 0$, the approximation $x^{(k)}(t)$ is \mathcal{F}_t -measurable and continuous on J , then it follows from (H1)-(H2) that the integrals in (3.3) are meaningful and that the resulting process $x^{(k+1)}(t)$ is \mathcal{F}_t -measurable and continuous on J . As $x^{(0)}(t)$ is obviously \mathcal{F}_t -measurable and continuous on J , it follows by induction that so too is each $x^{(k)}(t)$ for $k = 1, 2, \dots$. It is clear that

$$\sup_{0 \leq t \leq T} \mathbb{E}(|x^{(0)}(t)|^2) < \infty.$$

Since x_0 is \mathcal{F}_t -measurable with $\mathbb{E}(|x_0|^2) < \infty$, by using the algebraic inequality, the Cauchy-Schwarz inequality, the Itô isometry and the linear growth

condition, we obtain from (3.3)

$$\begin{aligned} \mathbb{E}(|x^{(k+1)}(t)|^2) &\leq 4\mathbb{E}[|x_0|^2 + |p(0, x_0)|^2] + 4\mathbb{E}[|p(t, x^{(k)}(s))|^2] \\ &\quad + \frac{4}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \mathbb{E} \left[\int_0^t |b(s, x^{(k)}(s))|^2 ds \right] \\ &\quad + \frac{4}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \mathbb{E} \left[\int_0^t |\sigma(s, x^{(k)}(s))|^2 ds \right]. \end{aligned}$$

Therefore

$$\begin{aligned} \mathbb{E}(|x^{(k+1)}(t)|^2) &\leq 4(1 + K^2)\mathbb{E}[|x_0|^2] + 4K^2\mathbb{E}(1 + |x^{(k)}(t)|^2) \\ &\quad + 2K^2 \frac{4}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \mathbb{E} \left(\int_0^t (1 + |x^{(k)}(s)|^2) ds \right), \end{aligned}$$

for $k = 0, 1, 2, \dots$. By induction, we have

$$\sup_{0 \leq t \leq T} \mathbb{E}(|x^{(k)}(t)|^2) \leq C_5 < \infty,$$

for $k = 1, 2, 3, \dots$ and $C_5 > 0$. Let

$$d^{(k)}(t) = \mathbb{E}(|x^{(k+1)}(t) - x^{(k)}(t)|).$$

We claim that

$$d^{(k)}(t) \leq \frac{(Mt)^{(k+1)}}{(k+1)!}, \text{ for all } k = 0, 1, 2, \dots,$$

for some constant M depending on K, L and x_0 . From equation (3.3) by applying the Schwarz inequality, Itô isometry and the Lipchitz condition we obtain

$$\begin{aligned} d^{(k)}(t) &= \mathbb{E}[|x^{(k+1)}(t) - x^{(k)}(t)|^2] \\ &\leq 3\mathbb{E}(|p(s, x^{(k)}(s)) - p(s, x^{(k-1)}(s))|^2) \\ &\quad + \frac{3}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \int_0^t \mathbb{E} \left[|b(s, x^{(k)}(s)) - b(s, x^{(k-1)}(s))|^2 \right] ds \\ &\quad + \frac{3}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \int_0^t \mathbb{E} \left[|\sigma(s, x^{(k)}(s)) - \sigma(s, x^{(k-1)}(s))|^2 \right] ds \\ &\leq 3L^2 \mathbb{E} \left[|x^{(k)}(t) - x^{(k-1)}(t)|^2 \right] \\ &\quad + 3 \frac{L^2}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \int_0^t \mathbb{E} \left[|x^{(k)}(s) - x^{(k-1)}(s)|^2 \right] ds \\ &\quad + 3 \frac{L^2}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \int_0^t \mathbb{E} \left[|x^{(k)}(s) - x^{(k-1)}(s)|^2 \right] ds. \end{aligned} \tag{3.4}$$

By applying again the Schwarz inequality, the Itô isometry together with the growth conditions for $k = 0$, one gets

$$\begin{aligned}
 d^{(0)}(t) &= \mathbb{E}[|x^{(1)}(t) - x^{(0)}(t)|^2] \\
 &\leq 3\mathbb{E}(|p(t, x_0) - p(0, x_0)|^2) + \frac{3}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \int_0^t \mathbb{E} \left[|b(s, x^{(0)}(s))|^2 \right] ds \\
 &\quad + \frac{3}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \int_0^t \mathbb{E} \left[|\sigma(s, x^{(0)}(s))|^2 \right] ds \\
 &\leq 3L^2t + 2K^2 \frac{3}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \mathbb{E} \left(\int_0^t (1 + |x_0|^2) ds \right) \\
 &\leq 3L^2t + 2K^2 \frac{3}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} (t)(1 + \mathbb{E}(|x_0|^2)). \tag{3.5}
 \end{aligned}$$

Now, for $k = 1$, replacing $\mathbb{E}[|x^{(1)}(t) - x^{(0)}(t)|^2]$ in the inequality (3.4) with the value on the right hand side of inequality (3.5) and integrating, we obtain

$$\begin{aligned}
 \mathbb{E}[|x^{(2)}(t) - x^{(1)}(t)|^2] &\leq 3L^2\mathbb{E}(|x^{(1)}(t) - x^{(0)}(t)|^2) \\
 &\quad + 2L^2 \frac{3}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \int_0^t \mathbb{E}[|x^{(1)}(s) - x^{(0)}(s)|^2] ds \\
 &\leq 3L^2 [K_1 + K^2(1 + \mathbb{E}(|x_0|^2))] \left(\frac{2}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \right)^2 \int_0^t s ds \\
 &\leq 3L^2 [K_1 + K^2(1 + \mathbb{E}(|x_0|^2))] \left(\frac{2}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \right)^2 \times \frac{t^2}{2!},
 \end{aligned}$$

where $K_1 < \infty$. For $k = 2$, proceeding as before, we have

$$\mathbb{E}[|x^{(3)}(t) - x^{(2)}(t)|^2] \leq 3L^2 [K_1 + K^2(1 + \mathbb{E}(|x_0|^2))] \left(\frac{2}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \right)^3 \times \frac{t^3}{3!}.$$

Thus by the principle of mathematical induction, we have

$$d^{(k)}(t) = \mathbb{E}[|x^{(k+1)}(t) - x^{(k)}(t)|^2] \leq \frac{BM^{k+1}t^{(k+1)}}{(k+1)!}, \quad k = 0, 1, 2, \dots, 0 \leq t \leq T,$$

where $B = 3L^2 [K_1 + K^2(1 + \mathbb{E}(|x_0|^2))]$ and $M = \left(\frac{2}{(\Gamma(\alpha))^2} \frac{T^{2\alpha-1}}{2\alpha-1} \right)$ are constants depending only on α, T, L^2, K^2 and $\mathbb{E}(|x_0|^2)$.

Note that

$$\begin{aligned} & \mathbb{E} \left(\max_{0 \leq t \leq T} |x^{(k+1)}(t) - x^{(k)}(t)|^2 \right) \\ & \leq 3L^2 \left(\max_{0 \leq t \leq T} |x^{(k)}(t) - x^{(k-1)}(t)|^2 \right) \\ & \quad + 3L^2 \frac{T^{2\alpha-1}}{2\alpha-1} \mathbb{E} \left(\max_{0 \leq t \leq T} \int_0^t |x^{(k)}(s) - x^{(k-1)}(s)|^2 ds \right) \\ & \quad + 3\mathbb{E} \left(\max_{0 \leq t \leq T} \int_0^t (t-s)^{\alpha-1} |\sigma(s, x^{(k)}(s)) - \sigma(s, x^{(k-1)}(s))|^2 dW(s) \right). \end{aligned}$$

Use of second part of the Theorem 2.9 gives

$$\begin{aligned} & \mathbb{E} \left(\max_{0 \leq t \leq T} |x^{(k+1)}(t) - x^{(k)}(t)|^2 \right) \\ & \leq 3L^2 \left(\max_{0 \leq t \leq T} |x^{(k)}(t) - x^{(k-1)}(t)|^2 \right) \\ & \quad + 3L^2 \frac{T^{2\alpha-1}}{2\alpha-1} \mathbb{E} \left(\int_0^T |x^{(k)}(s) - x^{(k-1)}(s)|^2 ds \right) \\ & \quad + 12L^2 \frac{T^{2\alpha-1}}{2\alpha-1} \mathbb{E} \left(\int_0^T |x^{(k)}(s) - x^{(k-1)}(s)|^2 ds \right) \\ & \leq B \frac{M^{k+1}}{(k+1)!} T^{(k+1)}. \end{aligned} \tag{3.6}$$

Use of Chebyshev’s inequality gives

$$\mathcal{P} \left(\max_{0 \leq t \leq T} |x^{(k+1)}(t) - x^{(k)}(t)|^2 > \frac{1}{k^2} \right) \leq \frac{1}{(1/k^2)^2} \mathbb{E} \left(\max_{0 \leq t \leq T} |x^{(k+1)}(t) - x^{(k)}(t)|^2 \right).$$

Using the equation (3.6) and summing up the resultant inequalities, one gets

$$\sum_{k=0}^{\infty} \mathcal{P} \left(\max_{0 \leq t \leq T} |x^{(k+1)}(t) - x^{(k)}(t)|^2 > \frac{1}{k^2} \right) \leq \sum_{k=0}^{\infty} \frac{BM^{k+1}k^4T^{(k+1)}}{(k+1)!},$$

where the series on the right side converges by ratio test. Hence the series on the left side also converges, so by the Borel-Cantelli lemma, conclude that $\max_{0 \leq t \leq T} (|x^{(k+1)}(t) - x^{(k)}(t)|^2)$ converges to 0, almost surely, that is, the successive approximations $\{x^{(k)}(t)\}$ converge, almost surely, uniformly on J to a limit $x(t)$ defined by

$$\lim_{n \rightarrow \infty} \left(x^{(0)}(t) + \sum_{k=1}^n (x^{(k)}(t) - x^{(k-1)}(t)) \right) = \lim_{n \rightarrow \infty} x^{(n)}(t) = x(t). \tag{3.7}$$

From (3.3), we have

$$\begin{aligned}
 x(t) = & x_0 - p(0, x_0) + p(t, x(t)) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} b(s, x(s)) ds \\
 & + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sigma(s, x(s)) dW(s), \tag{3.8}
 \end{aligned}$$

for all $t \in J$. This completes the proof of the existence of solution of (3.1).

Next, we have to prove the uniqueness of solution. The uniqueness follows from the Itô isometry and the Lipschitz conditions with $L < 1/2$.

Let $x(t, \omega)$ and $y(t, \omega)$ be solution processes through the initial data $(0, x_0)$ and $(0, y_0)$, respectively, that is, $x(0, \omega) = x_0(\omega)$ and $y(0, \omega) = y_0(\omega)$, $\omega \in \Omega$. Let

$$\begin{aligned}
 a(s, \omega) &= b(s, x(s)) - b(s, y(s)), \\
 \gamma(s, \omega) &= \sigma(s, x(s)) - \sigma(s, y(s)).
 \end{aligned}$$

Then, by virtue of the Schwarz inequality and the Itô isometry, we have

$$\begin{aligned}
 \mathbb{E}[|x(t) - y(t)|^2] &\leq 4L_1^2 \mathbb{E}[|x_0 - p(0, x_0) - y_0 + p(0, y_0)|^2] \\
 &\quad + \frac{4L_1^2}{(\Gamma(\alpha))^2} \frac{t^{2\alpha-1}}{2\alpha-1} \mathbb{E} \left[\int_0^t |a(s, \omega)|^2 ds \right] \\
 &\quad + \frac{4L_1^2}{(\Gamma(\alpha))^2} \frac{t^{2\alpha-1}}{2\alpha-1} \mathbb{E} \left[\int_0^t |\gamma(s, \omega)|^2 ds \right] \\
 &\leq 4L_1^2 \mathbb{E}[|x_0 - p(0, x_0) - y_0 + p(0, y_0)|^2] \\
 &\quad + 2L_1^2 \frac{4L^2}{(\Gamma(\alpha))^2} \frac{t^{2\alpha-1}}{2\alpha-1} \int_0^t \mathbb{E}[|x(s) - y(s)|^2] ds,
 \end{aligned}$$

with $L_1^2 = \frac{1}{1-4L^2}$ and $L < 1/2$. We define $v(t) = \mathbb{E}[|x(t) - y(t)|^2]$. Then the function v satisfies

$$v(t) \leq F + A \int_0^t v(s) ds,$$

where $F = 4L_1^2 \mathbb{E}[|x_0 - p(0, x_0) - y_0 + p(0, y_0)|^2]$ and $A = 2L_1^2 \frac{4L^2}{(\Gamma(\alpha))^2} \frac{t^{2\alpha-1}}{2\alpha-1}$.

By the application of the Gronwall's inequality, we conclude that

$$v(t) \leq F \exp(At).$$

Now assume that $x_0 = y_0$. Then $F = 0$ and so $v(t) = 0$ for all $t \geq 0$. That is,

$$\mathbb{E}[|x(t) - y(t)|^2] = 0,$$

which gives

$$\int_{\Omega} |x(t) - y(t)|^2 d\mathcal{P} = 0.$$

This implies that $x(t) = y(t)$ a.s for all $t \in J$. That is,

$$\mathcal{P}\{|x(t, \omega) - y(t, \omega)| = 0 \text{ for all } t \in J\} = 1,$$

that is, the solution is unique. This completes the proof of existence and uniqueness of solution of the given stochastic fractional neutral differential equation (3.1). \square

4. STABILITY OF SYSTEM WITH GAUSSIAN NOISE

In this section, we study the exponential asymptotic stability in the quadratic mean of a trivial solution. Consider the stochastic fractional neutral differential equation with Gaussian noise of the form

$$\left. \begin{aligned} {}^C D^\alpha(x(t) - p(t, x(t))) &= Ax(t) + f(t, x(t)) + \sigma(t, x(t)) \frac{dW(t)}{dt}, \\ x(0) &= x_0, \end{aligned} \right\} \quad (4.1)$$

where $t \in J = [0, T]$, $\alpha \in (1/2, 1)$, $b(t, x(t)) = Ax(t) + f(t, x(t))$, $p \in C(J \times \mathbb{R}^n, \mathbb{R}^n)$, $f \in C(J \times \mathbb{R}^n, \mathbb{R}^n)$, $\sigma \in C(J \times \mathbb{R}^n, \mathbb{R}^{n \times m})$ and $W = \{W(t), t \geq 0\}$ is an m -dimensional Brownian motion on a complete probability space $(\Omega, \mathcal{F}, \mathcal{P})$. $A \in \mathbb{R}^{n \times n}$ is a diagonal stability matrix. Assume from now on that $b(t, 0) = \sigma(t, 0) = p(t, 0) \equiv 0$ a.e t so that equation (4.1) admits a trivial solution.

Definition 4.1. ([13]) The trivial solution of equation (4.1) is said to be exponentially stable in the quadratic mean if there exist positive constants C, ν such that

$$\mathbb{E}(|x(t)|^2) \leq C\mathbb{E}(|x(0)|^2) \exp(-\nu t), \quad t \geq 0.$$

Theorem 4.2. *Let the assumptions of Theorem 3.1 hold. Then the solution of equation (4.1) is exponentially stable in the quadratic mean provided*

$$a > \beta = \beta(a, K, M_1) = \left(5M_1^2(K^2/a(\Gamma(\alpha))^2T + (K^2/a + K^2) \frac{T^{2\alpha-1}}{2\alpha-1} \right).$$

Proof. The integral form of the equation (4.1) is

$$\begin{aligned} x(t) &= E_\alpha(At^\alpha)[x_0 - p(0, x_0)] + p(t, x(t)) \\ &\quad + \int_0^t A(t-s)^{\alpha-1} E_{\alpha,\alpha}(A(t-s)^\alpha) p(s, x(s)) ds \\ &\quad + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}(A(t-s)^\alpha) f(s, x(s)) ds \\ &\quad + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}(A(t-s)^\alpha) \sigma(s, x(s)) dW(s). \end{aligned} \quad (4.2)$$

By using algebraic inequality and Lemmas 2.10 and 2.11, we get

$$\begin{aligned} \mathbb{E}[|x(t)|^2] &\leq 10M^2 \exp(-at)(1 + K^2)\mathbb{E}(|x_0|^2) + 5\mathbb{E}(|p(t, x(t))|^2) \\ &\quad + 5(\Gamma(\alpha))^2 TM^2/a \int_0^t \exp(-a(t-s))\mathbb{E}(|p(s, x(s))|^2)ds \\ &\quad + 5(M^2/a) \frac{T^{2\alpha-1}}{2\alpha-1} \int_0^t \exp(-a(t-s))\mathbb{E}|f(s, x(s))|^2 ds \\ &\quad + 5M^2 \frac{T^{2\alpha-1}}{2\alpha-1} \int_0^t \exp(-a(t-s))\mathbb{E}|\sigma(s, x(s))|^2 ds. \end{aligned}$$

Linear growth assumption when $f(t, 0) = \sigma(t, 0) = p(t, 0) \equiv 0$ a.e t yields

$$\begin{aligned} \exp(at)\mathbb{E}|x(t)|^2 &\leq 10M_1^2(1 + K^2)\mathbb{E}(|x_0|^2) \\ &\quad + 5M_1^2 \left[K^2/a(\Gamma(\alpha))^2 T + (K^2/a + K^2) \frac{T^{2\alpha-1}}{2\alpha-1} \right] \\ &\quad \times \int_0^t \exp(as)\mathbb{E}(|x(s)|^2)ds \end{aligned}$$

with $M_1 = M^2/(1 - 5K^2)$ and $K^2 < 1/5$. Applying Gronwall’s inequality, we obtain

$$\begin{aligned} \exp(at)\mathbb{E}|x(t)|^2 &\leq 10M_1^2(1 + K^2)\mathbb{E}|x_0|^2 \\ &\quad \times \exp \left(5M_1^2(K^2/a(\Gamma(\alpha))^2 T + (K^2/a + K^2) \frac{T^{2\alpha-1}}{2\alpha-1} t) \right). \end{aligned}$$

Consequently

$$\mathbb{E}|x(t)|^2 \leq C\mathbb{E}|x_0|^2 \exp(-\nu t), \quad t \geq 0,$$

where $\nu = a - \beta$ and $C = 10M_1^2(1 + K^2)$. □

5. NONLINEAR EQUATION WITH LÉVY NOISE

Consider the stochastic fractional neutral differential equation with Lévy noise of the form:

$$\begin{aligned} {}^C D^\alpha(x(t) - p(t, x(t))) &= b(t, x(t)) + \sigma(t, x(t)) \frac{dW(t)}{dt} + \int_z g(t, x(t), z) \frac{d\tilde{N}(t, z)}{dt}, \\ x(0) &= x_0, \end{aligned} \tag{5.1}$$

where $t \in J$, $\alpha \in (1/2, 1)$ and $z \in \mathbb{R}_0^n = \mathbb{R}^n/\{0\}$. Here $b, p : J \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\sigma : J \times \mathbb{R}^n \rightarrow \mathbb{R}^{n \times m}$, $g : J \times \mathbb{R}^n \times \mathbb{R}_0^n \rightarrow \mathbb{R}^{n \times l}$ are given functions such that $b(t, x(t))$, $\sigma(t, x(t))$, $p(t, x(t))$ and $g(t, x(t), z)$ are \mathcal{F}_t measurable for all $x \in \mathbb{R}^n$ and $z \in \mathbb{R}_0^n$, $\tilde{N}(dt, dz) = N(dt, dz) - \nu(dz)dt$ which is the l -dimensional compensated jump measure of $\eta(\cdot)$ an independent compensated Poisson random

measure and $W = \{W(t), t \geq 0\}$ is an m -dimensional Brownian motion on a complete probability space $\Omega \equiv (\Omega, \mathcal{F}, \mathcal{P})$. The integral form of the equation (5.1) is

$$\begin{aligned} x(t) = & x_0 - p(0, x_0) + p(t, x(t)) + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} b(s, x(s)) ds \\ & + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sigma(s, x(s)) dW(s) \\ & + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \int_z g(s, x(s), z) d\tilde{N}(s, z). \end{aligned} \tag{5.2}$$

Assume that the following conditions hold true:

(H4) $b : J \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, $p : J \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $\sigma : J \times \mathbb{R}^n \rightarrow \mathbb{R}^{n \times m}$ are continuous and there exists constant $K_2 > 0$ such that

$$|b(t, x)|^2 + |\sigma(t, x)|^2 + |p(t, x)|^2 + \int_z |g(t, x, z)|^2 \nu(dz) \leq K_2^2 (1 + |x|^2)$$

for all $x \in \mathbb{R}^n$.

(H5) There exists a constant $L_2 > 0$ such that

$$\begin{aligned} |b(t, x) - b(t, y)|^2 + |\sigma(t, x) - \sigma(t, y)|^2 + \int_z |g(t, x, z) - g(t, y, z)|^2 \nu(dz) \\ + |p(t, x) - p(t, y)|^2 \leq L_2^2 (|x - y|^2) \end{aligned}$$

for all $x, y \in \mathbb{R}^n$.

Theorem 5.1. *If the hypotheses (H4) – (H5) are satisfied and if x_0 is a random variable defined on $(\Omega, \mathcal{F}, \mathcal{P})$ and independent of the σ -algebra $\mathcal{F}_s^t \subset \mathcal{F}$ generated by $\{W(s), t \geq s \geq 0\}$ such that $\mathbb{E}(|x_0|^2) < \infty$. Then the initial value problem (5.1) has a unique solution which is t -continuous with the property that $x(t, \omega)$ is adapted to the filtration $\mathcal{F}_t^{x_0}$ generated by x_0 and $\{W(s)(\cdot), s \leq t\}$ and*

$$\sup_{0 \leq t \leq T} \mathbb{E}[|x(t)|^2] < \infty.$$

Proof. To prove uniqueness, let $x(t, \omega)$ and $y(t, \omega)$ be solution processes through the initial data $(0, x_0)$ and $(0, y_0)$, respectively, that is, $x(0, \omega) = x_0(\omega)$ and $y(0, \omega) = y_0(\omega)$, $\omega \in \Omega$. Let

$$\begin{aligned} \gamma_1(s, \omega) &= b(s, x(s)) - b(s, y(s)), \\ \gamma_2(s, \omega) &= \sigma(s, x(s)) - \sigma(s, y(s)), \\ \gamma_3(s, \omega) &= \int_z g(s, x(s), z) \nu(dz) - \int_z g(s, y(s), z) \nu(dz). \end{aligned}$$

By the use of Schwarz inequality and the Itô isometry, we have

$$\begin{aligned} \mathbb{E}[|x(t) - y(t)|^2] &\leq 5L_3^2 \mathbb{E}[|x_0 - p(0, x_0) - y_0 + p(0, y_0)|^2] \\ &\quad + \frac{5L_3^2}{(\Gamma(\alpha))^2} \frac{t^{2\alpha-1}}{2\alpha-1} \mathbb{E}\left[\int_0^t |\gamma_1(s, \omega)|^2 ds\right] \\ &\quad + \frac{5L_3^2}{(\Gamma(\alpha))^2} \frac{t^{2\alpha-1}}{2\alpha-1} \mathbb{E}\left[\int_0^t |\gamma_2(s, \omega)|^2 ds\right] \\ &\quad + \frac{5L_3^2}{(\Gamma(\alpha))^2} \frac{t^{2\alpha-1}}{2\alpha-1} \mathbb{E}\left[\int_0^t |\gamma_3(s, \omega)|^2 ds\right] \\ &\leq \frac{5L_3^2}{(\Gamma(\alpha))^2} \mathbb{E}[|x_0 - y_0|^2] \\ &\quad + 5L_2^2 \frac{3L_3^2}{(\Gamma(\alpha))^2} \frac{t^{2\alpha-1}}{2\alpha-1} \int_0^t \mathbb{E}[|x(s) - y(s)|^2] ds, \end{aligned}$$

with $L_3^2 = \frac{1}{1-5L_2^2}$ and $L_2^2 < 1/5$. We define $v(t) = \mathbb{E}[|x(t) - y(t)|^2]$. Then the function v satisfies $v(t) \leq F + A \int_0^t v(s) ds$, where $F = 5L_1^2 \mathbb{E}[|x_0 - p(0, x_0) - y_0 + p(0, y_0)|^2]$ and $A = 5L_2^2 \frac{3L_3^2}{(\Gamma(\alpha))^2} \frac{t^{2\alpha-1}}{2\alpha-1}$. By the application of Gronwall's inequality, we conclude that

$$v(t) \leq F \exp(At).$$

Now assume that $x_0 = y_0$. Then $F = 0$ and so $v(t) = 0$ for all $t \geq 0$. That is,

$$\mathbb{E}[|x(t) - y(t)|^2] = 0,$$

which gives

$$\int_{\Omega} |x(t) - y(t)|^2 d\mathcal{P} = 0.$$

This implies that $x(t) = y(t)$ a.s for all $t \in J$. That is,

$$\mathcal{P}\{|x(t, \omega) - y(t, \omega)| = 0 \text{ for all } t \in J\} = 1,$$

that is, the solution is unique.

Existence of solution of (5.1) is proved as before using Lipschitz condition, growth condition and Cauchy Schwarz inequality by approximation. \square

6. STABILITY OF SYSTEM WITH LÉVY NOISE

In this section we study the exponentially asymptotic stability in the quadratic mean of a trivial solution. Consider the following stochastic fractional

nonlinear system with Lévy noise of the form

$$\begin{aligned}
 {}^C D^\alpha(x(t) - p(t, x(t))) &= Ax(t) + f(t, x(t)) + \sigma(t, x(t)) \frac{dW(t)}{dt} \\
 &\quad + \int_z g(t, x(t), z) \frac{d\tilde{N}(t, z)}{dt}, \quad t \in J, \\
 x(0) &= x_0,
 \end{aligned} \tag{6.1}$$

where $\alpha \in (1/2, 1)$, $z \in \mathbb{R}_0^n = \mathbb{R}^n / \{0\}$, $b(t, x(t)) = Ax(t) + f(t, x(t))$, $f, p \in C(J \times \mathbb{R}^n, \mathbb{R}^n)$, $\sigma \in C(J \times \mathbb{R}^n, \mathbb{R}^{n \times m})$, $g \in C(J \times \mathbb{R}^n \times \mathbb{R}_0^n, \mathbb{R}^{n \times l})$, $\tilde{N}(dt, dz)$ is as in (5.1), $A \in \mathbb{R}^{n \times n}$ is a diagonal stability matrix. Assume from now on that $b(t, 0) = \sigma(t, 0) = p(t, 0) \equiv 0$ a.e t so that equation (6.1) admits a trivial solution.

Definition 6.1. ([13]) The trivial solution of equation (6.1) is said to be exponentially stable in the quadratic mean if there exist positive constants C, ν such that

$$\mathbb{E}(|x(t)|^2) \leq C\mathbb{E}(|x_0|^2) \exp(-\nu t), \quad t \geq 0.$$

Theorem 6.2. *Let the assumptions of Theorem 5.1 hold. Then the solution of equation (6.1) is exponentially stable in the quadratic mean provided*

$$a > \beta = \beta(a, K, M_1) = 6M_1^2 \left[K^2/a(\Gamma(\alpha))^2 T + (2K^2/a + K^2) \frac{T^{2\alpha-1}}{2\alpha - 1} \right].$$

Proof. The integral form of the equation (6.1) is

$$\begin{aligned}
 x(t) &= E_\alpha(At^\alpha)[x_0 - p(0, x_0)] + p(t, x(t)) \\
 &\quad + \int_0^t A(t-s)^{\alpha-1} E_{\alpha,\alpha}(A(t-s)^\alpha) p(s, x(s)) ds \\
 &\quad + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}(A(t-s)^\alpha) f(s, x(s)) ds \\
 &\quad + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}(A(t-s)^\alpha) \sigma(s, x(s)) dW(s) \\
 &\quad + \int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}(A(t-s)^\alpha) \int_z g(s, x(s), z) \tilde{N}(ds, dz). \tag{6.2}
 \end{aligned}$$

Applying assumption (H) when $f(t, 0) = \sigma(t, 0) = g(t, 0, z) = p(t, 0) \equiv 0$ a.e t yields

$$\begin{aligned} \exp(at)\mathbb{E}(|x(t)|^2) &\leq 12M_1^2(1 + K^2)\mathbb{E}(|x_0|^2) \\ &\quad + 6(M_1^2/a)K^2(\Gamma(\alpha))^2T \int_0^t \exp(as)\mathbb{E}(|x(s)|^2)ds \\ &\quad + 6M_1^2(2K^2/a + K^2)\frac{T^{2\alpha-1}}{2\alpha - 1} \int_0^t \exp(as)\mathbb{E}(|x(s)|^2)ds \\ &\leq 12M_1^2(1 + K^2)\mathbb{E}|x_0|^2 + M_2 \int_0^t \exp(as)\mathbb{E}(|x(s)|^2)ds, \end{aligned}$$

where $M_1 = M/(1 - 6K^2)$ with $K^2 < 1/6$ and $M_2 = 6M_1^2 \left[K^2/a(\Gamma(\alpha))^2T + (2K^2/a + K^2)\frac{T^{2\alpha-1}}{2\alpha - 1} \right]$. Applying Gronwall's inequality, we obtain

$$\exp(at)\mathbb{E}(|x(t)|^2) \leq 12M_1^2(1 + K^2)\mathbb{E}(|x_0|^2) \exp(M_2). \tag{6.3}$$

Consequently,

$$\mathbb{E}(|x(t)|^2) \leq C\mathbb{E}(|x_0|^2) \exp(-\nu t), \quad t \geq 0, \tag{6.4}$$

where $\nu = a - \beta$ and $C = 12M_1^2(1 + K^2)$. □

7. EXAMPLES

In this section, we provide two examples to support the theory developed in the previous sections.

Example 7.1. Consider the following stochastic fractional neutral differential equation of the form

$$\left. \begin{aligned} {}^C D^{0.7} \left(x(t) - \frac{1}{5\sqrt{2}} \cos x \right) &= -0.2x(t) + \frac{1}{5\sqrt{2}} \sin x + \frac{1}{5\sqrt{2}} x(t) \frac{dW(t)}{dt}, \\ x(0) &= 5, \quad t \in J. \end{aligned} \right\} \tag{7.1}$$

Here $\alpha = 0.7$, $b(t, x(t)) = 0.2x(t) + \frac{1}{5\sqrt{2}} \sin x$, $p(t, x(t)) = \frac{1}{5\sqrt{2}} \cos x$ and $\sigma(t, x(t)) = \frac{1}{5\sqrt{2}} x(t)$. It can be easily seen that $b(t, x(t))$, $p(t, x(t))$ and $\sigma(t, x(t))$ satisfy the assumptions of Theorem 3.1. Hence, by Theorem 3.1, the stochastic fractional differential equation (7.1) has a unique solution. Also the equation (7.1) satisfies the condition of Theorem 4.2. So from Theorem 4.2, the stochastic fractional differential equation with $A = -0.2$ and $f(t, x(t)) = \frac{1}{5\sqrt{2}} \sin x$ is exponentially stable.

Example 7.2. Consider the following stochastic fractional neutral differential equation with Lévy noise of the form

$$\begin{aligned} {}^C D^{0.6} \left(x(t) - \frac{1}{15} \exp(-x(t)) \right) &= -0.5x(t) + \frac{1}{15} \left(\frac{1}{1+t} \right) \frac{dW(t)}{dt} \\ &+ \int_{\mathbb{R}/\{0\}} \frac{1}{15} tz \frac{d\tilde{N}(t, z)}{dt}, \\ x(0) &= 1. \end{aligned} \quad (7.2)$$

Here $\alpha = 0.6$, $b = 0.5x(t)$, $p(t, x(t)) = \frac{1}{15} \exp(-x(t))$, $\sigma = \frac{1}{15} \left(\frac{1}{1+t} \right)$ and $g = \frac{1}{15} tz$. It can be easily seen that $b(t, x(t))$, $\sigma(t, x(t))$ and $g(t, x(t), z)$ satisfies the hypothesis (H4) and (H5). Hence by the Theorem 5.1 the stochastic fractional differential equation (7.2) has a unique solution. Also the equation (7.2) satisfies the condition of Theorem 6.2. So from Theorem 6.2 the stochastic fractional differential equation with $A = -0.5$ is exponentially stable.

8. CONCLUSION

Existence and uniqueness results for neutral stochastic fractional differential equations with Gaussian noise and Lévy noise are obtained. The Picard-Lindelöf successive approximation technique is used to obtain the existence results. Further sufficient conditions for the stability of neutral systems with Gaussian noise and Lévy noise are established by deriving the equivalent nonlinear integral equation corresponding to the system. Examples are given to illustrate the theory.

REFERENCES

- [1] R.P. Agarwal, Y. Zhou and Y. He, *Existence of fractional neutral functional differential equations*, *Compu. Math. Appl.*, **59** (2010), 1095-1100.
- [2] B. Boufoussi and S. Hajji, *Successive approximation of neutral functional stochastic differential equations with jumps*, *Statistics Prob. Lett.*, **80** (2010), 324-332.
- [3] A. Chadha and S.N. Bora, *Stability analysis for neutral stochastic differential equation of second order driven by Poisson jumps*, *J. Math. Physics*, **58** (2017), 112703.
- [4] Y. Chen, W.X. Zheng and A. Xue, *A new result on stability analysis for stochastic neutral systems*, *Automatica*, **46** (2010), 2100-2104.
- [5] S. Cong, *On exponential stability conditions of linear neutral stochastic differential systems with time-varying delay*, *Inter.J. Robust and Nonlinear Control*, **23** (2013), 1265-1276.
- [6] J. Cui and L. Yan, *Existence result for fractional neutral stochastic integro-differential equations with infinite delay*, *J. Physics A: Mathematical and Theoretical*, **44** (2011), 335201.
- [7] V. Govindaraj, S. Priyadarshini, P. Sureshkumar and K. Balachandran, *Asymptotic stability of fractional Langevin systems*, *J. Appl. Nonlinear Dyna.*, **11** (2022), 635-650.

- [8] L. Huang and X. Mao, *Delay-dependent exponential stability of neutral stochastic delay systems*, IEEE Trans. Autom. Control, **54** (2009), 147-152.
- [9] R. Joice Nirmala, K. Balachandran and J.J. Trujillo, *Null controllability of fractional dynamical systems with constrained control*, Fractional Calcul. Appl. Anal., **20** (2017), 153-167.
- [10] C. Kausika, K. Balachandran, N. Annapoorani and J.K. Kim, *Existence and stability results of generalized fractional integrodifferential equations*, Nonlinear Funct. Anal. Appl., **26** (2021), 793-809.
- [11] Y. Li and B. Liu, *Existence of solution of nonlinear neutral stochastic differential inclusions with infinite delay*, Stochastic Anal. Appl., **25** (2007), 397-415.
- [12] R. Mabel Lizzy, K. Balachandran and J.J. Trujillo, *Controllability of nonlinear stochastic fractional neutral systems with multiple time varying delay in control*, Chaos, Solitons & Fractals, **102** (2017), 162-167.
- [13] X. Mao, *Stochastic Differential Equation and Applications, Second Edition*, Woodhead Publishing Limited, Cambridge, 2010.
- [14] X. Mao, Y. Shen and C. Yuan, *Almost surely asymptotic stability of neutral stochastic differential delay equations with Markovian switching*, Stochastic Proc. Their Appl., **118** (2008), 1385-1406.
- [15] J.Y. Park, K. Balachandran and N. Annapoorani, *Existence results for impulsive neutral functional integrodifferential equations with infinite delay*, Nonlinear Anal., **71** (2009), 3152-3162.
- [16] D. Qian, C. Li, R.P. Agarwal and P.J.Y. Wong, *Stability analysis of fractional differential system with Riemann-Liouville derivative*, Math. Comput Model., **52** (2010), 862-874.
- [17] A.G. Radwan, A.M. Soliman, A.S. Elwakil and A. Sedeek, *On the stability of linear systems with fractional order elements*, Chaos, Solitons Fractals, **40** (2009), 2317-2328.
- [18] P. Ramesh, M. Sambath, M.H. Mohd and K. Balachandran, *Stability analysis of the fractional order pre-predator model with infection*, Inter. J. Model. Simul., **41** (2021) 434-450.
- [19] A. Rathinasamy, K. Balachandran and J.K. Kim, *Existence and stability of solutions of neutral hybrid stochastic infinite delay differential equations with Poisson jumps*, Nonlinear Funct. Anal. Appl., **16** (2011), 123-139.
- [20] Y. Ren and L. Chen, *A note on the neutral stochastic functional differential equation with infinite delay and Poisson jumps in an abstract space*, J. Math. Physics, **50** (2009), 082704.
- [21] S.G. Samko, A.A. Kilbas and O.I. Marichev, *Fractional Integrals and Derivatives, Theory and Applications*, Gordon and Breach Science Publishers, Amsterdam, 1993.
- [22] P. Umamaheswari, K. Balachandran and N. Annapoorani, *Existence and stability results for Caputo fractional stochastic differential equations with Lévy noise*, Filomat, **34** (2020), 1739-1751.