

Invited Mini Review

The role of myokines in cancer: crosstalk between skeletal muscle and tumor

Se-Young Park^{1,2,3}, Byeong-Oh Hwang^{1,2,3} & Na-Young Song^{1,2,3,4,*}¹Department of Applied Life Science, The Graduate School, Yonsei University, Seoul 03722, ²BK21 Four Project, Yonsei University College of Dentistry, Seoul 03722, ³Department of Oral Biology, Yonsei University College of Dentistry, Seoul 03722, ⁴Oral Cancer Research Institute, Yonsei University College of Dentistry, Seoul 03722, Korea

Loss of skeletal muscle mass is a primary feature of sarcopenia and cancer cachexia. In cancer patients, tumor-derived inflammatory factors promote muscle atrophy via tumor-to-muscle effects, which is closely associated with poor prognosis. During the past decade, skeletal muscle has been considered to function as an autocrine, paracrine, and endocrine organ by releasing numerous myokines. The circulating myokines can modulate pathophysiology in the other organs, as well as in the tumor microenvironment, suggesting myokines function as muscle-to-tumor signaling molecules. Here, we highlight the roles of myokines in tumorigenesis, particularly in terms of crosstalk between skeletal muscle and tumor. Better understanding of tumor-to-muscle and muscle-to-tumor effects will shed light on novel strategies for the diagnosis and treatment of cancer. [BMB Reports 2023; 56(7): 365-373]

INTRODUCTION

Skeletal muscle comprises approximately 40% of the whole body weight, and contains the largest body protein pool (1). The major functions of skeletal muscle are to contract to produce movement, maintain body temperature, store nutrients, and stabilize joints (1). In this regard, skeletal muscle wasting is closely associated with numerous human diseases (2). Sarcopenia is characterized by the decline in skeletal muscle mass and strength, emerging as a major health problem, particularly related to aging (3). Thus, maintaining skeletal muscle mass is crucial for human health and wellness.

During the past decade, skeletal muscle has been recognized as an endocrine organ (4). Skeletal muscle is composed

of various types of cells, such as myocytes, fibroblasts, adipocytes, neurons, and connective tissues, which can release bioactive molecules referred to as 'myokines' (5). The myokines are peptides that are synthesized, expressed, and secreted by skeletal muscle fibers, including cytokines, interleukins (ILs), and neurotrophins (5). These myokines can regulate extracellular matrix organization, angiogenesis, and metabolism in both paracrine and endocrine manners (6).

Notably, most of the patients with advanced cancers represent sarcopenia with or without loss of fat mass, distinct from age-related sarcopenia or malnutrition, which is termed 'cachexia' (7). The cancer-associated cachexia is a multi-organ metabolic syndrome, which is not fully recovered by nutritional support, and accounts for 20% of cancer deaths (8). In this regard, skeletal muscle is suggested as a crucial player in carcinogenesis; however, its underlying mechanism is not fully understood. Herein, we focus on how skeletal muscle regulates cancer promotion and progression by affecting the tumor microenvironment (TME), particularly through myokines.

SKELETAL MUSCLE MASS AND CANCER: TUMOR-TO-MUSCLE EFFECTS

Cancer cachexia: tumor-to-muscle effects

Cancer cachexia is a devastating syndrome of remarkable weight loss, muscle wasting, and anorexia, which occurs in about 30% of total cancer patients, and in 70-80% of advanced cancer patients (8). Both cancer cachexia and sarcopenia show loss of skeletal muscle mass, sharing a complex pathophysiology, while cachexia is involved with more inflammatory responses than sarcopenia (9). Cancer has long been considered as a chronic and systemic inflammatory status that creates TME more favorable to tumor promotion and progression, leading to metabolic alterations in multiple organs (10). Among the pro-inflammatory factors released from tumors, IL-6 and tumor necrosis factor α (TNF- α) are well known to trigger skeletal muscle wasting (11, 12). Notably, serum levels of IL-6 and TNF- α were dramatically increased in pancreatic cancer patients with weight loss, compared to those without weight change (13, 14). A prospective cohort study further demonstrated that the plasma IL-6 level was markedly higher

*Corresponding author. Tel: +82-2-2228-3056; Fax: +82-2-364-7113; E-mail: nysong608@yuhs.ac

<https://doi.org/10.5483/BMBRep.2023-0064>

Received 16 April 2023, Revised 2 May 2023,
Accepted 1 June 2023, Published online 13 June 2023

Keywords: Cancer cachexia, Myokine, Sarcopenia, Skeletal muscle, Tumor

in cachectic lung cancer patients than in non-cachectic patients (15). Higher IL-6 and TNF- α levels were associated with cachexia, as well as mortality in cancer patients (16, 17). In tumor-bearing murine models, the elevated serum levels of IL-6 and/or TNF- α were also correlated with skeletal muscle loss, further supporting the sarcopenic effects of cancer-derived inflammatory cytokines (18, 19). However, ablation of IL-6 attenuated skeletal muscle wasting in tumor-bearing mice (19, 20). Similarly, administration of an anti-TNF- α antibody improved cachectic conditions in tumor-bearing mice (21). As illustrated in Fig. 1, these data suggest that cancer-derived IL-6 and/or TNF- α promote skeletal muscle wasting via tumor-to-muscle effects.

Skeletal muscle mass as a cancer prognostic marker

Notably, low skeletal muscle mass is correlated with poor prognosis in various types of cancer, including pancreatic and oesophageal cancers (8, 22, 23). In contrast, increased skeletal muscle mass was associated with longer survival in pediatric patients with malignant solid cancers (24). Thus, skeletal muscle mass is considered a putative marker for predicting cancer prognosis.

Moreover, skeletal muscle mass can affect chemotherapy-induced toxicity in cancer patients. Cancer patients with loss of skeletal muscle mass presented susceptibility to chemotherapy-induced toxicity and poor prognosis, compared to patients with lean body mass (25, 26). A meta-analysis including 48 studies has demonstrated that low skeletal muscle mass is associated with dose-limiting toxicities in cancer patients under neoadjuvant/adjuvant chemotherapies or curative radio-chemotherapies (27). In cachectic tumor-bearing mice, 5-fluorouracil

chemotherapy exacerbated skeletal muscle wasting (28). Likewise, sarcopenia has been reported as a poor prognostic marker in cancer patients treated with immune checkpoint inhibitors, implying that skeletal muscle mass is crucial for optimizing chemotherapeutic regimens (29, 30). Taken together, skeletal muscle mass can reflect the body status in cancer patients and skeletal muscle might prevent and/or block carcinogenic processes, possibly through its secretome, while tumors induce skeletal muscle atrophy via secretion of IL-6 and TNF- α .

MYOKINES: MUSCLE-TO-TUMOR EFFECTS

Myokines, exercise, and cancer

In addition to tumor-to-muscle effects, skeletal muscle can modulate TME through its secretome. These skeletal muscle-derived molecules, termed myokines, include more than 600 proteins, such as cytokines (IL-8; IL-15) and neurotrophins (brain-derived neurotrophic factor, BDNF) (5). Myokines exert biological functions in autocrine, paracrine, and endocrine fashions (6). In particular, exercise can continuously stimulate skeletal muscle contraction, resulting in the secretion of a large amount of myokines (31). These myokines enhance skeletal muscle hypertrophy in an autocrine manner (31). Once released into the systemic circulation, myokines facilitate communication between muscle and the other organs (6).

It is well established that exercise is inversely correlated to cancer risk (32, 33). Exercise and/or regular physical activities can inhibit tumor promotion and progression in animal models, as well as cancer patients (34-36). The secreted myokines during the physical activities can regulate metabolic pathways,




Fig. 1. Tumor-to-muscle effects. Tumor tissues constantly release various types of pro-inflammatory molecules into the bloodstream, maintaining the systemic and chronic inflammatory status. The circulating tumor-derived inflammatory factors, such as IL-6 and TNF- α , promote skeletal muscle atrophy through activating ubiquitin-dependent proteasomal degradation, apoptosis, autophagy, and/or lysosomal degradation. Loss of skeletal muscle is closely associated with poor survival in cancer patients.

immune system, and inflammation in TME (37). Thus, myokines play a key role in muscle-to-tumor effects, which mediates the beneficial effects of exercise in cancer patients. In the following sections, we discuss how skeletal muscle-derived myokines modulate cancer promotion and progression by affecting TME. Among various myokines, we focus on myokines that are known to be released by exercise from skeletal muscle and to function in TME in both direct and/or indirect manners.

Effects of myokines on TME

Irisin: Irisin, a recently discovered myokine, is a cleaved form of fibronectin type III domain-containing 5 (FNDC5) (38). During exercise, muscle contraction can upregulate the plasma levels of FNDC5 and its proteolytic product irisin (38, 39). Notably, it has been reported that the serum level of irisin was significantly reduced in patients with breast and liver cancers, compared to healthy individuals, implying the tumor-suppressive role of irisin (40, 41). In line with this notion, irisin treatment inhibited cell proliferation in breast and lung cancer cell lines (42, 43). Moreover, irisin prevented epithelial-to-mesenchymal transition (EMT), a major driver of cancer metastasis (43, 44). Exposure to irisin or overexpression of FNDC5/irisin diminished migration and invasion in different types of human cancer cell lines (43, 45). In ovarian cancer cell lines, irisin blocked hypoxia-inducible factor 1 α signaling pathways responsible for the expression of metastasis markers, supporting its anti-metastatic activities (46). In the xenograft glioblastoma mouse model, administration of irisin dramatically repressed tumor growth (47). Furthermore, bladder cancer patients with lower serum irisin levels showed higher mortality (48). Taken together, these data suggest that irisin is a tumor-suppressive myokine.

Decorin: Decorin (DCN) is a member of the small leucine-rich repeat proteoglycan family that is ubiquitously expressed in connective tissues (49). During exercise, muscle contraction can upregulate DCN expression in human tendon and muscle, and also induce its secretion into the systemic circulation (50, 51). DCN has been reported to act as a soluble inhibitor of pan-receptor tyrosine kinases, such as Met receptor and epidermal growth factor receptor (EGFR), pro-survival signals for tumor cells (52). DCN treatment or overexpression reduced cell proliferation, migration, and EMT marker expression in various types of cancer cells, which can be mediated by the inhibition of Met/EGFR pathways (53-56). In consistent manner, systemic injection or overexpression of DCN suppressed tumor growth through blocking Met/EGFR pathways in the orthotopic xenograft models (54, 57). DCN significantly reduced angiogenesis and pulmonary metastasis in the tumor xenograft models (58, 59). In contrast, DCN ablation showed more and larger tumors in mice subcutaneously injected with murine colon cancer cells (53). Moreover, it has been reported that DCN is down-regulated in tumors from patients with invasive breast and gastric cancers, which is associated with poor prognosis (60, 61). Overall, these data suggest that muscle-derived DCN can exert anti-carcinogenic activities in an endocrine manner, parti-

cularly in patients with DCN-low tumors.

IL-15: IL-15, a cytokine that belongs to the four α -helix bundle family, is responsible for natural killer (NK) and T cell immunity (62). Besides the immune cells, IL-15 is abundantly expressed in skeletal muscle, acting as a myokine. Exercise increased IL-15 levels in skeletal muscle and circulation in human subjects (63-65). It is noteworthy that exercise-induced IL-15 expression was correlated with better prognosis in pancreatic cancer cohorts, suggesting IL-15 as an anti-tumorigenic myokine (66). Consistently, the serum IL-15 level was significantly increased by exercise in prostate cancer patients (35). The pooled sera isolated from the exercising patients reduced cell growth in human prostate cancer DU-145 cells (35). Its anti-cancer effects could be attributed to immune modulation in TME. In tumor-bearing mice, IL-15 was co-localized with CD8 T and NK cells in TME (67). Heterodimeric IL-15 injection suppressed metastatic burden through increasing intratumoral CD8 T and NK cells in several different mouse cancer models, implying the involvement of CD8 T and NK cells in immune modulation by IL-15 (68). Likewise, subcutaneous injection of recombinant human IL-15 induced the number of circulating CD8 T and NK cells, while decreasing the number of leukemic cells in the blood of patients with T-cell malignancies (69). Recently, Kurz *et al.* demonstrated that exercise enhanced CD8 T cell immunity via increasing IL-15, which suppressed tumor growth in a pancreatic cancer mouse model (70). Overall, these data support that skeletal muscle-derived IL-15 promotes anti-tumor immunity.

Secreted Protein Acidic and Rich in Cysteine: Secreted Protein Acidic and Rich in Cysteine (SPARC), also known as osteonectin, is a calcium-binding matricellular glycoprotein that is involved in development, wound repair, tissue remodeling, differentiation, and proliferation (71). It has been reported that exercise induced circulating SPARC levels in healthy adult men (72). Exercise increased skeletal muscle expression and serum level of SPARC in both human and mice, which suppressed colon tumorigenesis (34). In wild type (WT) mice, regular exercise significantly reduced the number of aberrant crypts driven by a chemical carcinogen azoxymethane (34). However, SPARC-null mice developed more colon tumors than WT mice, which was not attenuated by exercise (34). In a rat model bearing colon cancers, high-intensity swimming training enhanced serum SPARC levels, finally suppressing tumorigenesis (73). Likewise, exercise significantly stimulated SPARC release from the skeletal muscle in patients with metastatic castrate-resistant prostate cancer (35). The serum isolated from these patients suppressed cell growth in human prostate cancer cell lines (35). These data suggest that exercise-induced SPARC can act as an anti-carcinogenic muscle-to-tumor signaling molecule.

SPARC can suppress multiple stages of carcinogenesis. SPARC inhibited cell proliferation in different types of cancer cell lines (74-76). In consistent manner, SPARC-null mice showed accelerated growth of tumor grafts, compared with WT mice (75, 77). Moreover, overexpression of SPARC hampered the expres-

sion of EMT markers and migratory capability in gastric cancer cells (78). In athymic nude mice subcutaneously implanted with gastric cancer cells, SPARC overexpression diminished tumor growth and angiogenesis (78, 79). Said *et al.* demonstrated that SPARC silencing promoted tumor growth and invasiveness, while reducing stromal collagen in the transgenic prostate cancer mouse model (80). Furthermore, SPARC enhanced chemosensitivity *in vivo* (78, 81). Taken together, these data suggest SPARC as a tumor suppressive myokine.

In contrast, SPARC can act as a tumor promoter as well. SPARC is highly expressed in metastatic tumors, such as glioblastomas and melanoma (82). SPARC increased the expression of EMT markers, enhancing invasion and migration abilities in cancer cells (83, 84). Tumor-derived SPARC induced vascular permeability and lung metastasis in mice inoculated with melanoma cells via tail vein injection (85). Moreover, SPARC overexpression promoted cell proliferation in liver and pancreatic cancer cells, further supporting tumor-derived SPARC as a tumor promoter (86, 87). It has been reported that certain types of cancers display contradictory compartmentalized expression of SPARC in TME, which contributes to the complex functions of SPARC in tumorigenesis (88). In the case of pancreatic cancer, stromal expression of SPARC was associated with poor prognosis, while its tumoral expression showed no significance (89). Furthermore, Pan *et al.* suggested that the oncogenic roles of SPARC might be due to its autocrine secretion into TME (87). Therefore, the roles of SPARC on TME can vary depending on the site of secretion, either TME or skeletal muscle, as well as its localization in TME.

Oncostatin M: Oncostatin M (OSM) is a member of the IL-6 family cytokines that exerts biological functions through binding to its heterodimeric receptor complex comprising OSM receptor β (OSMR β) and gp130 (90). Interestingly, exercise remarkably increased the serum level of OSM in healthy individuals and cancer patients (35, 91). OSM has been reported to suppress cell growth in breast cancer, lung adenocarcinoma, and glioblastoma cells, implying OSM as an anti-tumorigenic myokine (92-94). In the orthotopic mouse cancer model, OSM-expressing glioma cells were unable to generate tumor mass (95). In prostate cancer patients undertaking androgen deprivation therapy, the exercise program significantly enhanced the serum level of OSM, while the expression levels of the other myokines were not altered (96). Human prostate cancer DU145 cells exposed to the sera from these patients showed retarded cell growth, further supporting the anti-cancer effect of exercise-induced OSM (96). Similarly, the sera pooled from exercising mice contained abundant OSM levels, which inhibited the proliferation of human breast cancer MCF-7 cells (97). OSM-containing conditioned media reduced EMT phenotypes and cancer stemness in lung adenocarcinoma cells (98). Moreover, treatment with OSM suppressed migration and invasion in human lung adenocarcinoma cell lines, and also blocked tumor metastasis *in vivo* (92). These data suggest that skeletal muscle-secreted OSM during exercise can act as an

anti-tumorigenic myokine.

However, it is controversial as to whether OSM is anti-tumorigenic or pro-tumorigenic. In addition to muscle-to-tumor effects, OSM can be released within TME by tumor cells and/or stromal cells (99, 100). TME-derived OSM seems to exert pro-tumorigenic functions. Treatment with OSM induced proliferation in ovarian and prostate cancer cell lines (101, 102). OSM upregulated the expression of EMT markers and cancer stemness in pancreatic and cervical cancer cells (103, 104). In consistent manner, paracrine expression of OSM promoted invasiveness and angiogenesis of the grafts *in vivo* (105, 106). Of note, OSM suppressed or maintained the tumor sizes in xenograft models, while promoting metastatic features (104, 106). This might be explained by the lack of OSMR complex. OSM inhibited the growth of OSMR-intact cell growth, while cells with OSMR loss showed OSM resistance (102, 107). Thus, OSM might differentially affect tumorigenesis, depending on the cellular and molecular contexts.

BDNF: BDNF is a member of the nerve growth factor family that is mainly produced by neurons, regulating neuronal survival, differentiation, and apoptosis (108). In addition to neuron cells, skeletal muscle can synthesize BDNF (109). BDNF-null mice exhibited abnormal differentiation of myoblasts, decreased myotube size, and delayed muscle regeneration, suggesting that BDNF is a myokine responsible for skeletal muscle physiology (110). Of note, BDNF can be secreted from stromal cells in TME (111, 112). Higher expression of BDNF was associated with poor prognosis in various types of cancers (112, 113). Moreover, BDNF promoted cancer cell proliferation, migration, and invasion (112, 114). These data suggest that TME-derived BDNF can act as an oncogenic regulator.

In the case of skeletal-muscle derived BDNF, exercise can induce the skeletal muscle expression of BDNF in murine and human (109, 115). However, exercise was not able to increase plasma BDNF levels in human subjects, which means that muscle-derived BDNF might not be directly involved in muscle-to-tumor signaling (115). Rather, exercise seems to stimulate BDNF production in the brain (116). Hypothalamic BDNF expression inhibited tumor growth, while enhancing the anti-cancer CD8 T-cell immunity in C57BL/6 mice subcutaneously implanted with mouse melanoma cells (117, 118). Furthermore, brain infusion of BDNF reduced migration and intratumoral macrophage infiltration in the glioma mouse model (119). Although there is no strong evidence yet that skeletal muscle-derived BDNF directly affects tumorigenesis, exercise-induced hypothalamic BDNF expression can inhibit tumor promotion and progression. Taken together, these data imply that BDNF exerts tumor suppressive functions rather indirectly through the muscle-to-brain effects. Further investigation is required to elucidate a direct role of skeletal muscle-derived BDNF as a myokine.




Fig. 2. Muscle-to-tumor effects. During exercise, muscle contraction stimulates the production of myokines. Muscle-derived myokines enter the circulation, and affect tumor promotion and progression by modulating TME in an endocrine fashion. In particular, BDNF seems to regulate TME rather indirectly through muscle-to-brain effects. Overall, exercise-derived myokines exert anti-tumorigenic functions in both direct and/or indirect manners.

CONCLUDING REMARKS

Currently, skeletal muscle is well recognized as a secretory organ (4). During exercise, skeletal muscle releases various types of myokines that control exercise adaptations in the autocrine and/or paracrine fashions (5). Moreover, myokines can travel through the circulation to the other organs, such as adipose tissues and brain, modulating pathophysiology (120, 121). Similarly, myokines play a crucial role in the interplay between skeletal muscle and tumor, which regulates tumor promotion and progression. While tumor-derived IL-6 and/or TNF- α contribute to skeletal muscle atrophy (Fig. 1), skeletal muscle-secreted myokines promote anti-tumor immunities and suppress tumor growth and metastatic abilities (Fig. 2). Thus, maintaining and/or recovery of skeletal muscle mass, a source of myokines, would be a novel and proactive approach to cancer treatment. Taken together, the integrative understanding of crosstalk between skeletal muscle and tumor will provide effective diagnosis/prognosis markers and novel chemotherapeutic targets.

ACKNOWLEDGEMENTS

This research was supported by National Research Foundation of Korea (NRF) Grants funded by the Korean Government (grant numbers NRF-2020R1C1C1003338 and NRF-2022M3A9F3016364 to NYS), and by the Yonsei Signature Research Cluster Program (2023-22-0011).

CONFLICTS OF INTEREST

The authors have no conflicting interests.

REFERENCES

1. Frontera WR and Ochala J (2015) Skeletal muscle: a brief review of structure and function. *Calcif Tissue Int* 96, 183-195
2. Cohen S, Nathan JA and Goldberg AL (2015) Muscle wasting in disease: molecular mechanisms and promising therapies. *Nat Rev Drug Discov* 14, 58-74
3. Prokopidis K, Giannos P, Reginster JY et al (2023) Sarcopenia is associated with a greater risk of polypharmacy and number of medications: a systematic review and meta-analysis. *J Cachexia Sarcopenia Muscle* 14, 671-683
4. Iizuka K, Machida T and Hirafuji M (2014) Skeletal muscle is an endocrine organ. *J Pharmacol Sci* 125, 125-131
5. Pedersen BK and Febbraio MA (2012) Muscles, exercise and obesity: skeletal muscle as a secretory organ. *Nat Rev Endocrinol* 8, 457-465
6. Severinsen MCK and Pedersen BK (2020) Muscle-organ crosstalk: the emerging roles of myokines. *Endocr Rev* 41, 594-609
7. Argiles JM, Busquets S, Stemmler B and Lopez-Soriano FJ (2015) Cachexia and sarcopenia: mechanisms and potential targets for intervention. *Curr Opin Pharmacol* 22, 100-106
8. Baracos VE, Martin L, Korc M, Guttridge DC and Fearon KCH (2018) Cancer-associated cachexia. *Nat Rev Dis Primers* 4, 17105
9. Ali S and Garcia JM (2014) Sarcopenia, cachexia and aging: diagnosis, mechanisms and therapeutic options - a mini-review. *Gerontology* 60, 294-305
10. Greten FR and Grivnickov SI (2019) Inflammation and cancer: triggers, mechanisms, and consequences. *Immunity* 51, 27-41
11. Narsale AA and Carson JA (2014) Role of interleukin-6

- in cachexia: therapeutic implications. *Curr Opin Support Palliat Care* 8, 321-327
12. Patel HJ and Patel BM (2017) TNF-alpha and cancer cachexia: molecular insights and clinical implications. *Life Sci* 170, 56-63
 13. Karayiannakis AJ, Syrigos KN, Polychronidis A, Pitiakoudis M, Bounovas A and Simopoulos K (2001) Serum levels of tumor necrosis factor-alpha and nutritional status in pancreatic cancer patients. *Anticancer Res* 21, 1355-1358
 14. Okada S, Okusaka T, Ishii H et al (1998) Elevated serum interleukin-6 levels in patients with pancreatic cancer. *Jpn J Clin Oncol* 28, 12-15
 15. Srdic D, Plestina S, Sverko-Peternac A, Nikolac N, Simundic AM and Samarzija M (2016) Cancer cachexia, sarcopenia and biochemical markers in patients with advanced non-small cell lung cancer-chemotherapy toxicity and prognostic value. *Support Care Cancer* 24, 4495-4502
 16. Bilir C, Engin H, Can M, Temi YB and Demirtas D (2015) The prognostic role of inflammation and hormones in patients with metastatic cancer with cachexia. *Med Oncol* 32, 56
 17. Utech AE, Tadros EM, Hayes TG and Garcia JM (2012) Predicting survival in cancer patients: the role of cachexia and hormonal, nutritional and inflammatory markers. *J Cachexia Sarcopenia Muscle* 3, 245-251
 18. Matsuyama T, Ishikawa T, Okayama T et al (2015) Tumor inoculation site affects the development of cancer cachexia and muscle wasting. *Int J Cancer* 137, 2558-2565
 19. Rupert JE, Narasimhan A, Jengelle DHA et al (2021) Tumor-derived IL-6 and trans-signaling among tumor, fat, and muscle mediate pancreatic cancer cachexia. *J Exp Med* 218, e20190450
 20. Pototschnig I, Feiler U, Diwoky C et al (2023) Interleukin-6 initiates muscle- and adipose tissue wasting in a novel C57BL/6 model of cancer-associated cachexia. *J Cachexia Sarcopenia Muscle* 14, 93-107
 21. Kang EA, Park JM, Jin W, Tchahc H, Kwon KA and Hahm KB (2022) Amelioration of cancer cachexia with preemptive administration of tumor necrosis factor-alpha blocker. *J Clin Biochem Nutr* 70, 117-128
 22. Choi MH and Yoon SB (2022) Sarcopenia in pancreatic cancer: effect on patient outcomes. *World J Gastrointest Oncol* 14, 2302-2312
 23. Fang P, Zhou J, Xiao X et al (2023) The prognostic value of sarcopenia in oesophageal cancer: a systematic review and meta-analysis. *J Cachexia Sarcopenia Muscle* 14, 3-16
 24. Omori A, Kawakubo N, Takemoto J et al (2022) Effects of changes in skeletal muscle mass on the prognosis of pediatric malignant solid tumors. *Pediatr Surg Int* 38, 1829-1838
 25. Becker JN, Hermann R, Wichmann J, Sonnhoff M, Christiansen H and Bruns F (2023) Low skeletal muscle mass is predictive of dose-limiting toxicities in head and neck cancer patients undergoing low-dose weekly cisplatin chemoradiotherapy. *PLoS One* 18, e0282015
 26. de Jong C, Chargin N, Herder GJM et al (2022) The association between skeletal muscle measures and chemotherapy-induced toxicity in non-small cell lung cancer patients. *J Cachexia Sarcopenia Muscle* 13, 1554-1564
 27. Surov A, Pech M, Gessner D et al (2021) Low skeletal muscle mass is a predictor of treatment related toxicity in oncologic patients. A meta-analysis. *Clin Nutr* 40, 5298-5310
 28. Murphy KT, Swiderski K, Ryall JG et al (2021) Mechanisms of chemotherapy-induced muscle wasting in mice with cancer cachexia. *JCSM Rapid Communications* 5, 102-116
 29. Cortellini A, Bozzetti F, Palumbo P et al (2020) Weighing the role of skeletal muscle mass and muscle density in cancer patients receiving PD-1/PD-L1 checkpoint inhibitors: a multicenter real-life study. *Sci Rep* 10, 1456
 30. Li S, Liu Z, Ren Y et al (2022) Sarcopenia was a poor prognostic predictor for patients with advanced lung cancer treated with immune checkpoint inhibitors. *Front Nutr* 9, 900823
 31. Hoffmann C and Weigert C (2017) Skeletal muscle as an endocrine organ: the role of myokines in exercise adaptations. *Cold Spring Harb Perspect Med* 7, a029793
 32. McTiernan A, Friedenreich CM, Katzmarzyk PT et al (2019) Physical activity in cancer prevention and survival: a systematic review. *Med Sci Sports Exerc* 51, 1252-1261
 33. Moore SC, Lee IM, Weiderpass E et al (2016) Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Intern Med* 176, 816-825
 34. Aoi W, Naito Y, Takagi T et al (2013) A novel myokine, secreted protein acidic and rich in cysteine (SPARC), suppresses colon tumorigenesis via regular exercise. *Gut* 62, 882-889
 35. Kim JS, Taaffe DR, Galvao DA et al (2022) Acute effect of high-intensity interval aerobic exercise on serum myokine levels and resulting tumour-suppressive effect in trained patients with advanced prostate cancer. *Prostate Cancer Prostatic Dis* doi: 10.1038/s41391-022-00624-4
 36. Kim J-S, Galvão DA, Newton RU, Gray E and Taaffe DR (2021) Exercise-induced myokines and their effect on prostate cancer. *Nat Rev Urol* 18, 519-542
 37. Huang Q, Wu M, Wu X, Zhang Y and Xia Y (2022) Muscle-to-tumor crosstalk: the effect of exercise-induced myokine on cancer progression. *Biochim Biophys Acta Rev Cancer* 1877, 188761
 38. Boström P, Wu J, Jedrychowski MP et al (2012) A PGC1- α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature* 481, 463-468
 39. Huh JY, Panagiotou G, Mougios V et al (2012) FNDC5 and irisin in humans: I. Predictors of circulating concentrations in serum and plasma and II. mRNA expression and circulating concentrations in response to weight loss and exercise. *Metabolism* 61, 1725-1738
 40. Provatopoulou X, Georgiou GP, Kalogera E et al (2015) Serum irisin levels are lower in patients with breast cancer: association with disease diagnosis and tumor characteristics. *BMC Cancer* 15, 898
 41. Pazgan-Simon M, Zuwala-Jagiello J, Kukla M, Grzebyk E and Simon K (2020) Serum concentrations of selected adipokines in virus-related liver cirrhosis and hepatocellular carcinoma. *Clin Exp Hepatol* 6, 235-242

42. Gannon NP, Vaughan RA, Garcia-Smith R, Bisoffi M and Trujillo KA (2015) Effects of the exercise-inducible myokine irisin on malignant and non-malignant breast epithelial cell behavior in vitro. *Int J Cancer* 136, 197-202
43. Shao L, Li H, Chen J et al (2017) Irisin suppresses the migration, proliferation, and invasion of lung cancer cells via inhibition of epithelial-to-mesenchymal transition. *Biochem Biophys Res Commun* 485, 598-605
44. Zhang Y and Weinberg RA (2018) Epithelial-to-mesenchymal transition in cancer: complexity and opportunities. *Front Med* 12, 361-373
45. Xu L, Ye Y, Sun Y et al (2022) Low FNDC5/Irisin expression is associated with aggressive phenotypes in gastric cancer. *Front Pharmacol* 13, 981201
46. Alizadeh Zarei M, Seyed Hosseini E, Haddad Kashani H, Ahmad E and Nikzad H (2023) Effects of the exercise-inducible myokine irisin on proliferation and malignant properties of ovarian cancer cells through the HIF-1 alpha signaling pathway. *Sci Rep* 13, 170
47. Huang CW, Chang YH, Lee HH et al (2020) Irisin, an exercise myokine, potently suppresses tumor proliferation, invasion, and growth in glioma. *FASEB J* 34, 9678-9693
48. Esawy MM and Abdel-Samd KM (2020) The diagnostic and prognostic roles of serum irisin in bladder cancer. *Curr Probl Cancer* 44, 100529
49. Sofeu Feugaing DD, Gotte M and Viola M (2013) More than matrix: the multifaceted role of decorin in cancer. *Eur J Cell Biol* 92, 1-11
50. Heinemeier KM, Bjerrum SS, Schjerling P and Kjaer M (2013) Expression of extracellular matrix components and related growth factors in human tendon and muscle after acute exercise. *Scand J Med Sci Sports* 23, 150-161
51. Kanzleiter T, Rath M, Gorgens SW et al (2014) The myokine decorin is regulated by contraction and involved in muscle hypertrophy. *Biochem Biophys Res Commun* 450, 1089-1094
52. Neill T, Schaefer L and Iozzo RV (2015) Decoding the matrix: instructive roles of proteoglycan receptors. *Biochemistry* 54, 4583-4598
53. Bi X, Pohl NM, Qian Z et al (2012) Decorin-mediated inhibition of colorectal cancer growth and migration is associated with E-cadherin in vitro and in mice. *Carcinogenesis* 33, 326-330
54. Hu X, Villodre ES, Larson R et al (2021) Decorin-mediated suppression of tumorigenesis, invasion, and metastasis in inflammatory breast cancer. *Commun Biol* 4, 72
55. Chen H, Wang Z, Yang N, Zhang J and Liang Z (2022) Decorin inhibits proliferation and metastasis in human bladder cancer cells by upregulating P21. *Medicine (Baltimore)* 101, e29760
56. Jia Y, Feng Q, Tang B et al (2021) Decorin suppresses invasion and EMT phenotype of glioma by inducing autophagy via c-Met/Akt/mTOR axis. *Front Oncol* 11, 659353
57. Buraschi S, Pal N, Tyler-Rubinstein N, Owens RT, Neill T and Iozzo RV (2010) Decorin antagonizes Met receptor activity and down-regulates beta-catenin and Myc levels. *J Biol Chem* 285, 42075-42085
58. Reed CC, Waterhouse A, Kirby S et al (2005) Decorin prevents metastatic spreading of breast cancer. *Oncogene* 24, 1104-1110
59. Grant DS, Yenisey C, Rose RW, Tootell M, Santra M and Iozzo RV (2002) Decorin suppresses tumor cell-mediated angiogenesis. *Oncogene* 21, 4765-4777
60. Basak D, Jamal Z, Ghosh A et al (2021) Reciprocal interplay between asporin and decorin: implications in gastric cancer prognosis. *PLoS One* 16, e0255915
61. Troup S, Njue C, Kliewer EV et al (2003) Reduced expression of the small leucine-rich proteoglycans, lumican, and decorin is associated with poor outcome in node-negative invasive breast cancer. *Clin Cancer Res* 9, 207-214
62. Di Sabatino A, Calarota SA, Vidali F, Macdonald TT and Corazza GR (2011) Role of IL-15 in immune-mediated and infectious diseases. *Cytokine Growth Factor Rev* 22, 19-33
63. Tamura Y, Watanabe K, Kantani T, Hayashi J, Ishida N and Kaneki M (2011) Upregulation of circulating IL-15 by treadmill running in healthy individuals: is IL-15 an endocrine mediator of the beneficial effects of endurance exercise? *Endocr J* 58, 211-215
64. Riechman SE, Balasekaran G, Roth SM and Ferrell RE (2004) Association of interleukin-15 protein and interleukin-15 receptor genetic variation with resistance exercise training responses. *J Appl Physiol* (1985) 97, 2214-2219
65. Nielsen AR, Mounier R, Plomgaard P et al (2007) Expression of interleukin-15 in human skeletal muscle effect of exercise and muscle fibre type composition. *J Physiol* 584, 305-312
66. Luo Z, He Z, Qin H et al (2022) Exercise-induced IL-15 acted as a positive prognostic implication and tumor-suppressed role in pan-cancer. *Front Pharmacol* 13, 1053137
67. Reyes AF, Goldusky J, Bhimalli P, Marzo AL and Schneider JR (2022) Tracking fluorescently labeled IL-15 and anti-PD-1 in the tumor microenvironment and draining lymph nodes. *J Immunol Methods* 505, 113253
68. Stravokefalou V, Stellas D, Karaliota S et al (2022) Heterodimeric IL-15 (hetIL-15) reduces circulating tumor cells and metastasis formation improving chemotherapy and surgery in 4T1 mouse model of TNBC. *Front Immunol* 13, 1014802
69. Miljkovic MD, Dubois SP, Muller JR et al (2023) Interleukin-15 augments NK cell-mediated ADCC of alemtuzumab in patients with CD52+ T-cell malignancies. *Blood Adv* 7, 384-394
70. Kurz E, Hirsch CA, Dalton T et al (2022) Exercise-induced engagement of the IL-15/IL-15Ralpha axis promotes anti-tumor immunity in pancreatic cancer. *Cancer Cell* 40, 720-737 e725
71. Bradshaw AD and Sage EH (2001) SPARC, a matricellular protein that functions in cellular differentiation and tissue response to injury. *J Clin Invest* 107, 1049-1054
72. Miyamoto T, Shimizu Y, Matsuo Y et al (2022) Effects of exercise intensity and duration on a myokine, secreted protein acidic and rich in cysteine. *Eur J Sport Sci* 22, 1401-1410
73. Matsuo K, Sato K, Suemoto K et al (2017) A mechanism underlying preventive effect of high-intensity training on colon cancer. *Med Sci Sports Exerc* 49, 1805-1816
74. Yiu GK, Chan WY, Ng SW et al (2001) SPARC (secreted

- protein acidic and rich in cysteine) induces apoptosis in ovarian cancer cells. *Am J Pathol* 159, 609-622
75. Brekken RA, Puolakkainen P, Graves DC, Workman G, Lubkin SR and Sage EH (2003) Enhanced growth of tumors in SPARC null mice is associated with changes in the ECM. *J Clin Invest* 111, 487-495
 76. Schultz C, Lemke N, Ge S, Golembieski WA and Rempel SA (2002) Secreted protein acidic and rich in cysteine promotes glioma invasion and delays tumor growth in vivo. *Cancer Res* 62, 6270-6277
 77. Puolakkainen PA, Brekken RA, Muneer S and Sage EH (2004) Enhanced growth of pancreatic tumors in SPARC-null mice is associated with decreased deposition of extracellular matrix and reduced tumor cell apoptosis. *Mol Cancer Res* 2, 215-224
 78. Ma J, Ma Y, Chen S et al (2021) SPARC enhances 5-FU chemosensitivity in gastric cancer by modulating epithelial-mesenchymal transition and apoptosis. *Biochem Biophys Res Commun* 558, 134-140
 79. Zhang JL, Chen GW, Liu YC et al (2012) Secreted protein acidic and rich in cysteine (SPARC) suppresses angiogenesis by down-regulating the expression of VEGF and MMP-7 in gastric cancer. *PLoS One* 7, e44618
 80. Said N, Frierson H, Chernauskas D, Conaway M, Motamed K and Theodorescu D (2009) The role of SPARC in the TRAMP model of prostate carcinogenesis and progression. *Oncogene* 28, 3487-3498
 81. Rahman M, Chan AP, Tang M and Tai IT (2011) A peptide of SPARC interferes with the interaction between caspase8 and Bcl2 to resensitize chemoresistant tumors and enhance their regression in vivo. *PLoS One* 6, e26390
 82. Feng J and Tang L (2014) SPARC in tumor pathophysiology and as a potential therapeutic target. *Curr Pharm Des* 20, 6182-6190
 83. Grant JL, Fishbein MC, Hong LS et al (2014) A novel molecular pathway for snail-dependent, SPARC-mediated invasion in non-small cell lung cancer pathogenesis. *Cancer Prev Res (Phila)* 7, 150-160
 84. Lopez-Moncada F, Torres MJ, Lavanderos B, Cerda O, Castellon EA and Contreras HR (2022) SPARC induces E-cadherin repression and enhances cell migration through integrin α v β 3 and the transcription factor ZEB1 in prostate cancer cells. *Int J Mol Sci* 23, 5874
 85. Tichet M, Prod'Homme V, Fenouille N et al (2015) Tumour-derived SPARC drives vascular permeability and extravasation through endothelial VCAM1 signalling to promote metastasis. *Nat Commun* 6, 6993
 86. Gao ZW, Liu C, Yang L et al (2021) SPARC Overexpression promotes liver cancer cell proliferation and tumor growth. *Front Mol Biosci* 8, 775743
 87. Pan K, Huang X and Jia X (2021) SPARC promotes pancreatic cancer cell proliferation and migration through autocrine secretion into the extracellular milieu. *Oncol Lett* 21, 485
 88. Arnold SA and Brekken RA (2009) SPARC: a matricellular regulator of tumorigenesis. *J Cell Commun Signal* 3, 255-273
 89. Neuzillet C, Tijeras-Raballand A, Cros J, Faivre S, Hamel P and Raymond E (2013) Stromal expression of SPARC in pancreatic adenocarcinoma. *Cancer Metastasis Rev* 32, 585-602
 90. Richards CD (2013) The enigmatic cytokine oncostatin m and roles in disease. *ISRN Inflamm* 2013, 512103
 91. Hwang JH, McGovern J, Minett GM et al (2020) Mobilizing serum factors and immune cells through exercise to counteract age-related changes in cancer risk. *Exerc Immunol Rev* 26, 80-99
 92. Pan CM, Wang ML, Chiou SH, Chen HY and Wu CW (2016) Oncostatin M suppresses metastasis of lung adenocarcinoma by inhibiting SLUG expression through coordination of STATs and PIASs signalings. *Oncotarget* 7, 60395
 93. Grant SL, Hammacher A, Douglas AM et al (2002) An unexpected biochemical and functional interaction between gp130 and the EGF receptor family in breast cancer cells. *Oncogene* 21, 460-474
 94. Halfter H, Friedrich M, Resch A et al (2006) Oncostatin M induces growth arrest by inhibition of Skp2, Cks1, and cyclin A expression and induced p21 expression. *Cancer Res* 66, 6530-6539
 95. Friedrich M, Hoss N, Stogbauer F et al (2001) Complete inhibition of in vivo glioma growth by oncostatin M. *J Neurochem* 76, 1589-1592
 96. Kim JS, Wilson RL, Taaffe DR, Galvão DA, Gray E and Newton RU (2022) Myokine expression and tumor-suppressive effect of serum after 12 wk of exercise in prostate cancer patients on ADT. *Med Sci Sports Exerc* 54, 197
 97. Hojman P, Dethlefsen C, Brandt C, Hansen J, Pedersen L and Pedersen BK (2011) Exercise-induced muscle-derived cytokines inhibit mammary cancer cell growth. *Am J Physiol Endocrinol Metab* 301, E504-E510
 98. Wang ML, Pan CM, Chiou SH et al (2012) Oncostatin m modulates the mesenchymal-epithelial transition of lung adenocarcinoma cells by a mesenchymal stem cell-mediated paracrine effect. *Cancer Res* 72, 6051-6064
 99. Tripathi C, Tewari BN, Kanchan RK et al (2014) Macrophages are recruited to hypoxic tumor areas and acquire a pro-angiogenic M2-polarized phenotype via hypoxic cancer cell derived cytokines Oncostatin M and Eotaxin. *Oncotarget* 5, 5350-5368
 100. Araujo AM, Abaurrea A, Azcoaga P et al (2022) Stromal oncostatin M cytokine promotes breast cancer progression by reprogramming the tumor microenvironment. *J Clin Invest* 132, 148667
 101. Li Q, Zhu J, Sun F, Liu L, Liu X and Yue Y (2011) Oncostatin M promotes proliferation of ovarian cancer cells through signal transducer and activator of transcription 3. *Int J Mol Med* 28, 101-108
 102. Mori S, Murakami-Mori K and Bonavida B (1999) Oncostatin M (OM) promotes the growth of DU 145 human prostate cancer cells, but not PC-3 or LNCaP, through the signaling of the OM specific receptor. *Anticancer Res* 19, 1011-1015
 103. Smigiel JM, Parameswaran N and Jackson MW (2017) Potent EMT and CSC phenotypes are induced by oncostatin-M in pancreatic cancer. *Mol Cancer Res* 15, 478-488
 104. Kucia-Tran JA, Tulkki V, Smith S et al (2016) Overexpression of the oncostatin-M receptor in cervical squa-

- mous cell carcinoma is associated with epithelial-mesenchymal transition and poor overall survival. *Br J Cancer* 115, 212-222
105. Smith DA, Kiba A, Zong Y and Witte ON (2013) Interleukin-6 and oncostatin-M synergize with the PI3K/AKT pathway to promote aggressive prostate malignancy in mouse and human tissues. *Mol Cancer Res* 11, 1159-1165
 106. Lapeire L, Hendrix A, Lambein K et al (2014) Cancer-associated adipose tissue promotes breast cancer progression by paracrine oncostatin M and Jak/STAT3 signaling. *Cancer Res* 74, 6806-6819
 107. Lacreusette A, Nguyen JM, Pandolfino MC et al (2007) Loss of oncostatin M receptor beta in metastatic melanoma cells. *Oncogene* 26, 881-892
 108. Bathina S and Das UN (2015) Brain-derived neurotrophic factor and its clinical implications. *Arch Med Sci* 11, 1164-1178
 109. Gomez-Pinilla F, Ying Z, Roy RR, Molteni R and Edgerton VR (2002) Voluntary exercise induces a BDNF-mediated mechanism that promotes neuroplasticity. *J Neurophysiol* 88, 2187-2195
 110. Clow C and Jasmin BJ (2010) Brain-derived neurotrophic factor regulates satellite cell differentiation and skeletal muscle regeneration. *Mol Biol Cell* 21, 2182-2190
 111. Miknyoczki SJ, Lang D, Huang L, Klein-Szanto AJ, Dionne CA and Ruggeri BA (1999) Neurotrophins and Trk receptors in human pancreatic ductal adenocarcinoma: expression patterns and effects on in vitro invasive behavior. *Int J Cancer* 81, 417-427
 112. Sinkevicius KW, Kriegel C, Bellaria KJ et al (2014) Neurotrophin receptor TrkB promotes lung adenocarcinoma metastasis. *Proc Natl Acad Sci U S A* 111, 10299-10304
 113. Shen T, Cheng X, Xia C et al (2019) Erlotinib inhibits colon cancer metastasis through inactivation of TrkB-dependent ERK signaling pathway. *J Cell Biochem* 120, 11248-11255
 114. Zhang SY, Hui LP, Li CY, Gao J, Cui ZS and Qiu XS (2016) More expression of BDNF associates with lung squamous cell carcinoma and is critical to the proliferation and invasion of lung cancer cells. *BMC Cancer* 16, 171
 115. Matthews VB, Astrom MB, Chan MH et al (2009) Brain-derived neurotrophic factor is produced by skeletal muscle cells in response to contraction and enhances fat oxidation via activation of AMP-activated protein kinase. *Diabetologia* 52, 1409-1418
 116. Ibeas K, Herrero L, Mera P and Serra D (2021) Hypothalamus-skeletal muscle crosstalk during exercise and its role in metabolism modulation. *Biochem Pharmacol* 190, 114640
 117. Xiao R, Bergin SM, Huang W et al (2016) Environmental and genetic activation of hypothalamic BDNF modulates T-cell immunity to exert an anticancer phenotype. *Cancer Immunol Res* 4, 488-497
 118. Cao L, Liu X, Lin EJ et al (2010) Environmental and genetic activation of a brain-adipocyte BDNF/leptin axis causes cancer remission and inhibition. *Cell* 142, 52-64
 119. Garofalo S, D'Alessandro G, Chece G et al (2015) Enriched environment reduces glioma growth through immune and non-immune mechanisms in mice. *Nat Commun* 6, 6623
 120. Fang P, She Y, Yu M, Min W, Shang W and Zhang Z (2023) Adipose-muscle crosstalk in age-related metabolic disorders: the emerging roles of adipo-myokines. *Ageing Res Rev* 84, 101829
 121. Jena BP, Larsson L, Gatti DL, Ghiran I and Cho WJ (2022) Understanding brain-skeletal muscle crosstalk impacting metabolism and movement. *Discoveries (Craiova)* 10, e144