

ON LACUNARY Δ^m -STATISTICAL CONVERGENCE OF TRIPLE SEQUENCE IN INTUITIONISTIC FUZZY N-NORMED SPACE

ASIF HUSSAIN JAN* AND TANWEER JALAL

ABSTRACT. In this article, we construct lacunary Δ^m -statistical convergence for triple sequences within the context of intuitionistic fuzzy n-normed spaces (IFnNS). For lacunary Δ^m -statistical convergence of triple sequence in IFnNS, we demonstrate numerous results. For this innovative notion of convergence, we further built lacunary Δ^m -statistical Cauchy sequences and offered the Cauchy convergence criterion.

1. Introduction

Fast [11] introduced the idea of statistical convergence, which has since been studied by a large number of authors. Active study on this topic was started after Fridy's publication [12, 13]. In a number of areas, including approximation theory [3], finitely additive set functions [2], sequence space [14, 15], and statistical convergence for fuzzy numbers [1, 21], mathematicians have studied the characteristics of convergence and statistical convergence.

Zadeh [28] gave the concept of fuzziness. It has been one of the most active areas of research in many branches of sciences, with a sizable number of research publications based on the concept of fuzzy sets/numbers appearing in the literature. Intuitionistic fuzzy normed space was first described by Saadati and Park in [22]. In a recent study, R. Antal et al. [1] explored the idea of double sequence Δ -statistical convergence in intuitionistic fuzzy normed space.

There have been numerous studies on difference sequence spaces and related generalisations published in the literature [7–10, 26, 27]. B.C.Tripathy et.al. studied a new type of generalized Difference Cesaro Sequence Spaces [26] and new type of difference sequence spaces [27]. A.Esi studied the generalized difference sequence spaces defined by Orlicz functions [7] and strongly generalized difference $[V^\lambda, \Delta^m, p]$ -summable sequence spaces defined by a sequence of moduli [8]. Later on, several authors studied generalized Δ^m Statistical Convergence in Probabilistic Normed Space [9] and generalized Strongly difference convergent sequences associated with multiplier sequences [10], respectively.

Received October 4, 2022. Revised May 1, 2023. Accepted July 15, 2023.

2010 Mathematics Subject Classification: 40A35, 26E50, 40G15, 46S40.

Key words and phrases: Statistical convergence, Lacunary Δ^m -statistical convergence, Triple Sequence, Intuitionistic fuzzy n-normed space.

* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2023.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

The function $X : \mathbb{N} \times \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{R}(C)$ can be used to define a triple sequence (real or complex), where \mathbb{N}, \mathbb{R} and \mathbb{C} stand for the sets of natural, real, and complex numbers, respectively. At the beginning, Sahiner et al. [23] introduced and studied the many conceptions of triple sequences and their statistical convergence. Triple sequence statistical convergence on probabilistic normed space was recently introduced by Savas and Esi [6], where as statistical convergence of triple sequences in topological groups was later introduced by Esi [5]. For further study on triple sequence spaces, we may refer to [1, 14–16, 18, 19].

Kizmaz [20] introduced the difference sequence space $Z(\Delta)$ as given below

$$Z(\Delta) = \{y = (y_k) : (\Delta y_k) \in Z\}$$

for $Z = \ell_\infty, c, c_0$ i.e. spaces of all bounded, convergent and null sequences respectively, where $\Delta y = (\Delta y_k) = (y_k - y_{k+1})$. In particular, $\ell_\infty(\Delta), c(\Delta)$ and $c_0(\Delta)$ are also Banach spaces, relative to a norm induced by $\|y\|_\Delta = |y_1| + \sup_k |\Delta y_k|$.

The generalized difference sequence spaces $Z(\Delta^m)$ was introduced by M.Et et.al. [4] as follows :

$$Z(\Delta^m) = \{y = (y_k) : (\Delta^m y_k) \in Z\}$$

for $Z = \ell_\infty, c, c_0$ where $\Delta^m(y) == (\Delta^m y_k) = (\Delta_{m-1} y_k - \Delta_{m-1} y_{k+1})$. So that $\Delta^m y_k = \sum_{r=0}^p (-1)^r \binom{m}{r} x_{k+r}$.

The difference operator Δ on triple sequence x_{mnl} is defined as :

$$\Delta x_{mnl} = x_{mnl} - x_{(m+1)nl} - x_{m(n+1)l} - x_{mn(l+1)} = x_{(m+1)(n+1)l} + x_{(m+1)n(l+1)} + x_{m(n+1)(l+1)} - x_{(m+1)(n+1)(l+1)}$$

The generalized difference spaces for triple sequences can be approximated as:

$$Z(\Delta^m) = \{y = (y_{jkl}) : (\Delta^m y_{jkl}) \in Z\}$$

for $Z = \ell_\infty^3, c^3, c_0^3$ where $\Delta^m(y) == (\Delta^m y_{jkl}) = (\Delta_{m-1} y_{jkl} - \Delta_{m-1} y_{jk, (l+1)})$. So that $\Delta^m y_k = \sum_{r=0}^p (-1)^{r+s+u} \binom{m}{r} \binom{m}{s} \binom{m}{u} x_{j+r, k+s, l+u}$.

Here is a summary of the current endeavours. In Section 2, we go over the fundamental definitions of the intuitionistic fuzzy n-normed space. Lacunary Δ^m -statistical convergence in intuitionistic fuzzy n-normed space is presented in Section 3. Here, we established a number of results that show how generalised this convergence process is. For this innovative notion of convergence, we further built Lacunary Δ^m -statistical Cauchy sequences and provided the Cauchy convergence criterion.

2. Definitions and Preliminaries

Here we mention some basic definitions of intuitionistic fuzzy n-normed space and other preliminaries.

DEFINITION 2.1. [24] A continuous t-norm is the mapping $\otimes : [0, 1] \times [0, 1] \rightarrow [0, 1]$ such that

1. \otimes is continuous, associative, commutative and with identity 1 ,
2. $a_1 \otimes b_1 \leq a_2 \otimes b_2$ whenever $a_1 \leq a_2$ and $b_1 \leq b_2, \forall a_1, a_2, b_1, b_2 \in [0, 1]$.

DEFINITION 2.2. [24] A continuous -conorm is the mapping $\odot : [0, 1] \times [0, 1] \rightarrow [0, 1]$ such that

1. \odot is continuous, associative, commutative and with identity 0 ,
2. $a_1 \odot b_1 \leq a_2 \odot b_2$ whenever $a_1 \leq a_2$ and $b_1 \leq b_2, \forall a_1, a_2, b_1, b_2 \in [0, 1]$.

DEFINITION 2.3. [22] An intuitionistic fuzzy normed space (IFNS) is referred to the 5tuple $(X, \varphi, \vartheta, \otimes, \odot)$ with vector space X , fuzzy sets φ, ϑ on $X \times (0, \infty)$, continuous t-norm \otimes and continuous t-conorm \odot , if for each $y, z \in X$ and $s, t > 0$, we have

1. $\varphi(y, t) + \vartheta(y, t) \leq 1$
2. $\varphi(y, t) > 0$ and $\vartheta(y, t) \leq 1$,
3. $\varphi(y, t) = 1$ and $\vartheta(y, t) = 0 \iff y = 0$,
4. $\varphi(\alpha y, t) = \varphi\left(y, \frac{t}{|\alpha|}\right)$ for $\alpha \neq 0$,
5. $\varphi(y, s) \otimes \varphi(z, t) \leq \varphi(y + z, s + t)$ and $\vartheta(y, s) \odot \vartheta(z, t) \leq \vartheta(y + z, s + t)$,
6. $\vartheta(y, o) : (0, \infty) \rightarrow [0, 1]$ and $\vartheta(y, o) : (0, \infty) \rightarrow [0, 1]$ are continuous,
7. $\lim_{t \rightarrow \infty} \varphi(y, t) = 1, \lim_{t \rightarrow 0} \varphi(y, t) = 0, \lim_{t \rightarrow \infty} \vartheta(y, t) = 1$ and $\lim_{t \rightarrow 0} \vartheta(y, t) = 0$.

Then (φ, ϑ) is known as intuitionistic fuzzy norm.

DEFINITION 2.4. [22] Let $(X, \|o\|)$ be any normed space. For every $t > 0$ and $y \in X$, take $\varphi = \frac{t}{t + \|y\|}, \vartheta = \frac{\|y\|}{t + \|y\|}$. Also, $a \otimes b = ab$ and $a \odot b = \min\{a + b, 1\} \forall a, b \in [0, 1]$.

Then, a 5-tuple $(X, \varphi, \vartheta, \otimes, \odot)$ is an IFNS which satisfies the above mentioned conditions.

DEFINITION 2.5. [22] Let $(X, \varphi, \vartheta, \otimes, \odot)$ be an IFNS with norm (φ, ϑ) . A sequence $y = (y_k)$ in X is called convergent to some $\xi \in X$ with respect to the intuitionistic fuzzy norm (φ, ϑ) if there exists $k_0 \in \mathbb{N}$ for each $\epsilon > 0$ and $t > 0$ such that $\varphi(yk - \xi, t) > 1 - \epsilon$ and $\vartheta(yk - \xi, t) < \epsilon$ for all $k \geq k_0$. It is denoted by $(\varphi, \vartheta) - \lim_{k \rightarrow \infty} y_k = \xi$.

DEFINITION 2.6. [22] Let $(X, \varphi, \vartheta, \otimes, \odot)$ be an IFNS with norm (φ, ϑ) . A sequence $y = (y_k)$ in X is called convergent to some $\xi \in X$ with respect to the intuitionistic fuzzy norm (φ, ϑ) if there exists $k_0 \in \mathbb{N}$ for each $\epsilon > 0$ and $t > 0$

$$\delta(\{k \in \mathbb{N} : \varphi(y_k - \xi, t) \leq 1 - \epsilon \quad \text{or} \quad \vartheta(y_k - \xi, t) \geq \epsilon\}) = 0.$$

It is denoted by $S^{\varphi, \vartheta} - \lim_{k \rightarrow \infty} y_k = \xi$.

A subset E of the set \mathbb{N} of natural numbers is said to have a "natural density" $\delta(E)$ if

$$\delta(E) = \lim_n \frac{1}{n} |\{k \leq n : k \in E\}|,$$

where the vertical bars denote the cardinality of the enclosed set.

The number sequence $x = (x_k)$ is said to be statistically convergent to number l if for each $\epsilon > 0$,

$$\lim_n \frac{1}{n} |\{k \leq n : |x_k - l| \geq \epsilon\}| = 0,$$

and x is said to be statistically cauchy sequence if for every $\epsilon > 0$ there exists a number $N = N(\epsilon)$ such that

$$\lim_n \frac{1}{n} |\{k \leq n : |x_k - x_N| \geq \epsilon\}| = 0.$$

DEFINITION 2.7. [22] Let $(X, \varphi, \vartheta, \otimes, \odot)$ be an IFNS with norm (φ, ϑ) . A double sequence $y = (y_{jk})$ in X is called statistically convergent to some $\xi \in X$ with respect to the intuitionistic fuzzy norm (φ, ϑ) if there exists $k_0 \in \mathbb{N}$ for each $\epsilon > 0$ and $t > 0$

$$\delta(\{k \in \mathbb{N} : \varphi(y_{jk} - \xi, t) \leq 1 - \epsilon \text{ or } \vartheta(y_{jk} - \xi, t) \geq \epsilon\}) = 0.$$

It is denoted by $S(\varphi, \vartheta) - \lim_{k \rightarrow \infty} y_{jk} = \xi$.

DEFINITION 2.8. [6] The triple sequence $\theta_{j,k,l} = \{(j_r, k_s, l_t)\}$ is called the triple lacunary sequence if there exist three increasing sequences of integers such that

$$j_o = 0, h_r = j_r - j_{r-1} \rightarrow \infty \text{ as } r \rightarrow \infty, \\ k_o = 0, h_s = k_s - k_{s-1} \rightarrow \infty \text{ as } s \rightarrow \infty,$$

and

$$I_o = 0, h_t = I_t - I_{t-1} \rightarrow \infty \text{ as } t \rightarrow \infty$$

Let $k_{r,s,t} = j_r k_s I_t, h_{r,s,t} = h_r h_s h_t$ and $\theta_{j,k,l}$ is determined by

$$I_{r,s,t} = \{(j, k, l) : j_{r-1} < j \leq j_r, k_{s-1} < k \leq k_s \text{ and } I_{t-1} < I \leq I_t\} \\ q_r = \frac{j_r}{j_{r-1}}, q_s = \frac{k_s}{k_{s-1}}, q_t = \frac{l_t}{l_{t-1}} \text{ and } q_{r,s,t} = q_r q_s q_t.$$

Let $K \subset \mathbb{N} \times \mathbb{N} \times \mathbb{N}$. The number

$$\delta_3^\theta = \lim_{r,s,t} \frac{1}{h_{r,s,t}} |\{(j, k, l) \in I_{r,s,t} : (j, k, l) \in K\}|$$

is said to be the $\theta_{r,s,t}$ -density of K , provided the limit exists.

Below is the definition of n-mored space:

DEFINITION 2.9. Let $n \in \mathbb{N}$ and X be a real vector space of dimension $d \geq n$. (Here we allow d to be infinite). A real-valued function $\|, \dots, \|$ on X^n satisfying the following four properties:

- (1) $\|f_1, f_2, \dots, f_n\| = 0$ if and only if f_1, f_2, \dots, f_n are linearly dependent;
- (2) $\|f_1, f_2, \dots, f_n\|$ is invariant under permutation;
- (3) $\|f_1, f_2, \dots, f_{n-1}, \alpha f_n\| = |\alpha| \|f_1, f_2, \dots, f_{n-1}, f_n\|$ for any $\alpha \in R$;
- (4) $\|f_1, f_2, \dots, f_{n-1}, y + z\| \leq \|f_1, f_2, \dots, f_{n-1}, y + f_1, f_2, \dots, f_{n-1}, z\|$, is called an n-norm on X and the pair $(X, \|f_1, f_2, \dots, f_n\|)$ is called an n-normed space.

DEFINITION 2.10. [25] An IFnNLS is the five-tuple $(X, \mu, v, *, \circ)$ where X is a linear space over a field $F, *$ is a continuous t -norm, \circ is a continuous t -conorm, μ, v are fuzzy sets on $X^n \times (0, \infty), \mu$ denotes the degree of membership and v denotes the degree of nonmembership of $(x_1, x_2, \dots, x_n, t) \in X^n \times (0, \infty)$ satisfying the following conditions for every $(x_1, x_2, \dots, x_n) \in X^n$ and $s, t > 0$:

- (i) $\mu(x_1, x_2, \dots, x_n, t) + v(x_1, x_2, \dots, x_n, t) \leq 1,$
- (ii) $\mu(x_1, x_2, \dots, x_n, t) > 0,$
- (iii) $\mu(x_1, x_2, \dots, x_n, t) = 1$ if and only if x_1, x_2, \dots, x_n are linearly dependent,
- (iv) $\mu(x_1, x_2, \dots, x_n, t)$ is invariant under any permutation of $x_1, x_2, \dots, x_n,$
- (v) $\mu(x_1, x_2, \dots, cx_n, t) = \mu(x_1, x_2, \dots, x_n, \frac{t}{|c|})$ for all $c \neq 0, c \in F,$
- (vi) $\mu(x_1, x_2, \dots, x_n, s) * \mu(x_1, x_2, \dots, x'_n, t) \leq \mu(x_1, x_2, \dots, x_n + x'_n, s + t),$
- (vii) $\mu(x_1, x_2, \dots, x_n, t) : (0, \infty) \rightarrow [0, 1]$ is continuous in $t,$
- (viii) $\lim_{t \rightarrow \infty} \mu(x_1, x_2, \dots, x_n, t) = 1$ and $\lim_{t \rightarrow 0} \mu(x_1, x_2, \dots, x_n, t) = 0,$
- (ix) $v(x_1, x_2, \dots, x_n, t) < 1$

- (x) $v(x_1, x_2, \dots, x_n, t) = 0$ if and only if x_1, x_2, \dots, x_n are linearly dependent,
- (xi) $v(x_1, x_2, \dots, x_n, t)$ is invariant under any permutation of x_1, x_2, \dots, x_n ,
- (xii) $v(x_1, x_2, \dots, cx_n, t) = v\left(x_1, x_2, \dots, x_n, \frac{t}{|c|}\right)$ for all $c \neq 0, c \in F$,
- (xiii) $v(x_1, x_2, \dots, x_n, s) \circ v(x_1, x_2, \dots, x'_n, t) \geq v(x_1, x_2, \dots, x_n + x'_n, s + t)$
- (xiv) $v(x_1, x_2, \dots, x_n, t) : (0, \infty) \rightarrow [0, 1]$ is continuous in t ,
- (xv) $\lim_{t \rightarrow \infty} v(x_1, x_2, \dots, x_n, t) = 0$ and $\lim_{t \rightarrow 0} v(x_1, x_2, \dots, x_n, t) = 1$.

EXAMPLE 2.11. [25] Let $(X, \|\bullet, \dots, \bullet\|)$ be an n -normed linear space. Also let $a * b = ab$ and $a \circ b = \min\{a + b, 1\}$ for all $a, b \in [0, 1]$

$$\mu(x_1, x_2, \dots, x_n, t) = \frac{t}{t + \|x_1, x_2, \dots, x_n\|} \quad \text{and} \quad v(x_1, x_2, \dots, x_n, t) = \frac{\|x_1, x_2, \dots, x_n\|}{t + \|x_1, x_2, \dots, x_n\|}.$$

Then $(X, \mu, v, *, \circ)$ is an IFnNLS.

3. Triple Lacunary Δ^m -statistical convergence in IFnNS.

In the context of intuitionistic fuzzy normed spaces for triple sequences, we define Lacunary Δ^m -statistical convergence and establish certain results.

DEFINITION 3.1. Let $(X, \varphi, \vartheta, \otimes, \odot)$ be a IFnNS with norm $(\varphi, \vartheta)^n$ and $\theta_{j,k,l}$ be a triple lacunary sequence. A triple sequence $y = (y_{jkl})$ in X is called lacunary Δ^m -statistically convergent to some $\xi \in X$ with respect to the intuitionistic fuzzy norm (φ, ϑ) if for each $\epsilon > 0, t > 0$ and $f_1, f_2, \dots, f_{n-1} \in X$.

$$(1) \quad \delta_3^\theta(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq 1 - \epsilon \text{ or } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq \epsilon\}) = 0$$

or equivalently

$$(1^*) \quad \delta_3^\theta(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) > 1 - \epsilon \text{ or } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) < \epsilon\}) = 1.$$

In this case, we write $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$ or $X_{jkl} \xrightarrow{(\varphi, \vartheta)^n} \xi (S_{\theta_{j,k,l}})$ and denote the set of all $S_{\theta_{j,k,l}}$ -convergent triple sequences in the intuitionistic fuzzy normed space by $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n}$.

DEFINITION 3.2. Let $(X, \varphi, \vartheta, \otimes, \odot)$ be a IFnNS with norm $(\varphi, \vartheta)^n$ and $\theta_{j,k,l}$ be a triple lacunary sequence. A triple sequence $y = (y_{jkl})$ in X is called lacunary Δ^m -statistically Cauchy with respect to the intuitionistic fuzzy norm (φ, ϑ) if there exists $j_0, k_0, l_0 \in \mathbb{N}$ for each $\epsilon > 0$ and $t > 0$ such that for all $j, r \geq j_0, k, s \geq k_0, l, u \geq l_0$ and $f_1, f_2, \dots, f_{n-1} \in X$, we have

$$\delta_3^\theta(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{rsu}, t) \leq 1 - \epsilon \text{ or } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{rsu}, t) \geq \epsilon\}) = 0.$$

It is denoted by $S_{jkl}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$.

From (1) and (1*), we have the following lemma.

Lemma 3.1. Let $(X, \varphi, \vartheta, \otimes, \odot)$ be an IFNS with norm $(\varphi, \vartheta)^n$ and $\theta_{j,k,l}$ be a triple lacunary sequence. Then the following statements are equivalent for triple sequence $y = (y_{jkl})$ in X whenever $\epsilon > 0$ and $t > 0$,

- (i) $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$,
- (ii) $\delta_3^\theta(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) > 1 - \epsilon\}) = 1$,
 $= \delta_3^\theta(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) < \epsilon\}) = 1$,
- (iii) $\delta_3^\theta(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq 1 - \epsilon\}) = 0$,
 $= \delta_3^\theta(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq \epsilon\}) = 0$,
- (iv) $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) = 1$
 and $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) = 0$.

THEOREM 3.3. Let $(X, \varphi, \vartheta, \otimes, \odot)$ be a IFNS with norm $(\varphi, \vartheta)^n$ and $\theta_{j,k,l}$ be a triple lacunary sequence. If $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$, then ξ is unique.

Proof. If possible, let $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi_1$ and $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi_2$.

For given $\epsilon \in (0, 1)$ and $t > 0$, take $\alpha > 0$ such that $(1 - \alpha) \otimes (1 - \alpha) > 1 - \epsilon$ and $\alpha \odot \alpha < \epsilon$.

Consider

$$\begin{aligned} K_{1,\varphi}(\alpha, t) &= \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi_1, t/2) \leq 1 - \alpha\}, \\ K_{2,\varphi}(\alpha, t) &= \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi_2, t/2) \leq 1 - \alpha\}, \\ K_{3,\vartheta}(\alpha, t) &= \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi_1, t/2) \geq \alpha\}, \\ K_{4,\vartheta}(\alpha, t) &= \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi_2, t/2) \geq \alpha\}. \end{aligned}$$

Using lemma 3.1, we have

$$\begin{aligned} \delta_3^\theta(K_{1,\varphi}(\alpha, t)) &= \delta_3^\theta(K_{3,\vartheta}(\alpha, t)) = 0. \\ \delta_3^\theta(K_{2,\varphi}(\alpha, t)) &= \delta_3^\theta(K_{4,\vartheta}(\alpha, t)) = 0. \end{aligned}$$

Let $K_{\varphi, \vartheta}(\alpha, t) = [K_{1,\varphi}(\alpha, t) \cup K_{2,\varphi}(\alpha, t)] \cap [K_{3,\vartheta}(\alpha, t) \cup K_{4,\vartheta}(\alpha, t)]$. Clearly,

$$\delta_3^\theta K_{\varphi, \vartheta}(\alpha, t) = 0.$$

Whenever $(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} - K_{\varphi, \vartheta}(\alpha, t)$, we have two possibilities, either $(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} - [K_{1,\varphi}(\alpha, t) \cup K_{2,\varphi}(\alpha, t)]$ or $(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} - [K_{3,\vartheta}(\alpha, t) \cup K_{4,\vartheta}(\alpha, t)]$.

First, we consider $(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} - [K_{1,\varphi}(\alpha, t) \cup K_{2,\varphi}(\alpha, t)]$. Then

$$\begin{aligned} \varphi(\xi_1 - \xi_2, t) &\geq \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi_1, t/2) \otimes \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi_2, t/2) \\ &> (1 - \alpha) \otimes (1 - \alpha) \\ &> 1 - \epsilon. \end{aligned}$$

As given $\epsilon \in (0, 1)$ was arbitrary, then $\varphi(\xi_1 - \xi_2, t) = 1$ for all $t > 0$, then $\xi_1 = \xi_2$.

Similarly, if $(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} - [K_{3,\vartheta}(\alpha, t) \cup K_{4,\vartheta}(\alpha, t)]$

$$\begin{aligned} \vartheta(\xi_1 - \xi_2, t) &\leq \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi_1, t/2) \odot \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi_2, t/2) \\ &< \alpha \odot \alpha \\ &< \epsilon. \end{aligned}$$

Since $\epsilon \in (0, 1)$ was arbitrary, then $\varphi(\xi_1, \xi_2, t) = 0$ for all $t > 0$, i.e., $\xi_1 = \xi_2$. Therefore $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$ exists uniquely. □

THEOREM 3.4. *Let $(X, \varphi, \vartheta, \otimes, \odot)$ be a IFNS with norm $(\varphi, \vartheta)^n$ and $\theta_{j,k,l}$ be a triple lacunary sequence. If $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$, then $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. But converse may not be true.*

Proof. Let $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. Then, there exists j_0, k_0 and $l_0 \in \mathbb{N}$ for given $\epsilon > 0$ and any $t > 0$ such that for all $j \geq j_0, k \geq k_0$ and $l \geq l_0$, we have $\varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) > 1 - \epsilon$ and $\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) < \epsilon$.

Further, the set $A(\epsilon, t) = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq 1 - \epsilon$ or $\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq \epsilon\}$, contains only finite number of elements. We know that natural density of any finite set is always zero. Therefore, $\delta_3^\theta(A(\epsilon, t)) = 0$ i.e. $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. □

But converse is not true, this can be justified with the example.

EXAMPLE 3.5. Let $(\mathbb{R}, ||)$ be the real normed space under the usual norm. Define $a \otimes b = ab$ and $a \odot b = \min\{a + b, 1\} \forall a, b \in [0, 1]$. Also for every $t > 0$ and all $y \in \mathbb{R}$, consider $\varphi(y, t) = \frac{t}{t+|y|}$ and $\vartheta(y, t) = \frac{|y|}{t+|y|}$. Then, clearly $(\mathbb{R}, \varphi, \vartheta, \otimes, \odot)$ is an IFNS. Define the sequence

$$\Delta^m x_{jkl} = \begin{cases} jkl, & \text{for } j_r - \lceil \sqrt{h_r} \rceil + 1 \leq j \leq j_r \\ k_s - \lceil \sqrt{h_s} \rceil + 1 \leq k \leq k_s \\ \text{and } l_t - \lceil \sqrt{h_t} \rceil + 1 \leq l \leq l_t \\ \xi, & \text{otherwise.} \end{cases}$$

By given $\epsilon > 0$ and $t > 0$, we obtain the below set for $\xi = 0$.

$$\begin{aligned} K(\epsilon, t) &= \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl}, t) \leq 1 - \epsilon \\ &\text{or } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl}, t) \geq \epsilon\} \\ &= \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : |\Delta^m y_{jkl}| \geq \frac{\epsilon t}{1 - \epsilon} > 0 \\ &= \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : |\Delta^m y_{jkl}| = jkl \\ &= \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : |\Delta^m y_{jkl}| \geq \frac{\epsilon t}{1 - \epsilon} > 0 \\ &= \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : j_r - \lceil \sqrt{h_r} \rceil + 1 \leq j \leq j, \\ &\quad k_s - \lceil \sqrt{h_s} \rceil + 1 \leq k \leq k_s \\ &\text{and } l_t - \lceil \sqrt{h_t} \rceil + 1 \leq l \leq l_t\} \end{aligned}$$

and so, we get

$$\lim_{r,s,t} \frac{1}{h_r, h_s, h_t} \mid \left\{ (j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : j_r - \lceil \sqrt{h_r} \rceil + 1 \leq j \leq j_r \right.$$

$$k_s - \lceil \sqrt{h_s} \rceil + 1 \leq k \leq k_s$$

$$\text{and } l_t - \lceil \sqrt{h_t} \rceil + 1 \leq l \leq l_t \left. \right\}$$

$$\leq \lim_{r,s,t} \frac{\sqrt{h_r} \sqrt{h_s} \sqrt{h_t}}{h_r h_s h_t} = 0.$$

Hence $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = 0$. By the above defined sequence $(\Delta^m y_{jkl})$, we get

$$\varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m x_{jkl}, t) = \begin{cases} \frac{t}{t+|jkl|}, & \text{for } r - \lceil \sqrt{h_r} \rceil + 1 \leq j \leq j_r \\ & k_s - \lceil \sqrt{h_s} \rceil + 1 \leq k \leq k_s \\ & \text{and } l_t - \lceil \sqrt{h_t} \rceil + 1 \leq l \leq l_t \\ 0, & \text{otherwise.} \end{cases}$$

i.e $\varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m x_{jkl}, t) \leq 1, \quad \forall j, k, l$. And

$$\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m x_{jkl}, t) = \begin{cases} \frac{|jkl|}{t+|jkl|}, & \text{for } j_r - \lceil \sqrt{h_r} \rceil + 1 \leq j \leq j_r, \\ & k_s - \lceil \sqrt{h_s} \rceil + 1 \leq k \leq k_s \\ & \text{and } l_t - \lceil \sqrt{h_t} \rceil + 1 \leq l \leq l_t \\ 0, & \text{otherwise.} \end{cases}$$

i.e $\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m x_{jkl}, t) \geq 0, \quad \forall j, k, l$.

This shows that $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} \neq 0$.

THEOREM 3.6. Let $(X, \varphi, \vartheta, \otimes, \odot)$ be a IFnNS with norm $(\varphi, \vartheta)^n$ and $\theta_{j,k,l}$ be a triple lacunary sequence. Then $S_{\theta_{j,k,l}, \vartheta} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi \iff$ there exists a set $P = \{(j_a, k_b, l_c) : a, b, c = 1, 2, 3, \dots\} \subset \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ such that $\delta(P) = 1$ and $(\varphi, \vartheta)^n - \lim_{j_a, k_b, l_c \rightarrow \infty} \Delta^m y_{j_a k_b l_c} = \xi$.

Proof. Assume that $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. For $t > 0$ and $\alpha \in \mathbb{N}$, we take $M(\alpha, t) = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) > 1 - 1/\alpha \text{ and } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) < 1/\alpha\}$, and

$$K(\alpha, t) = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq 1 - 1/\alpha$$

or

$$\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq 1/\alpha\},$$

as $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$, then $\delta_3(K(\alpha, t)) = 0$. Also, for any $t > 0$ and $\alpha \in \mathbb{N}$, evidently we get $M(\alpha, t) \supset M(\alpha + 1, t)$, and

$$(3.1) \quad \delta_3(M(\alpha, t)) = 1,$$

For $(j, k, l) \in M(\alpha, t)$, we prove $(\varphi, \vartheta)^n - \lim_{j_a, k_b, l_c \rightarrow \infty} \Delta^m y_{j_a k_b l_c} = \xi$.

On the contrary, suppose that triple sequence $y = (y_{jkl})$ is not Δ^m -convergent to ξ for all $(j, k, l) \in M(\alpha, t)$. So, there exists some $\alpha > 0$ and $k_0 \in \mathbb{N}$ such that

$$\begin{aligned} &\varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq 1 - \rho \\ &\text{or } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq \rho \} \text{ for all } j, k, l \geq k_0 \\ &\implies \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq 1 - \rho \text{ and } \vartheta(\Delta^m y_{jkl} - \xi, t) \leq \rho \} \text{ for all } j, k, l \geq k_0 \end{aligned}$$

Therefore, $\delta_3(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq 1 - \rho \text{ and } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq \rho\}) = 0$
 i.e. $\delta_3(M(\rho, t)) = 0$. Since $\rho > 1/\alpha$, then $\delta_3(M(\alpha, t)) = 0$ as $M(\alpha, t) \subset M(\rho, t)$, which is a contradiction to (3.1). This shows that there exists a set $M(\alpha, t)$ for which $\delta_3(M(\alpha, t)) = 1$ and the triple sequence $y = (y_{jkl})$ is statistically Δ^m -convergent to ξ .

Conversely, suppose there exists a subset $P = \{(j_a, k_b, l_c) : a, b, c = 1, 2, 3, \dots\} \subseteq \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ with $\delta_3(P) = 1$ and $(\varphi, \vartheta)^n - \lim_{j_a, k_b, l_c \rightarrow \infty} \Delta^m y_{j_a k_b l_c} = \xi$. i.e. for given $\rho > 0$ and any $t > 0$ we have $N_0 \in \mathbb{N}$, which gives

$$\varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) > 1 - \rho$$

and

$$\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) < \rho \text{ for all } j, k, l \geq N_0.$$

Now, let $K(\rho, t) = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq 1 - \rho \text{ or } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq \rho\}$. Then,

$$K(\rho, t) \subseteq \mathbb{N} - \{(j_{N_0+1}, k_{N_0+1}, l_{N_0+1}), \dots\}. \text{ As } \delta_3(P) = 1 \implies \delta_3(K(\alpha, t)) \leq 0.$$

Hence, $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. □

THEOREM 3.7. Let $(X, \varphi, \vartheta, \otimes, \odot)$ be a IFNS with norm $(\varphi, \vartheta)^n$ and $\theta_{j,k,l}$ be a triple lacunary sequence. Let $y = (y_{jkl})$ be any triple sequence. Then $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi \iff$ there is a triple sequence $x = (x_{jkl})$ such that $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$ and $\delta_3(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \Delta^m y_{jkl} = \Delta^m x_{jkl}\}) = 1$

Proof. Assume that $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. By Theorem (3.3), we set

$$P = \{(j_a, k_b, l_c) : a, b, c = 1, 2, 3, \dots\} \subseteq \mathbb{N} \times \mathbb{N} \times \mathbb{N}$$

with $\delta_3(P) = 1$ and $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{j_a k_b l_c} = \xi$. Consider the sequence

$$\Delta^m x_{jkl} = \begin{cases} \Delta^m y_{jkl}, & (j, k, l) \in P \\ \xi, & \text{otherwise,} \end{cases}$$

which gives the required result.

Conversely, consider $x = (x_{jkl})$ and $z = (z_{jkl})$ in X with $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$ and $\delta_3(\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \Delta^m y_{jkl} = \Delta^m x_{jkl}\}) = 1$. Then for each $\epsilon > 0$ and $t > 0$,

$$\{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq 1 - \epsilon$$

or

$$\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq \epsilon\} \subseteq A \cup B$$

where

$$A = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq 1 - \epsilon\}$$

or

$$\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq \epsilon\}$$

and

$$B = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : (\Delta^m y_{jkl} \neq \Delta^m x_{jkl})\}.$$

Since $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{j_a k_b l_c} = \xi$ then the set A contains at most finitely many terms. Also $\delta_3(B) = 0$ as $\delta_3(B^c) = 1$ where $B^c = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \Delta^m y_{jkl} = \Delta^m x_{jkl}\}$.

Therefore

$$\delta_3 \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \leq 1 - \epsilon\}$$

or

$$\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t) \geq \epsilon\}.$$

We get $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. □

THEOREM 3.8. *Let $(X, \varphi, \vartheta, \otimes, \odot)$ be a IFNS with norm $(\varphi, \vartheta)^n$ and $\theta_{j,k,l}$ be a triple lacunary sequence. Let $y = (y_{jkl})$ be any triple sequence. Then $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi \iff$ there exists two triple sequence $z = (z_{jkl})$ and $x = (x_{jkl})$ in X such that $\Delta^m y_{jkl} = \Delta^m z_{jkl} + \Delta^m x_{jkl}$ for all $(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ where $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{j_a k_b l_c} = \xi$ and $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$.*

Proof. Assume that $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. By Theorem (3.5), we set $P = \{(j_a, k_b, l_c) : a, b, c = 1, 2, 3, \dots\} \subseteq \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ with $\delta_3(P) = 1$ and $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{j_a k_b l_c} = \xi$.

Consider two triple sequences $z = (z_{jkl})$ and $x = (x_{jkl})$, then

$$\Delta^m z_{jkl} = \begin{cases} \Delta^m y_{jkl}, & (j, k, l) \in P \\ \xi, & \text{otherwise.} \end{cases}$$

and

$$\Delta^m x_{jkl} = \begin{cases} 0, & (j, k, l) \in P \\ \Delta^m y_{jkl} - \xi, & \text{otherwise,} \end{cases}$$

which gives the required result.

Conversely, consider $x = (x_{jkl})$ and $z = (z_{jkl})$ in X with $\Delta^m y_{jkl} = \Delta^m z_{jkl} + \Delta^m x_{jkl}$ for all $(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ where $(\varphi, \vartheta)^n - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$ and $S_{\theta_{j,k,l}}^{(\varphi, \vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. Then we get result using Theorem (3.6) and Theorem (3.7). □

THEOREM 3.9. *A triple sequence $y = (y_{jkl})$ in IFNS $(X, \varphi, \vartheta, \otimes, \odot)$ is lacunary Δ^m -statistically convergent with respect to $(\varphi, \vartheta)^n$ if and only if it is lacunary Δ^m -statistically Cauchy with respect to $(\varphi, \vartheta)^n$.*

Proof. Let $S_{\theta_{j,k,l}}^{(\varphi,\vartheta)^n} - \lim_{j,k,l \rightarrow \infty} \Delta^m y_{jkl} = \xi$. Then, for each $\epsilon > 0$ and $t > 0$, take $\alpha > 0$ such that $(1 - \alpha) \otimes (1 - \alpha) > 1 - \epsilon$ and $\alpha \odot \alpha < \epsilon$.

Let $K(\alpha, t) = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \leq 1 - \alpha \text{ or } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \geq \alpha\}$, therefore $\delta_3(K(\alpha, t)) = 0$ and $\delta_3([K(\alpha, t)]^c) = 1$.

Let $M(\epsilon, t) = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{rsu}, t) \leq 1 - \epsilon \text{ or } \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{rsu}, t) \geq \epsilon\}$.

Now, we prove $M(\epsilon, t) = K(\epsilon, t)$, for this if $(j, k, l) \in M(\epsilon, t) = K(\epsilon, t)$. Then we get $\varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \leq 1 - \alpha$ or $\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \geq \alpha$.

Also

$$\begin{aligned} & 1 - \epsilon \\ & \geq \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{rsu}, t) \\ & \geq \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \otimes \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \\ & > (1 - \alpha) \otimes (1 - \alpha) \\ & > 1 - \epsilon \end{aligned}$$

and

$$\begin{aligned} \epsilon & \geq \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{rsu}, t) \\ & \leq \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \odot \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \\ & < \alpha \odot \alpha \\ & < \epsilon. \end{aligned}$$

which is not possible. Therefore $M(\epsilon, t) \subset K(\alpha, t)$ and $\delta_3(M(\epsilon, t)) = 0$ i.e. $y = (y_{jkl})$ is Δ^m -statistically convergent with respect to (φ, ϑ) .

Coversely, assume that $y = (y_{jkl})$ is Δ^m -stastically Cauchy with respect to $(\varphi, \vartheta)^n$ but not Δ^m -stastically convergent with respect to $(\varphi, \vartheta)^n$. Thus for $\epsilon > 0$ and $t > 0$, $\delta_3(M(\epsilon, t)) = 0$, where

$$M(\epsilon, t) = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{j_0 k_0 l_0}, t) \leq 1 - \epsilon$$

or

$$\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{j_0 k_0 l_0}, t) \geq \epsilon\}.$$

Take $\alpha > 0$ such that $(1 - \alpha) \otimes (1 - \alpha) > 1 - \epsilon$ and $\alpha \odot \alpha < \epsilon$. Also, $\delta_3(K(\alpha, t)) = 0$, where $K(\alpha, t) = \{(j, k, l) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N} : \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \geq 1 - \alpha$

or $\vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) < \epsilon\}$. Now

$$\begin{aligned} & \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{j_0 k_0 l_0}, t) \\ & \geq \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \otimes \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{j_0 k_0 l_0} - \xi, t/2) \\ & > (1 - \alpha) \otimes (1 - \alpha) \\ & > 1 - \epsilon \end{aligned}$$

and

$$\begin{aligned} & \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \Delta^m y_{j_0 k_0 l_0}, t) \\ & \leq \vartheta(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{jkl} - \xi, t/2) \odot \varphi(f_1, f_2, \dots, f_{n-1}, \Delta^m y_{j_0 k_0 l_0} - \xi, t/2) \\ & < \alpha \odot \alpha \\ & < \epsilon. \end{aligned}$$

Therefore, $\delta_3([M(\epsilon, t)]^c) = 0$ i.e. $\delta_3(M(\epsilon, t)) = 1$, which is a contradiction as $y = (y_{jkl})$ is Δ^m -statistically Cauchy. Hence, $y = (y_{jkl})$ is Δ^m -statistically convergent with respect to $(\varphi, \vartheta)^n$. \square

4. Conclusion.

This work establishes certain conclusions and defines Lacunary Δ^m -statistical convergence on intuitionistic fuzzy n -normed space. Since any regular norm implies an intuitionistic fuzzy norm, the findings are more pervasive than in analogous normed spaces.

5. Declaration

Conflicts of interests: There is no conflict of interest.

Availability of data and materials: This paper has no associated data.

References

- [1] R.Antal, V.Kumar and B.Hazarika, *On Δ^m -statistical convergence double sequences in intuitionistic fuzzy normed spaces*, *Proyecciones journal of mathematics* **41** (3) (2022), 697–713.
- [2] J.Connor and J.Kline, *On statistical limit points and the consistency of statistical convergence*, *Journal of mathematical analysis and applications* **197** (2) (1996), 392–399.
- [3] O.Duman, M.K.Khan and C.Orhan, *A-statistical convergence of approximating operators*, *Mathematical inequalities and applications* **6** (2003), 689–700.
- [4] M.Et and R.Colak, *On some generalized difference difference sequence spaces*, *Soochow journal of mathematical* **24** (4) (1995), 377–386.
- [5] A.Esi, *Statistical convergence of triple sequences in topological groups*, *Annals of the University of Craiova-Mathematics and Computer Science Series* **40** (1) (2013), 29–33.
- [6] A.Esi and E.Savas, *On Lacunary Statistically Convergent Triple Sequences in Probabilistic Normed Space*, *Applied mathematics and information sciences* **5** (2015), 2529–2535.
- [7] A.Esi, *Generalized difference sequence spaces defined by Orlicz functions*, *General Mathematics* **17** (2) (2009), 53–66.
- [8] A.Esi, *Strongly generalized difference $[V^\lambda, \Delta^m, p]$ -summable sequence spaces defined by a sequence of moduli*, *Nihonkai Mathematical Journal* **20** (2) (2009), 99–108.
- [9] A.Esi and K.Ozdemir, *Generalized Δ^m -statistical convergence in Probabilistic Normed Space*, *Journal of Computational Analysis and Applications* **13** (5) (2011), 923–932.
- [10] A.Esi and B.C.Tripathy, *Generalized Strongly difference convergent sequences associated with multiplier sequences*, *Mathematica Slovaca* **57** (4) (2007), 339–348.
- [11] H.Fast, *Sur la convergence statistique*, *Colloquium Mathematicum* **2** (1951), 241–244.
- [12] J.A.Fridy and M.K.Khan, *Tauberian theorems via statistical convergence*, *Journal of mathematical analysis and applications* **228** (1) (1998), 73–95.
- [13] J.A.Fridy and C.Orhan, *Lacunary statistical summability*, *Journal of mathematical analysis and applications* **173** (2) (1993), 497–504.
- [14] T.Jalal and I.A.Malik, *I-convergent triple sequence spaces over n -normed space*, *Asia pacific journal of mathematics* **5** (2) (2018), 233–242.
- [15] T.Jalal and I.A.Malik, *Some new triple sequence spaces over n -normed space*, *Proyecciones (Antofagasta)* **37** (3) (2018), 547–564.
- [16] T.Jalal and I.A.Malik, *I-Convergence of triple difference sequence spaces over n -normed space*, *Tbilisi mathematical journal* **11** (4) (2018), 93–102

- [17] T.Jalal and I.A.Malik, *Topological and Algebraic Properties of Triple n-normed Spaces*, Proceedings of the fifth international conference on mathematics and computing, Springer Singapore (2021),187–196.
- [18] V.A.Khan, M.I.Idrisi and M.Ahmad, *On I-convergent triple sequence spaces defined by a compact operator and an Orlicz function*, TWMS Journal of Applied and Engineering Mathematics **2** (4) (2021), 1012–1022.
- [19] V.A.Khan, M.I.Idrisi and U.Tuba, *On ideal convergence of triple sequences in intuitionistic Fuzzy normed space defined by compact operator*, Proyecciones (Antofagasta) **40** (5) (2021), 1227–1247.
- [20] H.Kizmaz, *On certain sequence vspaces*, Canadian mathematical bulletin **24** (2) (1981), 169–176.
- [21] M.Mursaleen and S.A.Mohiuddine, *Statistical convergence of double sequences in intuitionistic fuzzy normed spaces*, Chaos, solitons and fractals **41** (5) (2009), 2414–2421.
- [22] R.Saadati and J.H.Park, *On the intuitionistic fuzzy topological spaces*, Chaos, solitons and fractals **27** (2) (2006), 331–344.
- [23] A.Sahiner, M.Gurdal and F.K.Duden, *Triple sequences and their statistical convergence*, Türkiye klinikleri psikiyatri dergisi **8** (2) (2007), 49–55.
- [24] B.Schweizer and A.Sklar, *Statistical Metric Spaces*, Pacific journal of mathematics **10** (1) (1960), 313–334.
- [25] M.Sen and P.Debnath, *Lacunary statistical convergence in intuitionistic fuzzy n-normed spaces*, Math. Comput. Model **54** (2011), 2978–2985.
- [26] B.C.Tripathy, A.Esi and B.Tripathy, *On a new type of generalized difference cesaro sequence spaces*, Soochow Journal of Mathematics **31** (3) (2005), 333–340.
- [27] B.C.Tripathy and A.Esi, *A new type difference sequence spaces*, International Journal of Science and Technology **1** (1) (2006), 11–14.
- [28] L.A.Zadeh, *Fuzzy sets*, Information and control **8** (3) (1965), 338–353.

Asif Hussain Jan

Department of Mathematics, National Institute of Technology,
 Hazratbal, Srinagar -190006, Jammu and Kashmir, India.
E-mail: asif_06phd20@nitsri.net

Tanweer Jalal

Department of Mathematics, National Institute of Technology,
 Hazratbal, Srinagar -190006, Jammu and Kashmir, India.
E-mail: tjalal@nitsri.ac.in