

## APPROXIMATE BEST PROXIMITY PAIR RESULTS ON METRIC SPACES USING CONTRACTION OPERATORS

R. THEIVARAMAN<sup>\*,†</sup>, P. S. SRINIVASAN, AND A. HERMINAU JOTHY

ABSTRACT. The aim of this paper is to prove some new approximate best proximity pair theorems on metric spaces using contraction mappings such as  $P$ -Bianchini contraction,  $P-B$  contraction and so on. A few examples are provided to exemplify our findings. Finally, we discuss some applications that are related to the main results.

### 1. Introduction

Best proximity point theory and fixed point theory are now crucial in many mathematics-related fields and its applications, notably in economics, astronomy, dynamical systems, decision theory, and parameter estimation. In 1922 [2], Banach proposed the Banach fixed point results. After that, various authors extended these principle and gave many results using contractive mappings on metric spaces (refer, [8], [9], [15], [19], [20], [21], [30] & [37]). In addition to that, many researchers found new approximate fixed point theorems on metric spaces that do not require completeness in both contractive and rational contractive operators (refer, [4], [5], [6], [10], [11], [18], [27] & [32]). On the other hand, the best proximity point theory also has the same importance as fixed point theory. In the absence of exact proximity points, approximate best proximity points may be used because the best proximity point results have overly strict limitations. There seem to be numerous problems in applied mathematics that can be handled using the concept of best proximity pair theory. Nonetheless, experience demonstrates that for many instances, an approximate computation is more than acceptable; hence, having the best proximity pair is not always necessary, but having an almost-best proximity pair is essential. Another type of growing challenge that leads to this approximate occurs when the requirements that must be enforced to ensure the presence of the best proximity pairings for the major challenge at hand are much more stringent. In [24], the authors achieved some results on the optimum proximity pairs. In the same way, the authors Antony

---

Received June 15, 2023. Revised September 7, 2023. Accepted September 20, 2023.

2010 Mathematics Subject Classification: 47H10, 54H25.

Key words and phrases: best approximate pairs,  $P-B$  contraction,  $P$ -Bianchini contraction,  $P$ -Reich contraction.

† This work is supported by Bharathidasan University, India, for its financial support under the URF scheme.

\* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2023.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

Eldred. A. et al [12], proved many results on proximity pairs. One can also refer to many results, which is also recent, about proximity point of the pairs and their theorems in [1], [3], [13], [14], [16], [17], [23], [29], [31], [33], [34], [35], [36]. Additionally,  $B$ -contraction and Bianchini contraction definitions are located in [7] & [25], and using these, we define  $P - B$  contraction and  $P$ -Bianchini contraction.

This manuscript is laid out as follows: In Section 2, we recall the basics from the previous literature. In Section 3, we present the main results, which include the approximate best proximity pairs in contraction operators such as  $P - B$  contraction,  $P$ -Bianchini contraction and so on. Also, we discuss diameter of an approximate best proximity point for the pair  $(W, V)$  by using various operators based on the results of [26] and [28]. In Section 4, we provide some applications of our main results in the field of applied mathematics. Finally, in Section 5, we reach a conclusion.

## 2. Preliminaries

In this section, some definitions and lemmas from earlier research are recalled. These are then employed throughout the remainder of the main findings of this manuscript.

**DEFINITION 2.1.** [26], [28] Let  $W$  and  $V$  be two nonempty subsets of a metric space  $M$  and  $B : W \cup V \rightarrow W \cup V$  such that  $B(W) \subseteq V$  and  $B(V) \subseteq W$ . Then  $w$  is said to be an approximate best proximity point of the pair  $(W, V)$ , if  $d_b(w, Bw) \leq d_b(W, V) + \epsilon$ .

**REMARK 2.2.** [26], [28] Let  $P_{B\epsilon}(W, V) = \{w \in (W, V) : d_b(w, Bw) < d_b(W, V) + \epsilon, \text{ for some, } \epsilon > 0\}$  be denotes the set of all approximate best proximity pairs of pair  $(W, V)$  for a given  $\epsilon > 0$ . Also the pair  $(W, V)$  is said to be an approximate best proximity pair property if  $d_b(w, Bw) \leq d_b(W, V) \neq 0$ .

**EXAMPLE 2.3.** Let us take  $M = \mathbb{R}^2$  and  $W = \{(w, v) \in M : (w - v)^2 + v^2 \leq 1\}$  and  $V = \{(-w, v) \in M : (w + v)^2 + v^2 \leq 1\}$  with  $B(w, v) = (-w, v)$  for  $(w, v) \in M$ . Then  $d_b(w, v), B(w, v) \leq d_b(W, V) + \epsilon$ , for some  $\epsilon > 0$ . Hence  $P_{B\epsilon}(W, V) \neq \emptyset$ .

**THEOREM 2.4.** [26], [28] Let  $W$  and  $V$  be two nonempty subsets of a metric space  $M$ . Suppose that the mapping  $B : W \cup V \rightarrow W \cup V$  satisfying  $B(W) \subseteq V$  and  $B(V) \subseteq W$  and  $\lim_{n \rightarrow \infty} d_b(B^n w, B^{n+1} w) = d_b(W, V)$ , for some  $w \in (W \cup V)$ . Then the pair  $(W, V)$  is called an approximate best proximity pair.

**DEFINITION 2.5.** [26], [28] Let  $B : W \cup V \rightarrow W \cup V$  be a continuous map such that  $B(W) \subseteq V, B(V) \subseteq W$  and  $\epsilon > 0$ . We define the diameter  $Dtr(P_{B\epsilon}(W, V))$ , i.e.,  $Dtr(P_{B\epsilon}(W, V)) = \sup\{d_b(w, v) : w, v \in P_{B\epsilon}(W, V)\}$ .

**THEOREM 2.6.** [26], [28] Let  $W$  and  $V$  be two non-empty subsets of a metric space  $M$ . Suppose that a mapping  $B : W \cup V \rightarrow W \cup V$  satisfying  $B(W) \subseteq V, B(V) \subseteq W$  is a  $P - \alpha$  contraction and  $\epsilon > 0$ . Suppose that:

- (i)  $P_{B\epsilon}(W, V) \neq \emptyset$ ;
- (ii) for every  $\varphi > 0, \exists \psi(\varphi) > 0$  such that  $d_b(w, v) - d_b(Bw, Bv) \leq \varphi \Rightarrow d_b(w, v) \leq \psi(\varphi)$ , for every  $w, v \in P_{B\epsilon}(W, V) \neq \emptyset$ .

Then,  $Dtr(P_{B\epsilon}(W, V)) \leq \psi(2d_b(W, V) + \epsilon)$ .

DEFINITION 2.7. A mapping  $B : W \cup V \rightarrow W \cup V$  satisfying  $B(W) \subseteq V$  and  $B(V) \subseteq W$  is a  $P - B$  contraction operator if there exists  $b_1, b_2, b_3 \in (0, 1)$  with  $b_1 + b_2 + b_3 < 1$  such that

$$(1) \quad \begin{aligned} d_b(Bw, Bv) \leq & b_1[d_b(w, Bw) + d_b(v, Bv)] + b_2[d_b(w, v)] \\ & + b_3[d_b(w, Bv) + d_b(v, Bw)], \text{ for all } w, v \in W \cup V. \end{aligned}$$

DEFINITION 2.8. A mapping  $B : W \cup V \rightarrow W \cup V$  satisfying  $B(W) \subseteq V$  and  $B(V) \subseteq W$  is a  $P$ -Bianchini contraction operator if there exists  $b_1 \in (0, 1)$  such that

$$(2) \quad \begin{aligned} d_b(Bw, Bv) \leq & b_1 B_{ia}(w, v), \\ \text{where } B_{ia}(w, v) = & \max\{d_b(w, Bw), d_b(v, Bv)\}, \text{ for all } w, v \in W \cup V. \end{aligned}$$

DEFINITION 2.9. A mapping  $B : W \cup V \rightarrow W \cup V$  satisfying  $B(W) \subseteq V$  and  $B(V) \subseteq W$  is a  $P$ -Hardy and Rogers contraction operator if there exists  $b_1, b_2, b_3, b_4, b_5 \in (0, 1)$  with  $b_1 + b_2 + b_3 + b_4 + b_5 < 1$  such that

$$(3) \quad \begin{aligned} d_b(Bw, Bv) \leq & b_1 d_b(w, v) + b_2 d_b(w, Bw) + b_3 d_b(v, Bv) \\ & + b_4 d_b(w, Bv) + b_5 d_b(v, Bw), \text{ for all } w, v \in W \cup V. \end{aligned}$$

### 3. Main Results

This section is divided into two parts. The first one deals with qualitative results, and the other one deals with quantitative results; both are related to the approximate best proximity points for the pairs  $(V, W)$  on metric spaces.

**3.1. Qualitative theorems for P-contraction operators:** In this subsection, we prove some qualitative theorems about the approximate best proximity point for the pair  $(V, W)$  by using contraction operators such as the  $P - B$  contraction operator, the  $P$ -Bianchini contraction operator, and the  $P$ -Hardy Rogers contraction operator on a metric space.

THEOREM 3.1. *Let  $W$  and  $V$  be two non-empty subsets of a metric space  $M$ . Suppose that a mapping  $B : W \cup V \rightarrow W \cup V$  satisfying  $B(W) \subseteq V$  and  $B(V) \subseteq W$  is a  $P - B$  contraction operator then for every  $\epsilon > 0, P_{B\epsilon}(W, V) \neq \emptyset$ .*

*Proof.* Let  $\epsilon > 0$  and  $w \in W \cup V$ . Consider,

$$\begin{aligned} d_b(B^n w, B^{n+1} w) &= d_b(B(B^{n-1} w), B(B^n w)) \\ &\leq b_1[d_b(B^{n-1} w, B^n w) + d_b(B^n w, B^{n+1} w)] + b_2[d_b(B^{n-1} w, B^n w)] \\ &\quad + b_3[d_b(B^{n-1} w, B^{n+1} w) + d_b(B^n w, B^{n+1} w)] \text{ [By equation (1)]} \\ &\leq b_1 d_b(B^{n-1} w, B^n w) + b_1 d_b(B^n w, B^{n+1} w) + b_2 d_b(B^{n-1} w, B^n w) \\ &\quad + b_3 d_b(B^{n-1} w, B^n w) + b_3 d_b(B^n w, B^{n+1} w) \\ &= \left( \frac{b_1 + b_2 + b_3}{1 - b_2 - b_3} \right) d_b(B^{n-1} w, B^n w) \\ &= \lambda d_b(B^{n-1} w, B^n w), \text{ where } \lambda = \frac{b_1 + b_2 + b_3}{1 - b_2 - b_3}. \end{aligned}$$

But  $b_1, b_2$  and  $b_3 \in (0, 1)$  implies that  $\lambda \in (0, 1)$ . Therefore,

$$\lim_{n \rightarrow \infty} d_b(B^n w, B^{n+1} w) = 0, \text{ for all } w \in W \cup V.$$

Hence, by Theorem 2.4, it follows that

$$P_{B\epsilon}(W, V) \neq \emptyset, \text{ for all } \epsilon > 0.$$

□

**THEOREM 3.2.** *Let  $W$  and  $V$  be two non-empty subsets of a metric space  $M$ . Suppose that a mapping  $B : W \cup V \rightarrow W \cup V$  satisfying  $B(W) \subseteq V$  and  $B(V) \subseteq W$  is a  $P$ -Bianchini contraction operator then for every  $\epsilon > 0$ ,  $P_{B\epsilon}(W, V) \neq \emptyset$ .*

*Proof.* Let  $\epsilon > 0$  and  $w \in W \cup V$ . Consider,

CASE 1. Suppose  $B_{ia}(w, v) = d_b(w, Bw)$ . Then the Definition 2.8 becomes

$$d_b(Bw, Bv) \leq b_1 d_b(w, Bw)$$

Substitute  $v = Bw$  we get,

$$d_b(Bw, B^2 w) \leq b_1 d_b(w, Bw)$$

Again substituting  $w = Bw$  implies,

$$\begin{aligned} d_b(B^2 w, B^3 w) &\leq b_1 d_b(Bw, B^2 w) \\ &\leq (b_1)^2 d_b(w, Bw) \end{aligned}$$

Continuing this process we have,

$$d_b(B^n w, B^{n+1} w) \leq (b_1)^n d_b(w, Bw)$$

CASE 2. Suppose  $B_{ia}(w, v) = d_b(v, Bv)$ . Then the Definition 2.8 becomes

$$d_b(Bw, Bv) \leq b_1 d_b(v, Bv)$$

Substitute  $v = Bw$  we get,

$$d_b(Bw, B^2 w) \leq b_1 d_b(w, B^2 w)$$

This is impossible because  $b_1 \in (0, 1)$ . Therefore, CASE 2 does not exist. Now using CASE 1 and Theorem 2.4, we have

$$\lim_{n \rightarrow \infty} d_b(B^n w, B^{n+1} w) = 0, \text{ for all } w \in W \cup V.$$

And it follows that

$$P_{B\epsilon}(W, V) \neq \emptyset, \text{ for all } \epsilon > 0.$$

□

**COROLLARY 3.3.** *Let  $W$  and  $V$  be two non-empty subsets of a metric space  $M$ . Suppose that a mapping  $B : W \cup V \rightarrow W \cup V$  satisfying  $B(W) \subseteq V$  and  $B(V) \subseteq W$  and defined by  $d_b(Bw, Bv) \leq b_1 d_b(w, Bw)$  operator then for every  $\epsilon > 0$ ,  $P_{B\epsilon}(W, V) \neq \emptyset$ .*

*Proof.* It is a direct consequence of Theorem 3.2. □

**THEOREM 3.4.** *Let  $W$  and  $V$  be two non-empty subsets of a metric space  $M$ . Suppose that a mapping  $B : W \cup V \rightarrow W \cup V$  satisfying  $B(W) \subseteq V$  and  $B(V) \subseteq W$  is a  $P$ -Hardy Rogers operator then for every  $\epsilon > 0$ ,  $P_{B\epsilon}(W, V) \neq \emptyset$ .*

*Proof.* Let  $\epsilon > 0$  and  $w \in W \cup V$ . Consider,

$$\begin{aligned} d_b(B^n w, B^{n+1} w) &= d_b(B(B^{n-1} w), B(B^n w)) \\ &\leq b_1 d_b(B^{n-1} w, B^n w) + b_2 d_b(B^{n-1} w, B^n w) + b_3 d_b(B^n w, B^{n+1} w) \\ &\quad + b_4 d_b(B^{n-1} w, B^{n+1} w) + b_5 d_b(B^n w, B^n w) \text{ [ By equation (3) ]} \\ &\leq b_1 d_b(B^{n-1} w, B^n w) + b_2 d_b(B^{n-1} w, B^n w) + b_3 d_b(B^n w, B^{n+1} w) \\ &\quad + b_4 d_b(B^{n-1} w, B^n w) + b_4 d_b(B^n w, B^{n+1} w) \\ &= \left( \frac{b_1 + b_2 + b_4}{1 - b_3 - b_4} \right) d_b(B^{n-1} w, B^n w) \\ &= \lambda d_b(B^{n-1} w, B^n w), \text{ where } \lambda = \frac{b_1 + b_2 + b_4}{1 - b_3 - b_4}. \end{aligned}$$

But  $b_1, b_2, b_3, b_4$  and  $b_5 \in (0, 1)$  implies that  $\lambda \in (0, 1)$ . Therefore,

$$\lim_{n \rightarrow \infty} d_b(B^n w, B^{n+1} w) = 0, \text{ for all } w \in W \cup V.$$

Hence, by Theorem 2.4, it follows that

$$P_{B\epsilon}(W, V) \neq \emptyset, \text{ for all } \epsilon > 0.$$

□

- REMARK 3.5. 1. In Definition 2.7, substitute  $b_2 = \alpha$  and  $b_1 = b_3 = 0$ , then it becomes  $P - \alpha$  contraction operator and for every  $\epsilon > 0, P_{B\epsilon}(W, V) \neq \emptyset$ .  
 2. In Definition 2.7, substitute  $b_2 = b_3 = 0$ , then it becomes  $P$ -Kannan operator and for every  $\epsilon > 0, P_{B\epsilon}(W, V) \neq \emptyset$ .  
 3. In Definition 2.7, substitute  $b_1 = b_2 = 0$ , then it becomes  $P$ -Chatterjea operator and for every  $\epsilon > 0, P_{B\epsilon}(W, V) \neq \emptyset$ .  
 4. In Definition 2.8, substitute  $b_4 = b_5 = 0$ , then it becomes  $P$ -Reich operator and for every  $\epsilon > 0, P_{B\epsilon}(W, V) \neq \emptyset$ .  
 5. In Definition 2.8, substitute  $b_4 = b_5$ , then it becomes  $P$ -Ciric operator and for every  $\epsilon > 0, P_{B\epsilon}(W, V) \neq \emptyset$ .

**3.2. Quantitative results for P-contraction operators:** In this subsection, we prove some quantitative results of approximate best proximity point of the pairs  $(V, W)$  by using contraction operators such as the  $P - B$  contraction operator, the  $P$ -Bianchini contraction operator, and the  $P$ -Hardy Rogers contraction operator on a metric space.

THEOREM 3.6. *Let  $(M, d_b)$  be a metric space and  $B : W \cup V \rightarrow W \cup V$  satisfies the conditions of Theorem 3.1. Then,*

$$Dtr(P_{B\epsilon}(W, V)) \leq \frac{2(b_1 + b_3)d_b(W, V) + 2\epsilon(b_1 + b_3 + 1)}{1 - b_2 - 2b_3}, \text{ for all } \epsilon > 0.$$

*Proof.* Let  $\epsilon > 0$ . Also, condition (i) of Theorem 2.6 is proved by using Theorem 3.1. To show, condition (ii) of Theorem 2.6 holds. For that, take  $\varphi > 0$  and  $w, v \in P_{B\epsilon}(W, V)$ . Also  $d_b(w, v) - d_b(Bw, Bv) \leq \varphi$ . Then  $d_b(w, v) \leq d_b(Bw, Bv) + \varphi$ . since  $w, v \in P_{B\epsilon}(W, V)$  implies that  $d_b(w, Bw) \leq d_b(W, V) + \epsilon_1$  and  $d_b(v, Bv) \leq$

$d_b(W, V) + \epsilon_2$ . And choose  $\epsilon = \{\epsilon_1, \epsilon_2\}$ . Now,

$$\begin{aligned} d_b(w, v) &\leq d_b(Bw, Bv) + \varphi \\ &\leq b_1[d_b(W, V) + \epsilon + d_b(W, V) + \epsilon] + b_2[d_b(w, v)] \\ &\quad + b_3[d_b(w, v) + d_b(v, Bv) + d_b(v, w) + d_b(w, Bw)] + \varphi \\ &= b_1[2d_b(W, V) + 2\epsilon] + b_2d_b(w, v) + b_3[2d_b(w, v) + 2d_b(W, V) + 2\epsilon] + \varphi \\ &= \frac{2(b_1 + b_3)d_b(W, V) + 2\epsilon(b_1 + b_3) + \varphi}{1 - b_2 - 2b_3} \\ &= \psi(\varphi) \end{aligned}$$

Thus, for every  $\varphi > 0$ , there exists  $\psi(\varphi) > 0$  such that  $d_b(w, v) - d_b(Bw, Bv) \leq \varphi$  implies  $d_b(w, v) = \psi(\varphi)$ . Then by Theorem 2.6 ,

$$Dtr(P_{B\epsilon}(W, V)) \leq \psi(2\epsilon), \text{ for all } \epsilon > 0.$$

This means exactly

$$Dtr(P_{B\epsilon}(W, V)) \leq \frac{2(b_1 + b_3)d_b(W, V) + 2\epsilon(b_1 + b_3 + 1)}{1 - b_2 - 2b_3}, \text{ for all } \epsilon > 0.$$

□

**THEOREM 3.7.** *Let  $(M, d_b)$  be a metric space and  $B : W \cup V \rightarrow W \cup V$  satisfies the conditions of Theorem 3.2. Then,*

$$Dtr(P_{B\epsilon}(W, V)) \leq b_1d_b(W, V) + \epsilon(b_1 + 2), \text{ for all } \epsilon > 0.$$

*Proof.* Let  $\epsilon > 0$ . Also, condition (i) of Theorem 2.6 is proved by using Theorem 3.2. To show, condition (ii) of Theorem 2.6 holds. For that, take  $\varphi > 0$  and  $w, v \in P_{B\epsilon}(W, V)$ . Also  $d_b(w, v) - d_b(Bw, Bv) \leq \varphi$ . Then  $d_b(w, v) \leq d_b(Bw, Bv) + \varphi$ . since  $w, v \in P_{B\epsilon}(W, V)$  implies that  $d_b(w, Bw) \leq d_b(W, V) + \epsilon_1$  and  $d_b(v, Bv) \leq d_b(W, V) + \epsilon_2$ . And choose  $\epsilon = \{\epsilon_1, \epsilon_2\}$ . Now,

$$\begin{aligned} d_b(w, v) &\leq d_b(Bw, Bv) + \varphi \\ &\leq b_1d_b(w, Bw) + \varphi \\ &\leq b_1(d_b(W, V) + \epsilon) + \varphi \\ &= \psi(\varphi) \end{aligned}$$

Thus, for every  $\varphi > 0$ , there exists  $\psi(\varphi) > 0$  such that  $d_b(w, v) - d_b(Bw, Bv) \leq \varphi$  implies  $d_b(w, v) = \psi(\varphi)$ . Then the Theorem 2.6 gives,

$$Dtr(P_{B\epsilon}(W, V)) \leq \psi(2\epsilon), \text{ for all } \epsilon > 0.$$

This means exactly

$$Dtr(P_{B\epsilon}(W, V)) \leq b_1d_b(W, V) + \epsilon(b_1 + 2), \text{ for all } \epsilon > 0.$$

□

**THEOREM 3.8.** *Let  $(M, d_b)$  be a metric space and  $B : W \cup V \rightarrow W \cup V$  satisfies the conditions of Theorem 3.4. Then,*

$$Dtr(P_{B\epsilon}(W, V)) \leq \frac{(1 - b_1)d_b(W, V) + \epsilon(3 - b_1)}{b_2 + b_3}, \text{ for all } \epsilon > 0.$$

*Proof.*  $\epsilon > 0$ . Also, condition (i) of Theorem 2.6 is proved by using Theorem 3.4. To show, condition (ii) of Theorem 2.6 holds. For that, take  $\varphi > 0$  and  $w, v \in P_{B\epsilon}(W, V)$ . Also  $d_b(w, v) - d_b(Bw, Bv) \leq \varphi$ . Then  $d_b(w, v) \leq d_b(Bw, Bv) + \varphi$ . since  $w, v \in P_{B\epsilon}(W, V)$  implies that  $d_b(w, Bw) \leq d_b(W, V) + \epsilon_1$  and  $d_b(v, Bv) \leq d_b(W, V) + \epsilon_2$ . And choose  $\epsilon = \{\epsilon_1, \epsilon_2\}$ . Now,

$$\begin{aligned} d_b(w, v) &\leq d_b(Bw, Bv) + \varphi \\ &\leq b_1 d_b(w, v) + b_2 [d_b(W, V) + \epsilon] + b_3 [d_b(W, V) + \epsilon] + b_4 d_b(w, v) \\ &\quad + b_4 [d_b(W, V) + \epsilon] + b_5 d_b(w, v) + b_5 [d_b(W, V) + \epsilon] + \varphi \\ &= (b_1 + b_4 + b_5) d_b(w, v) + (b_2 + b_3 + b_4 + b_5) [d_b(W, V) + \epsilon] + \varphi \\ &= \frac{(b_2 + b_3 + b_4 + b_5) [d_b(W, V) + \epsilon] + \varphi}{1 - (b_1 + b_4 + b_5)} \\ &= \psi(\varphi) \end{aligned}$$

Thus, for every  $\varphi > 0$ , there exists  $\psi(\varphi) > 0$  such that  $d_b(w, v) - d_b(Bw, Bv) \leq \varphi$  implies  $d_b(w, v) = \psi(\varphi)$ . Then, the Theorem 2.6 gives

$$Dtr(P_{B\epsilon}(W, V)) \leq \psi(2\epsilon), \text{ for all } \epsilon > 0.$$

That is,

$$Dtr(P_{B\epsilon}(W, V)) \leq \frac{(b_2 + b_3 + b_4 + b_5) d_b(W, V) + \epsilon(b_2 + b_3 + b_4 + b_5 + 2)}{1 - (b_1 + b_4 + b_5)}, \text{ for all } \epsilon > 0.$$

This means exactly

$$Dtr(P_{B\epsilon}(W, V)) \leq \frac{(1 - b_1) d_b(W, V) + \epsilon(3 - b_1)}{b_2 + b_3}, \text{ for all } \epsilon > 0.$$

□

**COROLLARY 3.9.** *Let  $(M, d_b)$  be a metric space and  $B : W \cup V \rightarrow W \cup V$  satisfies the conditions of Corollary 3.3. Then,*

$$Dtr(P_{B\epsilon}(W, V)) \leq b_1 d_b(W, V) + \epsilon(b_1 + 2), \text{ for all } \epsilon > 0.$$

*Proof.* It is a direct consequence of Theorem 3.7. □

**REMARK 3.10.** 1. In  $P - \alpha$  contraction operator,

$$Dtr(P_{B\epsilon}(W, V)) \leq \frac{2(\epsilon + d_b(W, V))}{b_1}, \text{ for all } \epsilon > 0.$$

2. In  $P$ -Kannan operator,  $Dtr(P_{B\epsilon}(W, V)) \leq 2\epsilon(1 + b_1) + 2b_1 d_b(W, V)$ , for all  $\epsilon > 0$ .

3. In  $P$ -Chatterjea operator,  $Dtr(P_{B\epsilon}(W, V)) \leq \frac{2[\epsilon(1 + b_1) + b_1 d_b(W, V)]}{1 - 2b_1}$ , for all  $\epsilon > 0$ .

4. In  $P$ -Reich operator,  $Dtr(P_{B\epsilon}(W, V)) \leq \frac{(1 - b_1)(d_b(W, V)) + (3 - b_1)\epsilon}{1 - b_1}$ , for all  $\epsilon > 0$ .

5. In  $P$ -Ciric operator,  $Dtr(P_{B\epsilon}(W, V)) \leq \frac{(1 - b_1) d_b(W, V) + (3 - b_1)\epsilon}{b_2 + b_3}$ , for all  $\epsilon > 0$ .

#### 4. Applications

Approximate best proximity point theory covers a wide range of applications in mathematics, particularly differential geometry, numerical analysis, and so on. By reading [22] and the references there in, one can find a variety of applications involving approximate best proximity point results in the field of mathematics. The two examples below demonstrate how to apply approximate best proximity point findings in differential equations.

EXAMPLE 4.1. Consider  $z''(w) = 6z^2(w)$ ,  $0 \leq w \leq 1$  subject to  $z(0) = \frac{1}{4}$ ,  $z(1) = \frac{1}{9}$ . Exact solution is  $z_0(w) = \frac{-5w}{36} + \frac{1}{4}$ . Consider a mapping  $T : [0, 1] \rightarrow [0, 1]$  defined by

$$\begin{aligned} T(z) &= z + \int_0^1 G(w, v)[z''(v) - \phi(v, z(v), z'(v))]dv \\ &= \frac{-5w}{36} + \frac{1}{4} - \int_0^1 G(w, v)\phi(v, z(v), z'(v))dv \\ &= \frac{-5w}{36} + \frac{1}{4} - \int_0^1 G(w, v)6z''(v)dv \end{aligned}$$

Consider,

$$\begin{aligned} |T(z_1) - T(z_2)| &= 6 \left| - \int_0^1 G(w, v)z_1^2(v)dv + \int_0^1 G(w, v)z_2^2(v)dv \right| \\ &= 6 \left( \int_0^1 |G(w, v)|^2 dv \right)^{\frac{1}{2}} \left( \int_0^1 |z_2^2(v) - z_1^2(v)|^2 dv \right)^{\frac{1}{2}} \\ &\leq \frac{1}{4\sqrt{3}} \left( \int_0^1 |z_2^2(v) - z_1^2(v)|^2 dv \right)^{\frac{1}{2}} \\ &< \sup_{[0,1]} |z_1(v) - z_2(v)| \end{aligned}$$

Hence,  $T$  is a contraction and it has approximate best proximity point.

EXAMPLE 4.2. Consider  $z''(w) = \frac{3v^2(w)}{2}$ ,  $0 \leq v \leq 1$  subject to  $z(0) = 4$ ,  $z(1) = 1$ . Exact solution is  $z(w) = \frac{4}{(1+w)^2}$ . Consider a mapping  $W : [0, 1] \rightarrow [0, 1]$  by

$$(4) \quad W(z) = w_2 + \int_0^1 G(v, w)[z''(w) - \phi(w, z(w))]dw$$

Consider,  $z''(v) = 0$  which implies

$$(5) \quad z(v) = c_1v + c_2$$

By initial condition we have  $c_2 = 4$  and  $c_1 = -3$ . Then (5) becomes  $z(v) = -3w_1 + 4$ .

$$\begin{aligned} W(z) &= -3v + 4 + \int_0^1 G(v, w)[z''(w) - \phi(w, z(w))]dw \\ &= -3v + 4 + \int_0^1 G(v, w)z''(w)dw - \int_0^1 G(v, w)\phi(w, z(w))dw \\ &= -3v + 4 + \int_0^1 G(v, w)\frac{3}{2}z^2(w)dw \end{aligned}$$



Consider,

$$\begin{aligned}
 |W(z_1) - W(z_2)| &= \left| - \int_0^1 G(v, w) \frac{3}{2} z_2^2(w) dw + \int_0^1 G(v, w) \frac{3}{2} z_1^2(w) dw \right| \\
 &= \frac{3}{2} \left| \int_0^1 G(v, w) [z_2^2(w) - z_1^2(w)] dw \right| \\
 &\leq \frac{3}{2} \left( \int_0^1 |G(v, w)|^2 dw \right)^{\frac{1}{2}} \left[ \int_0^1 |z_2^2(w) - z_1^2(w)|^2 dw \right]^{\frac{1}{2}} \\
 &\leq \frac{3}{2} \left( \int_0^w w^2(1-v)^2 dv + \int_v^1 v^2(1-w)^2 dw \right)^{\frac{1}{2}} \left[ \int_0^1 |z_2^2(w) - z_1^2(w)|^2 dw \right]^{\frac{1}{2}} \\
 &\leq \frac{3}{2} \left\{ \frac{(1-v)^2 v^3}{3} + \frac{v^2(1-v)^3}{3} \right\}^{\frac{1}{2}} \left[ \int_0^1 |z_2^2(w) - z_1^2(w)|^2 dw \right]^{\frac{1}{2}} \\
 &\leq \frac{3}{2} \left\{ \frac{(1-v)^2}{3} [v^3 + v^2(1-v)] \right\}^{\frac{1}{2}} \left[ \int_0^1 |z_2^2(w) - z_1^2(w)|^2 dw \right]^{\frac{1}{2}} \\
 &\leq \frac{3}{2} \left\{ \frac{(1-v)^2 v^2}{3} \right\}^{\frac{1}{2}} \left[ \int_0^1 |z_2^2(w) - z_1^2(w)|^2 dw \right]^{\frac{1}{2}} \\
 &\leq \frac{3}{8\sqrt{3}} \left[ \int_0^1 |z_2^2(w) - z_1^2(w)|^2 dw \right]^{\frac{1}{2}} \\
 &\leq \frac{\sqrt{3}}{8} \left[ \int_0^1 |z_2^2(w) - z_1^2(w)|^2 dw \right]^{\frac{1}{2}} \\
 &\leq \frac{\sqrt{3}}{8} \sup_{[0,1]} |z_2(w) - z_1(w)| \\
 &\leq \sup_{[0,1]} |z_2(w) - z_1(w)|
 \end{aligned}$$

Hence,  $W$  is contraction, it has approximate best proximity point.

EXAMPLE 4.3. Let us consider a numerical problem  $\int_0^\pi \sin p dx$ . To solve this, by using Simpson's rule, we get  $p = 2.0008$ . Similarly, to solve this, by using Trapezoidal rule, we get  $p = 1.955$ . But the actual solution is  $p = 2$ . Therefore, in both methods, we get only an approximate solution.

### 5. Conclusion

This work provides a series of contraction mappings to demonstrate several approximate best proximity point theorems on metric spaces. It is essential to note that all of the conclusions made in the current paper generate better constrained approximations of best proximity points, mostly in minimising condition  $\epsilon \rightarrow 0$ . In order to confirm the presence of an approximate fixed points, alternative discoveries presented in the later can be demonstrated in a lower environment. Thus, the concept of an approximate best proximity point of the pair  $(W, V)$  is just as significant as the concept of best proximity point of the pair  $(W, V)$ .

### Author Contribution Statements

For making this article, all the author's contributed equally.

### Competing Interests

All the author's said that they have no competing interests.

### Acknowledgements

Also, all the authors thank the anonymous referee(s) and editorial members of the paper for their valuable recommendations. Once again, all the authors express their gratitude to the chief editor for giving us the opportunity to reset the manuscript in a nice way.

### References

- [1] Balraj, D., Marudai, M., Mitrovic, Z. D., Ege, O., Piramanantham, V., *Existence of best proximity points satisfying two constraint inequalities*, Electronic Research Archive **28** (1) (2020), 549–557.
- [2] Banach, S., *Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales*, Fund.Math. **3** (1922), 133–181.
- [3] Beer, G., Pai, D., *Proximal maps, prox maps and coincidence points*, Numerical Functional Analysis and Optimization **11** (5-6) (1990), 429–448.
- [4] Berinde, M., *Approximate fixed point theorems*, Stud. Univ. "Babes-Bolyai", Mathematica LI **1** (2006), 11–25.
- [5] Berinde, V., *Iterative approximation of fixed points*, Editura Efemeride, Baia Mare, 2002.
- [6] Berinde, V., *On the approximation of fixed points of weak contractive mappings*, Carpathian J. Math. **19** (1) (2003), 7–22.
- [7] Bianchini, R. M. T., *Su un problema di S. Reich riguardante la teoria dei punti fissi*, Bolletino U.M.I. **4** (5) (1972), 103–106.
- [8] Chatterjea, S.K., *Fixed point theorems*, C.R. Acad.Bulg. Sci. **25** (1972), 727–730.
- [9] Ćirić, L. B., *Generalized contractions and fixed point theorems*, Publ.Inst.Math.(Bulgr). **12** (26) (1971), 19–26.
- [10] Dass, B. K., Gupta, S., *An extension of Banach contraction principle through rational expression*, Communicated by F.C. Auluck, FNA., (1975).
- [11] Dey, D., Saha, M., *Approximate fixed point of Reich operator*, Acta Mathematica Universitatis Comenianae, (2012).
- [12] Eldred, A. A., Veeramani, P., *Existence and convergence of best proximity points*, Journal of Mathematical Analysis and Applications, **323** (2) (2006), 1001–1006.
- [13] Fallahi, K., Rad, G. S., Fulga, A., *Best proximity points for  $(\phi\psi)$ -weak contractions and some applications*, Filomat, **37** (6), (2023), 18351842.
- [14] Gardasevic-Filipovic, M., Kukic, K., Gardasevic, D., Mitrovic, D., *Some best proximity point results in the orthogonal O-complete b-metric-like spaces*, Journal of Contemporary Mathematical Analysis (Armenian Academy of Sciences) **58** (2023), 105115.
- [15] Hardy, G. E., Rogers, T. D., *A generalization of fixed point theorem of Reich*, Can. Math. Bull. **16** (1973), 201–206.
- [16] Iqbal, M., Batool, A., Ege, O., de la Sen, M., *Fixed point of generalized weak contraction in b-metric spaces*, Journal of Function Spaces, (2021), 8 pages, Article ID 2042162.
- [17] Iqbal, M., Batool, A., Ege, O., de la Sen, M., *Fixed point of almost contraction in b-metric spaces*, Journal of Mathematics (2020), 3218134, 1–6.
- [18] Jaggi, D. S., *Some unique fixed point theorems*, Indian Journal of Pure and Applied Mathematics, **8**, (1977), 223–230.
- [19] Kannan, R., *Some results on fixed points*, Bull Calcutta Math. Soc **60** (1968), 71–76.
- [20] Kannan, R., *Some results on fixed points, II*. Am. Math.Mon. **76** (1969), 405–408.

- [21] Khan, M. S., *A fixed point theorems for metric spaces*, Rendiconti Dell 'istituto di mathematica dell' Universtia di tresti, **8**, (1976), 69–72.
- [22] Khuri, S. A., Louhichi, I., *A novel Ishikawa-Green's fixed point scheme for the solution of BVPs*, Appl. Math. Lett. **82** (2018), 50–57.
- [23] Kim, W. K., Lee, K. H., *Corrigendum to existence of best proximity pairs and equilibrium pairs*, J. Math. Anal. Appl. **316** (2006), 433–446.
- [24] Kirk, W. A., Reich, S., Veeramani, P., *Proximinal retracts and best proximity pair theorems*, Numerical Functional Analysis and Optimization **24** (7-8), (2003).
- [25] Marudai, M., Bright V. S., *Unique fixed point theorem weakly B-contractive mappings*, Far East journal of Mathematical Sciences(FJMS) **98** (7) (2015), 897–914.
- [26] Mohsenalhosseini, S.A.M., *Approximate best proximate pairs in metric space for contraction maps*, Fixed Point Theory **4** (2) (2014), 310–324.
- [27] Mohsenalhosseini, S. A. M., Saheli. M., *Some of family of contractive type maps and approximate common fixed point*, J.Fixed Point Theory (2021), 2021:2.
- [28] Mohsenalhosseini, S.A.M., Mzaheri, H.,Dehghan, M. A., *Approximate best proximity pairs in metric space*, Abstr. Appl. Anal., (2011), Article ID 596971.
- [29] Nallaselli, G., Gnanaprakasam, A. J., Mani, G., Ege, O., *Solving integral equations via admissible contraction mappings*, Filomat, **36** (14), (2022), 4947–4961.
- [30] Reich, S., *Some remarks connecting contraction mappings*, Can. Math. Bull., **14**, (1971), 121–124.
- [31] Singer, I., *Best approximation in normed linear spaces by elements of linear subspaces*, Translated from the Romanian by Radu Georgescu. Die Grundlehren der mathematischen Wissenschaften, Band 171 Publishing House of the Academy of the Socialist Republic of Romania, Bucharest; Springer-Verlag, New York-Berlin, 1970.
- [32] Tis, S., Torre, A., Branzei, R., *Approximate fixed point theorems*, Libertas Mathematica **23** (2003), 35–39.
- [33] Vetrivel, V., Veeramani, P., Bhattacharyya, P., *Some extensions of Fan's best approximation theorem*, Numerical Functional Analysis and Optimization **13** (3-4), (1992), 397–402.
- [34] Włodarczyk, K., Plebaniak, R., Banach, A., *Best proximity points for cyclic and noncyclic set-valued relatively quasi-asymptotic contractions in uniform spacand es*, Nonlinear Analysis: Theory, Methods and Applications, 2009, 3332–3341.
- [35] Włodarczyk, K., Plebaniak, R., Obczyński, C., *Convergence theorems, best approximation and best proximity for set-valued dynamic systems of relatively quasi-asymptotic contractions in cone uniform spaces*, Nonlinear Analysis: Theory, Methods and Applications **72** (2) (2010), 794–805.
- [36] Xu, X. B., *A result on best proximity pair of two sets*, Journal of Approximation Theory **54** (3) (1988), 322–325.
- [37] Zamfirescu, T., *Fixed Point theorems in metric spaces*, Arch. Math.(Basel) **23** (1972), 292–298.

### R. Theivaraman

Department of Mathematics, Bharathidasan University, Tiruchirappalli-620 024,  
Tamilnadu, India

*E-mail:* deivaraman@gmail.com

### P. S. Srinivasan

Department of Mathematics, Bharathidasan University, Tiruchirappalli-620 024,  
Tamilnadu, India

*E-mail:* pssrini@bdu.ac.in

### A. Herminau Jothy

Department of Mathematics, Bharathidasan University, Tiruchirappalli-620 024,  
Tamilnadu, India

*E-mail:* jothyrose@gmail.com