

New record of two spirotrichous ciliate species (Protozoa, Ciliophora) collected from coasts of South Korea

Ji Hye Choi and Jae-Ho Jung*

Department of Biology, Gangneung-Wonju National University, Gangneung 25457, Republic of Korea

*Correspondent: jhjung@gwnu.ac.kr

Two newly recorded ciliates, *Aspidisca major* (Madsen, 1931) Kahl, 1932 and *Metaurostylopsis rubra* Song and Wilbert, 2002, were collected from eastern and southern coasts of South Korea, respectively. The morphology of these two species were studied based on observations of protargol-stained specimens. The morphological characteristics of the two species are as follows: (1) *Aspidisca major* is characterized by a size of 76–78 × 42–49 µm after protargol impregnation, two distinct macronuclei, seven frontoventral cirri in “polystyla-arrangement”, and dorsal kinety 1 extends to posterior end of dorsal kinety 2; and (2) *Metaurostylopsis rubra* is characterized by a size of 68–77 × 49–58 µm size after protargol impregnation, brick-reddish body color, and large number of marginal rows. In the present study, we provide a brief diagnosis, remarks, and photomicrographs.

Keywords: *Aspidisca major*, marine ciliates, *Metaurostylopsis rubra*, taxonomy

© 2024 National Institute of Biological Resources

DOI:10.12651/JSR.2024.13.4.373

INTRODUCTION

The genera *Aspidisca* Ehrenberg, 1830 and *Metaurostylopsis* Song *et al.*, 2001 are widely distributed globally in marine ecosystems. Up to date, about 60 nominal *Aspidisca* species and eight *Metaurostylopsis* species have been recorded (Wu and Curds, 1979; Fernandez-Leborans de Zaldumbide, 1987; Valbonesi, 1996; Berger, 2001; Song and Wilbert, 2002; Lei *et al.*, 2005; Shao *et al.*, 2008a; 2008b; Jankowski, 2009; Shen *et al.*, 2010; Chen *et al.*, 2011; Jung *et al.*, 2011; Jiang *et al.*, 2013; Lu *et al.*, 2016; Song *et al.*, 2020; Zheng *et al.*, 2022; Choi *et al.*, 2024). According to the National Institution of Biological Resources (2023), only 11 *Aspidisca* species (*A. aculeata* (Ehrenberg, 1838) Kahl, 1932; *A. cicada* (Müller, 1786) Claparède and Lachmann, 1858; *A. dentata* Kahl, 1928; *A. hexeris* Quennerstedt, 1869; *A. leptaspis* Fresenius, 1865; *A. lynceus* (Müller, 1773) Ehrenberg, 1830; *A. orthopogon* Deroux and Tuffrau, 1965; *A. polypoda* (Dujardin, 1841) Kahl, 1932; *A. polystyla* Stein, 1859; *A. steini* (Buddenbrock, 1920) Kahl, 1932; and *A. turrita* (Ehrenberg, 1838) Claparède and Lachmann, 1858) and only 4 *Metaurostylopsis* species (*M. cheni* Chen *et al.*, 2011; *M. marina* (Kahl, 1932) Song *et al.*, 2001; *M. salina* Lei *et al.*, 2005; and *M. struederkypkeae* Shao *et al.*, 2008) have been recorded

in South Korea, indicating undersampling of the Korean ciliate fauna (Shin and Kim, 1988; Lei *et al.*, 2005; Li *et al.*, 2010; Park *et al.*, 2012; Kim *et al.*, 2016; Kim and Jung, 2018; Kim *et al.*, 2018; 2020; Choi *et al.*, 2020; Choi *et al.*, 2023). Therefore, this study aims to further explore the diversity and distribution of these genera in Korean marine environments.

During a field survey, we identified two species, 1) *Aspidisca major* (Madsen, 1931) Kahl, 1932; and 2) *Metaurostylopsis rubra* Song and Wilbert, 2002, previously unrecorded in Korea. In this paper, we provide brief diagnosis, remarks, photomicrographs, and information about collection sites and conditions.

MATERIALS AND METHODS

The detailed information about collection sites and conditions are provided in ‘Material examined’ section for each species. After gently stirring up surface marine water, samples were collected with bottom sediments. All samples were transferred to the laboratory and kept in Plant culture dishes at room temperature (ca. 20°C) with 1–3 autoclaved rice grains. The morphology of each species was studied using a stereomicroscope (SZ11; Olympus, Tokyo, Japan), and photomicrographs were captured

using a digital camera (DP74; Olympus). The protargol powder was synthesized using the method of Pan *et al.* (2013) and Kim and Jung (2017). The protargol-impregnated specimens were prepared using the ‘procedure A’ method of Foissner (2014). The differential through-focal images of the protargol-impregnated specimens were merged using Helicon Focus ver. 8.1.0 software (Helicon-Soft Ltd, Kharkiv, Ukraine). The basic terminology and taxonomic classification mainly followed Lynn (2008), Song *et al.* (2001), and Wu and Curds (1979).

RESULTS AND DISCUSSION

Phylum Ciliophora Doflein, 1901

Class Spirotrichea Bütschli, 1889

Order Euplotida Small and Lynn, 1985

Suborder Euplotina Jankowski, 1979

Family Aspidiscidae Ehrenberg, 1830

Genus *Aspidisca* Ehrenberg, 1830

1. *Aspidisca major* (Madsen, 1931) Kahl, 1932 (Fig. 1)

Material examined. Marine water (salinity 37.8‰, temperature 18.3°C) collected from Sodol beach, Jumun-ri, Jumunjin-eup, Gangneung-si, Gangwon-do, Korea (37° 54'24.6"N, 128°49'41.7"E) on May 15, 2023.

Diagnosis. Size 76–78×42–49 µm after protargol im-

regnation ($n=3$); body bean-shaped; cortex rigid; peristomial spur and projections along left margin on ventral side lacking (Fig. 1A); anterior adoral zone of membranelles (AZM1) about 4 µm long with 4 membranelles, posterior adoral zone of membranelles (AZM2) about 24 µm long after protargol impregnation and with 15–18 membranelles; invariably seven frontoventral cirri in “polystyla-arrangement”; five transverse cirri, cilia of each transverse cirrus not separated (Fig. 1A); four dorsal kineties with 25–28, 18–22, 19–23, and 22–24 dikinetids in dorsal kineties 1–4, respectively (Fig. 1B); dorsal kinety 1 distinctly curved to right posteriorly and ends near posterior end of dorsal kinety 2 (Fig. 1C); two ellipsoid macronuclei, micronucleus not observed (Fig. 1D).

Distribution. Germany, South Korea

Remarks. The Korean population of *Aspidisca major* corresponds with the German population redescribed by Kahl (1932) in all available aspects. *Aspidisca major* is a special species among the congeners in having two macronuclei (Fig. 1D). Because of this feature, *A. major*, *A. binucleata* Kahl, 1932, *A. fusca* Kahl, 1928, and *A. mutans* Kahl, 1932 can be easily distinguished from other members of the genus *Aspidisca*. However, *A. major* differs from *A. binucleata* in the number of frontoventral cirri (7 vs. 9). *Aspidisca mutans* differs from *A. major* by the number of frontoventral cirri (12 vs. 7) and the number of transverse cirri (7 vs. 5) (Kahl, 1932). *Aspidisca fusca* differs from *A. major* by the presence of a peristomial spur




Fig. 1. *Aspidisca major* after protargol impregnation (A–D). A, B. Ventral (A) and dorsal (B) view, showing the body shape, anterior and posterior portion of adoral zone of membranelles, the transverse cirri, the frontoventral cirri in “polystyla-arrangement”, and dorsal kineties 1–4. C. Dorsal view showing the arrangement of dorsal kineties. Dorsal kinety 1 curved posteriorly and extends to posterior end of dorsal kinety 2. D. Two ellipsoidal macronuclei (arrowheads). AZM1 and 2, anterior and posterior adoral zone of membranelles; DK1–4, dorsal kineties 1–4; FVC, frontoventral cirri; TC, transverse cirri. Scale bars = 20 µm (A, B); 10 µm (C, D).




Fig. 2. *Metaurostylopsis rubra* after protargol impregnation (A, B). A. Ventral view showing the body shape, adoral zone of membranelles, frontal and transverse cirri, midventral pairs and row, and left and right marginal rows. Arrowhead denotes one buccal cirrus. B. Dorsal view showing the dorsal kinetics 1–3 and macronuclear nodules. Double arrowhead denotes fragment-like kinetics anterior to the leftmost right marginal row. AZM, adoral zone of membranelles; DK1–3, dorsal kinetics 1–3; FC, frontal cirri; LMR, left marginal rows; MA, macronuclear nodules; MP, midventral pairs; MR, midventral row; RMR, right marginal rows; TC, transverse cirri. Scale bar = 30 µm.

(present vs. absent) (Wu and Curds, 1979). Considering the arrangement of frontoventral cirri, number of membranelles in AZM1 and the presence of peristomial spur, the Korean population of *A. major* is similar to *A. steini* however, they differ mainly in the number of macronuclei (two vs. one) and larger body size ($76\text{--}78 \times 42\text{--}49$ µm vs. $20\text{--}35 \times 15\text{--}27$ µm) (Wu and Curds, 1979; Song and Wilbert, 1997).

Voucher slides. Two slides with protargol-impregnated specimens were deposited at the National Marine Biodiversity Institute of Korea (MABIK PR00045067, MABIK PR00045068).

Order Urostylida Jankowski, 1979

Family Urostylidae Bütschli, 1889

Genus *Metaurostylopsis* Song, Petz and Warren, 2001

2. *Metaurostylopsis rubra* Song and Wilbert, 2002 (Fig. 2)

Material examined. Marine water (salinity 36.1‰, temperature 16.8°C) collected from Sanyang-eup, Tong-

yeong-si, Gyeongsangnam-do, South Korea ($34^{\circ}49'16''$ N, $128^{\circ}26'27''$ E) on April 17, 2023.

Diagnosis. Size $68\text{--}77 \times 49\text{--}58$ µm after protargol impregnation ($n=3$); body oval to ellipsoidal with brick-redish color; body flexible and slightly contractile; adoral zone of membranelles occupies about 50% of body length; 36–44 adoral membranelles; 1 buccal cirrus; 3 frontal, 8–9 frontoterminal, and 5–6 transverse cirri; 13–16 midventral pairs and a midventral row with about 8–12 cirri, extending to the midline of body; 10–12 left and 6–7 right marginal rows (Fig. 2A); three dorsal kinetics (Fig. 2B); and about 70 macronuclear nodules.

Distribution. Antarctic, South Korea

Remarks. The Korean population of *Metaurostylopsis rubra* is similar to the type population of *M. rubra* described by Song and Wilbert (2002) in most features except for the slightly non-overlapping number of midventral pairs (13–16 vs. 8–11) and left marginal rows (10–12 vs. 6–9) (Song and Wilbert, 2002). *Metaurostylopsis rubra* can be easily distinguished from other congeners by the cell color and the number of midventral pairs and left marginal rows (Fig. 2A). Since the number of living

cells were not enough to examine in detail, we cannot determine whether the reddish color of the cells is due to the reddish cytoplasm or the cortical granules. Among the congeners, there is no species with more than six right marginal rows (Song *et al.*, 2020). Also, only two species, *M. struederkypkeae* and *M. parastruederkypkeae* Lu *et al.*, 2016, have reddish body color like *M. rubra*, but they differ from the present Korean population by the number of right marginal rows (3 vs. 3–5 vs. 6–7) (Shao *et al.*, 2008b; Lu *et al.*, 2016; Song *et al.*, 2020).

Voucher slides. Two slides with protargol-impregnated specimens were deposited at the National Marine Biodiversity Institute of Korea (MABIK PR00045065, MABIK PR00045066).

CONFLICTS OF INTEREST

The author of this paper has no affiliation with any interests and is solely responsible for the paper.

ACKNOWLEDGEMENTS

This work was supported by the management of Marine Fishery Bio-resources Center (2024) funded by the National Marine Biodiversity Institute of Korea (MABIK), the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBRE202406), and 2024 Academic Research Support Program in Gangneung-Wonju National University.

REFERENCES

- Berger, H. 2001. Catalogue of ciliate names 1. Hypotrichs. Verlag Helmet Berger, Salzburg.
- Chen, X., J. Huang and W. Song. 2011. Ontogeny and phylogeny of *Metaurostylopsis cheni* sp. n. (Protozoa, Ciliophora), with estimating the systematic position of *Metaurostylopsis*. *Zoologica Scripta* 40(1):99–111.
- Choi, J.H., A. Omar and J.-H. Jung. 2023. New record of three *Aspidisca* species (Protozoa, Ciliophora) from South Korea. *Journal of Species Research* 12(1):90–94.
- Choi, J.H., A. Omar and J.-H. Jung. 2024. Morphology, morphogenesis, and molecular phylogeny of *Aspidisca koreana* n. sp. (Ciliophora, Euplotida) from South Korea. *European Journal of Protistology* 92:126037.
- Choi, J.M., J.-H. Jung, J.-H. Kang and Y.-O. Kim. 2020. New record of four ciliates (Protozoa, Ciliophora) collected from rocky intertidal pools of South Korea. *Journal of Species Research* 9:455–461.
- Fernandez-Leborans, G. and M.C. de Zaldumbide. 1987. A new species of the genus *Aspidisca* (Ciliophora, Hypotrichida). *Journal of National History* 21:1293–1301.
- Foissner, W. 2014. An update of ‘basic light and scanning electron microscopic methods for taxonomic studies of ciliated protozoa’. *International Journal of Systematic and Evolutionary Microbiology* 64:271–292.
- Jankowski, A.W. 2009. *Aspidisca beringiana* sp. nov. and *Simbioidisca* subgen. nov. (Ciliophora, Euplotida), a symbiont of terebellid polychaetes in the Bering Sea. *Zoosystematica Rossica* 18:184–190.
- Jiang, J., J. Huang, L. Li, C. Shao, K.A.S. Al-Rasheid, S.A. Al-Farraj and Z. Chen. 2013. Morphology of two marine euplotids (Ciliophora: Euplotida), *Aspidisca fusca* Kahl, 1928 and *A. hexeris* Quennerstedt, 1869, with notes on their small subunit rRNA gene sequences. *European Journal of Protistology* 49:634–643.
- Jung, J.-H., Y.-S. Park, S.H. Kim, H.-G. Choi and G.-S. Min. 2011. A new marine ciliate, *Metaurostylopsis antarctica* nov. spec. (Ciliophora, Urostylida) from the Antarctic ocean. *Acta Protozoologica* 50:289–300.
- Kahl, A. 1932. Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 3. Spirotricha. *Tierwelt Deutschlands* 25:299–650.
- Kim, J.H. and J.-H. Jung. 2017. Cytological staining of protozoa: A case study on the impregnation of hypotrichs (Ciliophora: Spirotrichea) using laboratory-synthesized protargol. *Animal Cells and Systems* 21:412–418.
- Kim, J.H. and J.-H. Jung. 2018. New record of five ciliates (Protozoa, Ciliophora) collected in eastern areas of Gangwon-do province, Korea. *Journal of Species Research* 7:181–186.
- Kim, J.H., J.H. Moon and J.-H. Jung. 2018. New records of 24 ciliate species (Protozoa, Ciliophora) collected in South Korea. *Journal of Species Research* 7:291–314.
- Kim, J.H., A. Omar, J.H. Moon and J.-H. Jung. 2020. Taxonomy of 16 indigenous ciliate species (Protozoa, Ciliophora) from South Korea. *Journal of Species Research* 9:427–442.
- Kim, K.-S., K.-M. Park, J.-H. Jung and G.-S. Min. 2016. First description of three ciliates (Ciliophora: Stichotrichia) from Korea. *Journal of Species Research* 5(3):468–476.
- Lei, Y., J.K. Choi, K. Xu and W. Petz. 2005. Morphology and infraciliature of three species of *Metaurostylopsis* (Ciliophora, Stichotrichia): *M. songi* n. sp., *M. salina* n. sp., and *M. marina* (Kahl 1932) from sediments, saline ponds, and coastal waters. *Journal of Eukaryotic Microbiology* 52:1–10.
- Li, L., Q. Zhang, K.A.S. Al-Rasheid, C.B. Kwon and M.K. Shin. 2010. Morphological redescriptions of *Aspidisca magna* Kahl, 1932 and *A. leptaspis* Fresenius, 1865 (Ciliophora, Euplotida), with notes on ontogenesis in *A. magna*. *Acta Protozoologica* 49:327–337.
- Lu, B., C. Wang, J. Huang, Y. Shi and X. Chen. 2016. Morphology and SSU rDNA sequence analysis of two hypotrichous ciliates (Protozoa, Ciliophora, Hypotrichia)

- including the new species *Metaurostylopsis parastruederkyrkiae* n. sp. Journal of Ocean University of China 15(5):866-878.
- Lynn, D.H. 2008. The ciliated protozoa: Characterization, classification, and guide to the literature. New York, Springer.
- Pan, X., W.A. Bourland and W. Song. 2013. Protargol synthesis: An in-house protocol. Journal of Eukaryotic Microbiology 60:609-614.
- Park, K.-M., J.-H. Jung and G.-S. Min. 2012. Redescription of two urostylid ciliates (Ciliophora: Urostylida), *Anteholosticha pulchra* and *Metaurostylopsis struederkypkeae* from Korea. Animal Systematics, Evolution and Diversity 28(1):20-28.
- Shao, C., M. Miao, S. Weibo, W. Alan, K.A.S. Al-Rasheid, S.A. Al-Quraishi and S.A. Al-Farraj. 2008a. Studies on two marine *Metaurostylopsis* spp. from China with notes on the morphogenesis of *M. sinica* nov. spec. (Ciliophora, Urostylida). Acta Protozoologica 47:959-112.
- Shao, C., W. Song, K.A.S. Al-Rasheid, Z. Yi, X. Chen, S.A. Al-Farraj and S.A. Al-Quraishi. 2008b. Morphology and infraciliature of two new marine urostylid ciliates: *Metaurostylopsis struederkypkeae* n. sp. and *Thigmokeronopsis stoecki* n. sp. (Ciliophora, Hypotrichida) from China. Journal of Eukaryotic Microbiology 55:289-296.
- Shen, Z., J. Huang, X. Lin, Z. Yi, J. Li and W. Song. 2010. Morphological and molecular characterization of *Aspidisca hongkongensis* spec. nov. (Ciliophora, Euplotida) from the South China Sea. European Journal of Protistology 46:204-211.
- Shin, M.K. and H.S. Kim. 1988. Several hypotrichous ciliates inhabiting the Han River in Seoul. Korean Journal of Systematic Zoology Special Issue No. 2:67-85.
- Song, W., W. Petz and A. Warren. 2001. Morphology and morphogenesis of the poorly-known marine urostylid ciliate, *Metaurostylopsis marina* (Kahl, 1932) nov. gen., nov. comb. (Protozoa, Ciliophora, Hypotrichida). European Journal of Protistology 37:63-76.
- Song, W., Y. Qiao, J. Dong, W.A. Bourland, T. Zhang and X. Luo. 2020. Ontogeny and phylogeny of a new hypotrichous ciliate (Protista, Ciliophora), *Metaurostylopsis alrasheidi* n. sp., with establishment of a new genus *Monurostylopsis* n. gen. Frontiers in Marine Science 7:602317.
- Song, W. and N. Wilbert. 1997. Morphological investigations on some free living ciliates (Protozoa, Ciliophora) from China Sea with description of a new hypotrichous genus, *Hemigastrostyla* nov. gen. Archiv Für Protistenkunde 148: 413-444.
- Song, W. and N. Wilbert. 2002. Faunistic studies on marine ciliates from the Antarctic benthic area, including description of one epizoic form, 6 new species and, 2 new genera (Protozoa: Ciliophora). Acta Protozoologica 41:23-61.
- Valbonesi, A. 1996. Description of a new species of *Aspidisca*, *Aspidisca terranova* sp. n., from Antarctica (Ciliophora, Hypotrichida). Italian Journal of Zoology 63 (4):377-380.
- Wu, I.C.H. and C.R. Curds. 1979. A guide to the species of the genus *Aspidisca*. Bull. Br. Mus. nat. Hist. (Zool.) 36:1-34.
- Zheng, B., X. Zhao, T. Ye, J. Huang, A. Vallesi, Y. Jiang and X. Chen. 2022. Morphology and molecular phylogeny of two new *Aspidisca* species (Ciliophora, Spirotrichea, Euplotida) collected from subtropical coastal waters in China. Frontiers Marine Science 9:970692.

Submitted: July 9, 2024

Revised: August 6, 2024

Accepted: August 6, 2024