

Prediction of apparent total tract digestion of crude protein in adult dogs

Kangmin Seo[#], Hyun-Woo Cho[#], Min Young Lee, Chan Ho Kim, Ki Hyun Kim and Ju Lan Chun^{*}

Animal Welfare Research Team, National Institute of Animal Science, Rural Development Administration, Wanju 55365, Korea



Received: Jan 11, 2024
Revised: Feb 13, 2024
Accepted: Feb 14, 2024

[#] These authors contributed equally to this work.

*Corresponding author

Ju Lan Chun

Animal Welfare Research Team,
National Institute of Animal Science,
Rural Development Administration,
Wanju 55365, Korea.
Tel: +82-63-238-7053
E-mail: julanchun@korea.kr

Copyright © 2024 Korean Society of Animal Sciences and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Kangmin Seo

<https://orcid.org/0000-0001-6152-8536>

Hyun-Woo Cho

<https://orcid.org/0000-0002-3620-9952>

Min Young Lee

<https://orcid.org/0000-0003-4860-6290>

Chan Ho Kim

<https://orcid.org/0000-0003-2121-5249>

Ki Hyun Kim

<https://orcid.org/0000-0002-9834-2126>

Ju Lan Chun

<https://orcid.org/0000-0002-4618-586X>

Competing interests

No potential conflict of interest relevant to this article was reported.

Abstract

To predict the apparent total tract digestibility (ATTD) of crude protein (CP) in dogs we developed an *in vitro* system using an *in vitro* digestion method and a statistical analysis. The experimental diets used chicken meat powder as the protein source, with CP levels of 20% (22.01%, analyzed CP value as dry-based), 30% (31.35%, analyzed CP value as dry-based), and 40% (41.34%, analyzed CP value as dry-based). To simulate *in vivo* digestive processes a static *in vitro* digestion was performed in two steps; stomach and small intestine. To analyze ATTD the total fecal samples were collected in eight neutered beagle dogs during the experimental period. CP digestibility was calculated by measuring CP levels in dog food, *in vitro* undigested fraction, and dog feces. In result, CP digestibility at both *in vivo* and *in vitro* was increased with increasing dietary CP levels. To estimate *in vivo* digestibility the co-relation of *in vivo* ATTD and *in vitro* digestibility was investigated statistically and a regression equation was developed to predict the CP ATTD ($\% = 2.5405 \times \textit{in vitro}$ CP digestibility (%) + 151.8). The regression equation was evaluated its feasibility by using a commercial diet. The predicted CP digestibility which was calculated by the regression equation showed high index of similarity (100.16%) with that of *in vivo* in dogs. With that, it would be a feasible non-animal method to predict *in vivo* CP digestibility by using *in vitro* digestion method and the proposed linear regression equation in adult dogs.

Keywords: Dog food, Crude protein, *In vitro* digestion method, Linear regression equation, Prediction of CP apparent total tract digestibility (ATTD)

INTRODUCTION

The scale of the pet food industry has rapidly expanded as an increasing number of households worldwide have companion animals. Moreover, the change in the perception of companion animals (humanization trends, e.g., the pet treated as a member of a family) has prompted consumers to take an increasing interest in the quality of the diet of their pet [1]. Dog foods are composed of main nutrients such as proteins, lipids, carbohydrates, and trace nutrients such as minerals and vitamins. The European Pet Food Industry Federation (FEDIAF), National Research Council (NRC), and American Feed Control Officials (AAFCO) provide information on the recommended level of each nutrient for a

Funding sources

This work was supported out with the support of "Cooperative Research Program for Agriculture Science and Technology Development (PJ01569901)", Rural Development Administration, Republic of Korea.

Acknowledgements

This study was supported by 2024 Postdoctoral Fellowship Program of National Institute of Animal Science, Rural Development Administration, Korea.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Chun JL.
Data curation: Lee MY, Kim KH, Chun JL.
Formal analysis: Seo K, Cho HW, Kim CH.
Methodology: Seo K, Cho HW, Kim KH, Chun JL.
Software: Kim CH.
Validation: Lee MY, Kim KH, Chun JL.
Investigation: Seo K, Cho HW.
Writing - original draft: Seo K, Cho HW.
Writing - review & editing: Seo K, Cho HW, Lee MY, Kim CH, Kim KH, Chun JL.

Ethics approval and consent to participate

This study was approved by the Animal Care and Use Committee of the National Institute of Animal Science, Wanju, Republic of Korea (NIAS-2021-513, approved on 30 March 2021).

balanced nutrient supply for companion animals [2–4]. Proteins, in particular, are an essential nutrient, supplying essential amino acids necessary for the growth and metabolism of companion dogs [5]. Hence, the level and digestibility of crude protein (CP) can be an important indicator for evaluations of the dog food quality [6]. CP digestibility data can also offer information on nutrient bioavailability to consumers to enhance the reliability of the dog food quality and provide value from the perspective of animal health and welfare [7].

Ethical issues of animal experimentation have increased worldwide. Efforts to support replacement, reduction, or refinement (3Rs) of animal use have been made in research. Moreover, the Food and Drug Administration in the US has announced that alternate methods can be used in preclinical research to determine the safety and efficacy of drug before human trials. Moreover, it is especially difficult to use dogs for experimentation because many people consider dogs as a companion animal and a member of the family. Regardless, evaluation of the nutritional value of foods (raw materials) for companion animals demands abundant time, financial investment, and animal experiments [8]. In the pet food industry, these limitations entail practical challenges in product development, quality control, and data provision for consumers through adequate assessments of food value in the pet food industry [9]. Therefore, it is necessary to develop an alternate method to research the nutritional utilization in dogs without animal experimentation.

The biological accessibility of nutritive or non-nutritive factors and nutrient digestibility can be analyzed using an *in vitro* digestion method that can simulate the digestive process of the body, with advantages such as controlled selectivity and high reproducibility [6,10,11]. Therefore, this method is suitable for exploring and proving new hypotheses in nutritional research. Furthermore, it has attracted attention as an alternative to animal experiments in fields related to food and nutrition, which could help avoid the recent ethical debates [12]. *In vitro* digestion models are also advantageous because they allow the selective simulation of physiological conditions (pH, temperature, enzymes, and microorganisms) in the digestive tract of an animal (oral cavity, stomach, small intestine, and ileum), in addition to being easy to use, low-cost, and independent of special devices [13]. Previous studies on *in vitro* digestion have included a wide spectrum of research subjects ranging from humans [14–16] to livestock [17–19] and fish [20]. Some sophisticated computerized *in vitro* digestion models can even simulate physical or physiological aspects, from the rate of food migration to enzyme concentrations and pH changes [21,22]. Notably, different *in vitro* digestion models have been reported for predicting digestibility in economically important animals, such as livestock [17,23,24], whereas very few studies have been conducted to predict digestibility in companion dogs. Studies on companion dogs using *in vitro* digestion models have presented regression equations to predict the digestibility of several commercial diets [6,25,26]. However, they have limitations in reflecting the characteristics of protein digestion observed *in vivo* because of the diversity of protein sources in commercial diets [27]. Hence, using regression equations to predict digestibility, the digestibility of CP can be estimated for a wide range of dog foods; however, precise prediction might not always be possible.

Therefore, this study was performed to evaluate the potential feasibility of an *in vitro* method for predicting the apparent total tract digestibility (ATTD) of CP in dogs via a static *in vitro* digestion approach and a regression equation.

MATERIALS AND METHODS

Animals

All animal experiments were conducted in compliance with the methods approved by the Institutional Animal Care and Use Committee of the National Institute of Animal Science (NIAS),

Republic of Korea (approval number: NIAS-2021-513). All experiments were conducted on dogs owned by the NIAS (National Institute of Animal Science, Rural development Administration, Republic of Korea). All dogs were provided with an independent space (170 × 210 cm/dog) with a constant indoor temperature (22 °C–23 °C) and lighting (16 h light and 8 h dark cycle). The animals were provided with drinking water *ad libitum* and fed twice daily. The food intake was also measured daily. Each animal was provided with approximately 3 h of outdoor activity. The health of the dogs was monitored daily and was checked by a veterinarian in NIAS, if needed.

Experimental and commercial diets

The experimental diets were designed to meet the nutritional requirements suggested by AAFCO [28] for adult dogs (Table 1). The CP level in diets was formulated to 20% (22.01%, Analyzed CP value as dry-based), 30% (31.35%, Analyzed CP value as dry-based), and 40% (41.34%, Analyzed CP value as dry-based) [28], and the diet was prepared according to the procedure reported by Seo et al. [29]. All ingredients used in the experimental diets were purchased as commercial powdered products without any palatants. As Table 1 shows, all ingredients were mixed, steamed, and formulated. After preparation, the experimental diet was stored at –20 °C until use. The food was thawed and warmed to room temperature immediately prior to feeding. An extruded dry type of commercial dog food was purchased from a local brand in the Republic of Korea (chicken meat and rice flour as the base) containing 28.02% experimental CP (analyzed as the dry-based value) and was used to validate the linear regression equation for predicting the ATTD of CP. The chemical composition of the commercial dog diet is listed in Table 1.

In vivo digestibility

Feeding test

The CP ATTD of the experimental diet was evaluated using eight healthy beagle dogs (5 years old; neutered 4 males, spayed 4 females). Food supply was set to meet the maintenance energy requirements estimated using the equation (metabolizable energy [ME], 132 kcal kg body weight [BW]^{0.75} per day) suggested by the Association of American Feed Control Officials [28]. Experimental diets were provided to dogs for two weeks including three days to adjust it and four days to collect feces.

Apparent total tract digestibility

The total collection method was used to analyze the *in vivo* CP ATTD. Fecal samples were collected for four days and weighed daily during the study period. Fecal samples were stored at –20 °C until analyzed. To analyze the levels of water (Association of Official Analytical Chemists [AOAC] method 934.01) and CP (AOAC method 984.13) in fecal and food samples, the methods established by AOAC [30] were followed. The CP ATTD was calculated using equation 1 [31]:

$$\text{Equation 1: CP ATTD (\%)} = \left(\frac{\text{CP input (food)} - \text{CP output (fecal)}}{\text{CP input (food)}} \right) \times 100$$

In vitro digestibility

A static *in vitro* digestion

The *in vitro* digestion method comprised two digestive steps, in reference to the static *in vitro* digestion method suggested by Seo et al. [13]. Digestion progresses sequentially from the gastric digestion phase to the small intestinal phase. The experimental diet was dried in a dry oven (65 °C) and ground to a constant weight (<1 mm particle size). For the gastric digestion phase, a 10 g

Table 1. Ingredients and calculated chemical compositions of experimental and commercial diets

Items	Crude protein concentrations			Commercial dog food ¹⁾
	20%	30%	40%	
Ingredient composition (%)				
Rice flour	43.91	37.04	30.04	-
Chicken breast powder	6.60	14.00	21.60	-
Egg yolk powder	8.00	8.00	7.50	-
Lard	1.50	1.30	1.40	-
Seaweed (Enteromorpha)	1.00	1.00	1.00	-
Cabbage powder	1.00	1.00	1.00	-
Calcium phosphate	1.00	0.80	0.60	-
Calcium carbonate	0.70	0.80	0.89	-
Potassium citrate	0.51	0.35	0.30	-
Vitamin–mineral premix ²⁾	0.40	0.40	0.40	-
Salt	0.33	0.31	0.27	-
Tryptophan	0.05	0.00	0.00	-
Water	35.00	35.00	35.00	-
Chemical composition (% [DM basis, calculated])				
Dry matter	58.56	59.49	60.45	≤ 12
Crude protein	20.02	30.16	39.99	≥ 26
Ether extract	11.75	12.02	12.35	≥ 12
Crude fiber	0.29	0.28	0.28	≤ 4.5
Crude ash	2.19	2.47	2.68	≤ 7.0
NFE	65.75	55.07	44.70	-
Ca	0.81	0.81	0.80	≥ 1.0
P	0.69	0.68	0.67	≤ 2.0
Ca/P ratio	1.18	1.19	1.20	-
ME (kcal/kg)	4,001	4,005	4,014	3,732

These values were calculated as dry-based values.

All the experimental diets used chicken powder as the protein source, and each experimental diet was formulated to contain 20%, 30%, and 40% CP.

¹⁾The commercial food was an extruded dry type (chicken and rice base) for adult dogs, and information on its chemical composition was the same as that provided by the manufacturer.

²⁾Provided per kilogram of experimental diets: vitamin A, 5,250 IU; vitamin D₃, 375 IU; vitamin E, 37.5 mg; vitamin K, 0.078 mg; vitamin B₁ (thiamine), 4.2 mg; vitamin B₂ (riboflavin), 3.9 mg; vitamin B₆ (pyridoxine), 3 mg; vitamin B₁₂, 0.021 mg; D-calcium pantothenate, 9 mg; niacin, 45 mg; folic acid, 0.6 mg; biotin, 0.054 mg; taurine, 1,500 mg; FeSO₄·H₂O, 66 mg; MnSO₄·H₂O, 5.7 mg; ZnSO₄·H₂O, 75 mg; CuSO₄·H₂O, 11.25 mg; Na₂SeO₃, 0.27 mg; Ca(IO₃)₂, 1.35 mg.

CP, crude protein; DM, dry matter; EE, ether extract; NFE, nitrogen-free extract; ME, metabolizable energy (kcal/kg DM) = [(CP × 3.5) + (EE × 8.5) + (NFE × 3.5)] × 10.

sample (<1 mm), 250 mL (25 mL/g feed) of 0.1 M phosphate buffer (pH, 6.0), and 100 mL (10 mL/g feed) of 0.2 M HCl were placed in an Erlenmeyer glass flask (volume, 1,000 mL). The pH of the mixture was adjusted to 2.0 using 0.1 M HCl. Next, 1 mL of pepsin-HCl solution (100 mg/mL of 0.075 N HCl solution and pepsin from the porcine gastric mucosa, ≥ 250 units/mg, #P7000, Sigma-Aldrich, St. Louis, MO, USA) and 1 mL of chloramphenicol solution (2.5 mg/mL EtOH, #C-0378, Sigma-Aldrich) were added, and the mixture was incubated in a shaking incubator (39°C, 150 rpm) for 6 h. For the small intestinal phase, 100 mL (10 mL/g feed) of 0.2 M phosphate buffer (pH, 6.8) and 50 mL (5 mL/g feed) of 0.6 M NaOH were added to the digestive fluid after the gastric digestion phase. The pH of the mixture was adjusted to 7.5 using 0.1 M NaOH. Next, 12.5 g of bile salts (1.25 g/g feed, #13805, Sigma-Aldrich) and 2.50 g of pancreatin (0.25 g/g feed, #P7545, 8 × USP specifications, Sigma-Aldrich) were added, and the mixture

was incubated in a shaking incubator (39°C, 180 rpm) for 18 h. At the end of the entire digestive process, the undigested fraction was collected using a bottle top vacuum filter (pore size, 0.22 µm; polyethersulfone membranes, TPP Techno Plastic Products AG, Zurich, Switzerland) and dried in a dry oven (65°C) for subsequent analysis.

Analysis of *in vitro* crude protein digestibility

For the CP level and digestibility, the CP content (AOAC method 984.13) was measured in the undigested fraction from a static *in vitro* simulation of digestion and experimental diet, following the methods established by AOAC [30]. For these calculations, equation 2 was used [13].

$$\text{Equation 2: } \textit{in vitro} \text{ CP digestibility (\%)} = \left(\frac{\text{Food CP (g)} - \text{Undigested fraction CP (g)}}{\text{Food CP (g)}} \right) \times 100$$

Establishment of a regression equations for predicting the apparent total tract digestibility of crude protein in adult dogs

The linear regression equation for predicting the ATTD of CP in adult dogs was determined based on the co-relation between the average *in vitro* and *in vivo* CP digestibility data for the experimental diet (CP, 20%–40%). Briefly, to evaluate the regression equation, a commercial diet of the extruded dry type (chicken meat and rice flour as the base) containing 28.02% CP (analyzed value as dry-based; Table 1) was used.

Evaluation of the non-animal method to predict *in vivo* crude protein digestibility in dogs

First, a commercial diet was assessed the *in vitro* digestion process and the CP level in undigested fraction was measured using the same methods as those used for the experimental diet. *In vitro* digestibility was calculated according to Equation 2. Second, CP ATTD was analyzed by feeding the commercial diet to nine healthy beagle dogs (5 years old, neutered 1 male and spayed 8 females). Total feces were collected and CP level was measured. CP ATTD was calculated according to Equation 1. Third, the *in vitro* CP digestibility data for the commercial diet were applied to the regression equation to predict the ATTD of CP. In final, the predicted *in vivo* CP digestibility was compared with the ATTD of CP for the commercial diet tested using nine healthy beagle dogs. The index of similarity between the predicted digestibility and ATTD of CP was calculated according to equation 3 [32]. The index of similarity value closer to 100 means more similar between the two CP digestibility value.

$$\text{Index of similarity (\%)} = \left(\frac{\text{Predicted CP digestibility}}{\text{ATTD of CP}} \right) \times 100$$

Statistical analysis

SPSS (ver. 17.0, 2008; SPSS Statistics, Chicago, IL, USA) software was used to perform all statistical analyses, and the results were expressed as the mean ± standard error of the mean. One-way analysis of variance was used to analyze the changes in food intake and digestibility according to the variation in CP levels in the dog food. Tukey's multiple comparison test was used to test the significance of differences between groups. The Student's *t*-test was used to analyze the differences between *in vitro* and *in vivo* CP digestibility. The level of significance was set at $p < 0.05$. To analyze the correlation between CP levels and *in vitro* or *in vivo* CP digestibility, Pearson's bivariate correlation analysis was performed.

RESULTS

Differences in crude protein digestibility between *in vitro* and *in vivo* conditions

To assess *in vivo* CP digestibility, ATTD was evaluated in beagle dogs fed nutritionally controlled experimental diets. During the study period, the mean daily food intake, metabolic energy intake (MEI), and MEI per metabolic body weight (mBW) did not differ significantly among the experimental groups ($p > 0.05$; Table 2). In contrast, the mean daily CP intake was different among the groups fed an experimental diet with 20%, 30%, and 40% CP, with values of 46.48 ± 1.20 , 62.50 ± 2.46 , and 78.65 ± 5.43 g/day, respectively (Table 2). The mean daily CP intake increased according to the CP level in the experimental diet ($p < 0.001$).

The results of ATTD of CP was $89.26 \pm 0.60\%$ when fed 20% CP diet, indicative of the lowest digestibility when compared to that of the experimental diet with 30% CP ($92.19 \pm 0.68\%$) or 40% CP ($94.02 \pm 0.40\%$) ($p < 0.05$; Fig. 1A).

Next, CP digestibility assessed using the static *in vitro* digestion method was $94.86 \pm 0.24\%$, $96.13 \pm 0.23\%$, and $96.69 \pm 0.15\%$ for the experimental diets with 20%, 30%, and 40% CP diets, respectively. *In vitro* CP digestibility thus tend to increase in a concentration-dependent manner in diets containing 20%, 30%, or 40% CP ($p < 0.05$; Fig. 1A). In addition, CP digestibility was higher in the *in vitro* digestion method than that in ATTD for all experimental diets (20%–40% CP) ($p < 0.05$; Fig. 1A).

Correlation between the protein concentration in experimental diets and crude protein digestibility

CP digestibility under both *in vitro* and *in vivo* conditions tended to increase with increasing CP concentrations in the experimental diets (Fig. 1A). To determine the correlation between the CP concentration in the experimental diet and *in vitro* or *in vivo* digestibility, Pearson's correlation analysis was performed. The correlation coefficient between *in vitro* CP digestibility and CP levels in the experimental diet was 0.911 ($p = 0.001$; Fig. 1B). The coefficient between *in vivo* CP digestibility and the CP level in the experimental diet was 0.738 ($p < 0.001$; Fig. 1C).

A linear regression equation to predict crude protein apparent total tract digestibility

The *in vivo* and *in vitro* CP digestibility data of 20%, 30%, and 40% CP contained diets were subjected to a linear regression analysis to predict the ATTD of CP in adult dogs. The linear relationship among the three different CP concentrations is shown in Fig. 2, and the regression equation was calculated using equation 3.

Table 2. Food and ME intake

Items	CP concentrations (%)			p-value
	20	30	40	
ADFI (g/day)	309.46 ± 7.99^1	295.23 ± 11.62	284.97 ± 19.67	0.47
CP intake (g/day, DM)	46.48 ± 1.20^c	62.50 ± 2.46^b	78.65 ± 5.43^a	< 0.001
MEI (kcal/day)	725.50 ± 18.74	703.46 ± 27.70	690.87 ± 47.68	0.77
Calculated MEI/kg mBW	112.61 ± 0.00	109.14 ± 2.93	107.01 ± 6.41	0.62

Adult beagle dogs (n = 8/group) were fed experimental diets for 10 days.

All experimental diets had chicken powder as the protein source, and each experimental diet was formulated to contain 20%, 30%, and 40% CP.

¹Values are expressed as the mean \pm standard error of the mean (SEM).

^{a-c}Data without the same superscript numbers in the same row differ significantly ($p < 0.05$).

ME, metabolizable energy; ADFI, average daily food intake; MEI, metabolic energy intake; mBW, metabolic body weight.




Fig. 1. Effect of crude protein (CP) levels in the experimental diet (dry matter basis) on *in vitro* and *in vivo* CP digestibility. (A) Differences between *in vitro* and *in vivo* CP digestibility. The linear correlations between the experimental diets with the three protein levels (20%–40%) and (B) *in vitro* (n = 3) or (C) *in vivo* (n = 8) CP digestibility are shown. All experimental diets used chicken powder as the protein source, and each experimental diet was formulated to contain 20%, 30%, or 40% CP. *In vivo* CP digestibility values of experimental diets were evaluated in beagle dogs. The values are expressed as the mean ± standard error of the mean (SEM). ^{A,B}Significant differences in CP digestibility among dog diet groups containing 20%–40% CP in terms of *in vitro* digestion (p < 0.05). ^{a,b}Significant differences in CP digestibility among dog diet groups containing 20%–40% CP in terms of *in vivo* digestion (p < 0.05). *Significant differences in CP digestibility between *in vitro* and *in vivo* digestion (p < 0.05).




Fig. 2. Linear regression equation for predicting the apparent total tract digestibility (ATTD) of crude protein (CP) in adult dogs. Relationship between average *in vivo* and *in vitro* CP digestibility for experimental diets containing 20%, 30%, and 40% CP. Values are expressed as the mean ± SEM.

Table 3. Evaluation of regression equations for predicting CP of ATTD in adult dogs

Items	Commercial diet ¹⁾ (%)
<i>In vitro</i> CP digestibility	92.82 ± 0.09
Predicted <i>in vivo</i> CP digestibility (P) ²⁾	84.00 ± 0.23
Analyzed <i>in vivo</i> CP digestibility (A) ³⁾	83.87 ± 0.65
Index of similarity ⁴⁾	100.16

¹⁾The commercial diet used was a chicken- and rice-based adult dog diet (CP, 28.02%; analyzed as a dry-based value).

²⁾Predicted *in vivo* CP digestibility value was calculated by applying the *in vitro* CP digestibility value to the equation for predicting *in vivo* CP digestibility.

³⁾The *in vivo* CP digestibility value was evaluated for ATTD of the commercial diet in beagle dogs (n = 9). Values are expressed as the mean and SEM.

⁴⁾The index of similarity of CP digestibility was calculated as (P) / (A) × 100.

CP, crude protein; ATTD, apparent total tract digestibility.

Equation 3: equation for predicting ATTD of CP (%) = 2.5405 × *in vitro* CP digestibility (%) + 151.8

Evaluation of the linear regression equations for predicting crude protein apparent total tract digestibility

The regression equation developed to predict CP digestibility was validated using a commercial diet, which was an extruded dry type (chicken meat and rice flour as the base) containing 28.02% CP (analyzed value as dry-based). The CP digestibility of the commercial diet was 92.82 ± 0.09%, as calculated using the static *in vitro* digestion method. The CP digestibility results were applied to the regression equation (Eq. 3) and the *in vitro* CP digestibility was calculated to be 84.00 ± 0.23% (Table 3). Meanwhile, to obtain the *in vivo* CP digestibility of the same commercial diet used for *in vitro* digestibility, a feeding test was performed in dogs and the *in vivo* CP digestibility was calculated to be 83.87 ± 0.65% (Table 3). The index of similarity between the predicted CP digestibility using the regression equation after *in vitro* digestion and the measured ATTD of CP in dogs was shown to be 100.16 ± 0.65% (Table 3).

DISCUSSION

Data on nutrient bioavailability in the food are essential to provide well-balanced pet foods [6,7]. Proteins, in particular, comprise essential amino acids necessary for growth, homeostasis, and consistent immunity in companion animals. Amino acid content and types vary across different protein-based raw ingredients, and the *in vivo* amino acid availability or requirement can vary according to the type of ingested protein source [33]. Hence, it is necessary to evaluate the nutritional value of the food based on the protein source present in the dog food. Evaluations of ATTD have been widely used as a general method to obtain the value of *in vivo* CP digestibility [34]; however, this method requires time and is costly, with the most serious drawback being the social concern regarding animal experimentation. Consequently, despite consumer demand, the dog food industry is struggling to provide accurate and efficient data on the *in vivo* nutrient utilization of dog foods. Therefore, this study suggested the combination of a static *in vitro* simulation of digestion and a linear regression equation to improve the evaluation method for *in vivo* nutrient utilization, especially for proteins. As a result, the estimated *in vivo* digestibility of CP was highly similar to the *in vitro* digestibility of CP from ATTD in dogs.

To investigate the correlation between the estimated *in vitro* digestibility and measured *in vivo* digestibility, the digestibility of experimental diets containing varying levels of CP was assessed under *in vitro* and *in vivo* conditions. Experimental diets with 20%, 30%, and 40% CP

concentrations were prepared using chicken meat powder, and the variation in the contents of other nutrients was minimized by applying the nutritional requirements of AAFCO. *In vitro* digestibility was evaluated by a static *in vitro* simulation of digestion and *in vivo* digestibility was calculated using ATTD in dogs. The results of this study revealed that, for all experimental diets, CP digestibility was higher under *in vitro* conditions than under *in vivo* conditions. *In vivo* digestion is influenced by various parameters (e.g., protein types, bacteria, and endogenous proteins) and complex digestive mechanisms inside the body. For example, *in vivo* digestion is a dynamic process characterized by pH changes and the continuous production and release of digestive enzymes that occur with the flow of food through different compartments of the digestive organs (oral cavity, stomach, small intestine, and ileum) [12]. In contrast, *in vitro* digestion was performed under strictly controlled conditions of pH, temperature, enzyme content, and digestion time [6,25,26,35,36,37]. Therefore, the different degrees of digestibility under these two conditions can be explained. Despite clear limitations in simulating an actual digestive process using a static *in vitro* digestion model [38], this study demonstrated a correlation between *in vitro* and *in vivo* digestibility.

The results of this study also demonstrate a trend of increasing CP digestibility under the two digestion conditions (*in vitro* and *in vivo*), in line with the increase in CP concentration in the experimental diet. Pearson's correlation analysis showed that, under both *in vitro* ($r = 0.911$, $p = 0.001$) and *in vivo* ($r = 0.783$, $p < 0.001$) conditions, CP digestibility exhibited a significant linear correlation with CP levels in the experimental diet. This implies a definite correlation between the CP levels in dog food and CP digestibility. A previous study by Yamka et al. [39] reported similar findings *in vivo*. They examined the digestibility of foods (low-ash poultry meal base) containing varying concentrations of CP (10%–25%) in healthy adult dogs and showed that an increase in the CP level in the food increases CP digestibility [39]. Thus, the trend of increasing CP digestibility with the increase in CP concentration was observed. Moreover, the findings of this study support a strong correlation of digestibility between *in vitro* digestion using a static *in vitro* simulation method and *in vivo* digestion using ATTD.

For reasons related to their advantageous speed and cost, prediction models for *in vivo* digestibility using regression analysis are being widely investigated for economically important animals (cattle, poultry, pigs, and rabbits), companion animals, and humans [6,40–42]. Prediction models for *in vivo* digestibility using regression analysis have not only identified nutrients in foods, including carbohydrates [24], CP [43], and phosphorus [44], but have also verified the potential feasibility of prediction models using regression analysis for energy [23] and the postprandial glycemic index [45]. Nevertheless, there have been few studies on prediction models for *in vivo* digestibility using regression analysis in dogs. Biagi et al. [6] assessed 16 commercial diets (extruded dry type) and found that the coefficient of determination of the linear regression equation was low for the estimation of CP ATTD in adult dogs ($r^2 = 0.510$) [6]. Kawauchi et al. [38] proposed a CP digestibility prediction model based on 16 dog foods, including meat, bone meal (MBM), and poultry by-product meal (PM). Interestingly, they proposed an independent equation for each of the two protein-based raw materials; however, the reliability of both equations was low (MBM: $r^2 = 0.126$; PM: $r^2 = 0.216$) [38]. In this study, the regression equation was generated based on the correlation results of both static *in vitro* simulation and *in vivo* ATTD and showed a highly reliable coefficient value for the CP prediction equation for dogs ($r^2 = 0.9925$). Moreover, the predicted CP digestibility using the regression analysis with the data from the static *in vitro* digestion of CP had a significantly high similarity (100.16%) with that of *in vivo* CP digestibility in dogs. This suggests that the regression equation proposed in this study has potential utility in predicting ATTD of CP in adult dogs. To test the reproducibility of the regression equation, a commercial

dog food (extruded dry type) from a local brand in the Republic of Korea was used in this study. As the experimental diet was produced in lab, the digestibility of the commercial dog food was assessed using two methods: a static *in vitro* simulation method and ATTD in dogs. The *in vitro* digestibility was predicted using a regression equation and the *in vivo* digestibility was calculated as ATTD in dogs. The predicted CP digestibility of the commercial dog food using the regression equation was not significantly different from the *in vivo* ATTD measured in dogs and showed $100.16 \pm 0.65\%$ similarity. Therefore, the regression equation is suitable not only for experimental diets produced in the lab but also for commercially produced extruded dry type diets to predict the *in vivo* digestibility.

Animal feeding tests in the nutritional research field is a common and essential process, although it requires prohibitive cost, a safe animal facility, animal experts, and technical support. A static *in vitro* simulation of digestion is cost effective and requires minimum experimental skill and lab space, unlike animal experiments. The combination of a static *in vitro* simulation of digestion method and the regression analysis allows researchers to obtain information on the digestibility of CP without the necessity of animal experimentation. Thus, the data in this study are expected to prove the nutritional value of proteins for the development and production of dog food and in fields related to nutrition, where *in vivo* digestibility must be predicted according to varying protein content.

Evaluation of *in vivo* utilization of nutrients, especially proteins, is crucial not only for companion dog health management but also for the use of an optimized level of high-cost protein-based raw materials, contributing to more reasonable purchases by consumers. However, the pet food industry claims that it is difficult to evaluate the *in vivo* utilization of nutrients when they develop or produce pet food products because of the requirement of animal experiments. Therefore, the non-animal method to predict *in vivo* digestibility using an *in vitro* digestion technique and regression analysis is expected to satisfy both the animal feed industry and consumers. In conclusion, this approach may be a simple and reproducible potential method for assessing the CP digestibility of dog diet, but a further research considering the characteristics of the various protein sources that dog diet may contain is absolutely necessary.

REFERENCES

- Hill M, Shanoyan A, Aldrich G. Animal protein-based ingredients in pet food: analysis of supply chain and market drivers. In: 2022 Agricultural & Applied Economics Association Annual Meeting; 2022; Anaheim, CA.
- AAFCO [Association of American Feed Control Officials]. Official publication. Vol. 2021. Champaign, IL: AAFCO; 2021. p. 111-236.
- NRC [National Research Council]. Nutrient requirements of dogs and cats. Washington, DC: National Academies Press; 2006.
- FEDIAF [The European Pet Food Industry]. Nutritional guidelines for complete and complementary pet food for cats and dogs. Bruxelles: FEDIAF; 2021.
- Laflamme DP. Pet food safety: dietary protein. *Top Companion Anim Med.* 2008;23:154-7. <https://doi.org/10.1053/j.tcam.2008.04.009>
- Biagi G, Cipollini I, Grandi M, Pinna C, Vecchiato CG, Zaghini G. A new *in vitro* method to evaluate digestibility of commercial diets for dogs. *Ital J Anim Sci.* 2016;15:617-25. <https://doi.org/10.1080/1828051X.2016.1222242>
- Case LP, Carey DP, Hirakawa DA, Daristotle L. Canine and feline nutrition: a resource for companion animal professionals. 2nd ed. St. Louis, MO: Mosby; 2000.
- McCusker S, Buff PR, Yu Z, Fascetti AJ. Amino acid content of selected plant, algae and insect

- species: a search for alternative protein sources for use in pet foods. *J Nutr Sci*. 2014;3:e39. <https://doi.org/10.1017/jns.2014.33>
9. Corsato Alvarenga I, Aldrich CG. The effect of increasing levels of dehulled faba beans (*Vicia faba* L.) on extrusion and product parameters for dry expanded dog food. *Foods*. 2019;8:26. <https://doi.org/10.3390/foods8010026>
 10. Vors C, Capolino P, Guérin C, Meugnier E, Pesenti S, Chauvin MA, et al. Coupling in vitro gastrointestinal lipolysis and Caco-2 cell cultures for testing the absorption of different food emulsions. *Food Funct*. 2012;3:537-46. <https://doi.org/10.1039/C2FO10248J>
 11. Theysgeur S, Cudennec B, Deracinois B, Perrin C, Guiller I, Lepoudère A, et al. New bioactive peptides identified from a tilapia byproduct hydrolysate exerting effects on DPP-IV activity and intestinal hormones regulation after canine gastrointestinal simulated digestion. *Molecules*. 2020;26:136. <https://doi.org/10.3390/molecules26010136>
 12. Minekus M, Alminger M, Alvito P, Ballance S, Bohn T, Bourlieu C, et al. A standardised static in vitro digestion method suitable for food – an international consensus. *Food Funct*. 2014;5:1113-24. <https://doi.org/10.1039/C3FO60702J>
 13. Seo K, Cho HW, Jeon JH, Kim CH, Lim S, Jeong S, et al. Influence of bile salts and pancreatin on dog food during static in vitro simulation to mimic in vivo digestion. *Animals*. 2022;12:2734. <https://doi.org/10.3390/ani12202734>
 14. Guerra A, Etienne-Mesmin L, Livrelli V, Denis S, Blanquet-Diot S, Alric M. Relevance and challenges in modeling human gastric and small intestinal digestion. *Trends Biotechnol*. 2012;30:591-600. <https://doi.org/10.1016/j.tibtech.2012.08.001>
 15. Calvo-Lerma J, Fornés-Ferrer V, Heredia A, Andrés A. In vitro digestion of lipids in real foods: influence of lipid organization within the food matrix and interactions with nonlipid components. *J Food Sci*. 2018;83:2629-37. <https://doi.org/10.1111/1750-3841.14343>
 16. Brodkorb A, Egger L, Alminger M, Alvito P, Assunção R, Ballance S, et al. INFOGEST static in vitro simulation of gastrointestinal food digestion. *Nat Protoc*. 2019;14:991-1014. <https://doi.org/10.1038/s41596-018-0119-1>
 17. Boisen S, Fernández JA. Prediction of the total tract digestibility of energy in feedstuffs and pig diets by in vitro analyses. *Anim Feed Sci Technol*. 1997;68:277-86. [https://doi.org/10.1016/S0377-8401\(97\)00058-8](https://doi.org/10.1016/S0377-8401(97)00058-8)
 18. Mabweesh SJ, Cohen M, Arieli A. In vitro methods for measuring the dry matter digestibility of ruminant feedstuffs: comparison of methods and inoculum source. *J Dairy Sci*. 2000;83:2289-94. [https://doi.org/10.3168/jds.S0022-0302\(00\)75115-0](https://doi.org/10.3168/jds.S0022-0302(00)75115-0)
 19. Reggi S, Giromini C, Dell'Anno M, Baldi A, Rebutti R, Rossi L. In vitro digestion of chestnut and quebracho tannin extracts: antimicrobial effect, antioxidant capacity and cytomodulatory activity in swine intestinal IPEC-J2 cells. *Animals*. 2020;10:195. <https://doi.org/10.3390/ani10020195>
 20. Moyano FJ, Saénz de Rodríguez MA, Díaz M, Tacon AGJ. Application of in vitro digestibility methods in aquaculture: constraints and perspectives. *Rev Aquacult*. 2015;7:223-42. <https://doi.org/10.1111/raq.12065>
 21. Wickham M, Faulks R, Mills C. In vitro digestion methods for assessing the effect of food structure on allergen breakdown. *Mol Nutr Food Res*. 2009;53:952-8. <https://doi.org/10.1002/mnfr.200800193>
 22. Ménard O, Cattenoz T, Guillemin H, Souchon I, Deglaire A, Dupont D, et al. Validation of a new in vitro dynamic system to simulate infant digestion. *Food Chem*. 2014;145:1039-45. <https://doi.org/10.1016/j.foodchem.2013.09.036>
 23. Noblet J, Jaguelin-Peyraud Y. Prediction of digestibility of organic matter and energy in the

- growing pig from an in vitro method. *Anim Feed Sci Technol.* 2007;134:211-22. <https://doi.org/10.1016/j.anifeedsci.2006.07.008>
24. Weurding RE, Veldman A, Veen WAG, van der Aar PJ, Verstegen MWA. In vitro starch digestion correlates well with rate and extent of starch digestion in broiler chickens. *J Nutr.* 2001;131:2336-42. <https://doi.org/10.1093/jn/131.9.2336>
 25. Tonglet C, Jeusette I, Istasse L, Diez M. Prediction of protein digestibility in dog food by a multi-enzymatic method: a useful technique to develop. *J Anim Physiol Anim Nutr.* 2001;85:189-94. <https://doi.org/10.1046/j.1439-0396.2001.00334.x>
 26. Hervera M, Baucells MD, González G, Pérez E, Castrillo C. Prediction of digestible protein content of dry extruded dog foods: comparison of methods. *J Anim Physiol Anim Nutr.* 2009;93:366-72. <https://doi.org/10.1111/j.1439-0396.2008.00870.x>
 27. Dust JM, Grieshop CM, Parsons CM, Karr-Lilienthal LK, Schasteen CS, Quigley JD 3rd, et al. Chemical composition, protein quality, palatability, and digestibility of alternative protein sources for dogs. *J Anim Sci.* 2005;83:2414-22. <https://doi.org/10.2527/2005.83102414x>
 28. AAFCO [Association of American Feed Control Officials]. Official publication. Vol. 2016. Champaign, IL: AAFCO; 2016. p. 107-234
 29. Seo K, Cho HW, Chun J, Jeon J, Kim C, Kim M, et al. Evaluation of fermented oat and black soldier fly larva as food ingredients in senior dog diets. *Animals.* 2021;11:3509. <https://doi.org/10.3390/ani11123509>
 30. AOAC [Association of Official Analytical Chemists] International. Official methods of analysis of AOAC international. 18th ed. Gaithersburg, MD: AOAC International; 2006.
 31. Chen Y, Shen D, Zhang L, Zhong R, Liu Z, Liu L, et al. Supplementation of non-starch polysaccharide enzymes cocktail in a corn-miscellaneous meal diet improves nutrient digestibility and reduces carbon dioxide emissions in finishing pigs. *Animals.* 2020;10:232. <https://doi.org/10.3390/ani10020232>
 32. Lupatsch I, Kissil GW, Sklan D, Pfeffer E. Apparent digestibility coefficients of feed ingredients and their predictability in compound diets for gilthead seabream, *Sparus aurata* L. *Aquac Nutr.* 1997;3:81-9. <https://doi.org/10.1046/j.1365-2095.1997.00076.x>
 33. Zentek J, Fricke S, Hewicker-Trautwein M, Ehinger B, Amtsberg G, Baums C. Dietary protein source and manufacturing processes affect macronutrient digestibility, fecal consistency, and presence of fecal *Clostridium perfringens* in adult dogs. *J Nutr.* 2004;134:2158S-61S. <https://doi.org/10.1093/jn/134.8.2158S>
 34. Hervera M, Baucells MD, Blanch F, Castrillo C. Prediction of digestible energy content of extruded dog food by in vitro analyses. *J Anim Physiol Anim Nutr.* 2007;91:205-9. <https://doi.org/10.1111/j.1439-0396.2007.00693.x>
 35. Boisen S, Eggum BO. Critical evaluation of in vitro methods for estimating digestibility in simple-stomach animals. *Nutr Res Rev.* 1991;4:141-62. <https://doi.org/10.1079/NRR19910012>
 36. Crampton EW, Rutherford BE. Apparent digestibility of dietary protein as a function of protein level. *J Nutr.* 1954;54:445-51. <https://doi.org/10.1093/jn/54.3.445>
 37. Kawauchi IM, Sakomura NK, Pontieri CFF, Rebelato A, Putarov TC, Malheiros EB, et al. Prediction of crude protein digestibility of animal by-product meals for dogs by the protein solubility in pepsin method. *J Nutr Sci.* 2014;3:e36. <https://doi.org/10.1017/jns.2014.32>
 38. Bohn T, Carriere F, Day L, Deglaire A, Egger L, Freitas D, et al. Correlation between in vitro and in vivo data on food digestion. What can we predict with static in vitro digestion models? *Crit Rev Food Sci Nutr.* 2018;58:2239-61. <https://doi.org/10.1080/10408398.2017.1315362>
 39. Yamka RM, Jamikorn U, True AD, Harmon DL. Evaluation of low-ash poultry meal as a

- protein source in canine foods. *J Anim Sci.* 2003;81:2279-84. <https://doi.org/10.2527/2003.8192279x>
40. Zaefarian F, Cowieson AJ, Pontoppidan K, Abdollahi MR, Ravindran V. Trends in feed evaluation for poultry with emphasis on in vitro techniques. *Anim Nutr.* 2021;7:268-81. <https://doi.org/10.1016/j.aninu.2020.08.006>
 41. Tassone S, Fortina R, Mabrouki S, Hachana Y, Barbera S. Comparison of in vivo and in vitro digestibility in rabbits. *Animals.* 2021;11:3267. <https://doi.org/10.3390/ani11113267>
 42. Sousa R, Recio I, Heimo D, Dubois S, Moughan PJ, Hodgkinson SM, et al. In vitro digestibility of dietary proteins and in vitro DIAAS analytical workflow based on the INFOGEST static protocol and its validation with in vivo data. *Food Chem.* 2023;404:134720. <https://doi.org/10.1016/j.foodchem.2022.134720>
 43. Bryan DDSL, Classen HL. In vitro methods of assessing protein quality for poultry. *Animals.* 2020;10:551. <https://doi.org/10.3390/ani10040551>
 44. Soutar L, Coltherd JC, Steele VR, Staunton R, Carvell-Miller L, Hughes KR, et al. Comparisons of in vitro and in vivo digestibility assays for phosphorus in feline diets and associations with dietary nutrient content. *J Agric Food Chem.* 2021;69:10688-99. <https://doi.org/10.1021/acs.jafc.1c03308>
 45. Peng X, Liu H, Li X, Wang H, Zhang K, Li S, et al. Predicting the glycemic index of biscuits using static in vitro digestion protocols. *Foods.* 2023;12:404. <https://doi.org/10.3390/foods12020404>