

APPLICATIONS OF THE GAUSSIAN HYPERGEOMETRIC FUNCTION TO SOME SUBCLASSES OF ANALYTIC FUNCTIONS

ABEER O. BADGHAISH AND AMANI Z. BAJAMAL

ABSTRACT. In this paper, we derive the necessary and sufficient conditions for the Gaussian hypergeometric function to be in some subclasses of analytic functions.

1. Introduction

Let \mathcal{A} denote the family of functions that are analytic in the open unit disc $U = \{\zeta \in \mathbb{C} : |\zeta| < 1\}$ with normalized form

$$g(\zeta) = \zeta + \sum_{t=2}^{\infty} a_t \zeta^t. \quad (1.1)$$

We denote by \mathcal{S} the subclass of \mathcal{A} consisting of univalent functions. Further, let \mathcal{T} be a subclass of \mathcal{S} consisting of functions g of the form

$$g(\zeta) = \zeta - \sum_{t=2}^{\infty} a_t \zeta^t, \quad a_t \geq 0. \quad (1.2)$$

DEFINITION 1. [5] A function $g \in \mathcal{T}$ is in the class $\overline{H}(\nu, \tau)$ if it satisfies the following condition

$$\Re \left\{ \frac{\zeta g'(\zeta) + \nu \zeta^2 g''(\zeta)}{g(\zeta)} \right\} > \tau, \quad 0 \leq \nu, \tau < 1.$$

DEFINITION 2. [3] A function $g \in \mathcal{T}$ is in the class $\mathcal{J}^*(\nu, \tau)$ if it satisfies the following condition

$$\Re \left\{ \frac{(1-\nu)g(\zeta)}{\zeta} + \nu g'(\zeta) \right\} > \tau, \quad 0 \leq \nu, \tau < 1.$$

DEFINITION 3. [8] A function $g \in \mathcal{T}$ is in the class $\mathcal{R}^1(\nu, \tau)$ if it satisfies the following condition

$$\Re \{g'(\zeta) + \nu \zeta g''(\zeta)\} > \tau, \quad 0 \leq \nu, \tau < 1.$$

Received January 9, 2024. Revised January 18, 2024. Accepted February 10, 2024.

2010 Mathematics Subject Classification: 30C45, 30C50, 30C55.

Key words and phrases: Gaussian hypergeometric function, Analytic functions, Univalent functions, Starlike and convex functions.

© The Kangwon-Kyungki Mathematical Society, 2024.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

DEFINITION 4. A function $g \in \mathcal{T}$ is in the class $\mathcal{X}^*(\nu, \tau)$ if it satisfies the following condition

$$\Re \left\{ (1 - \nu) \frac{g(\zeta)}{\zeta g'(\zeta)} + \nu \left(1 + \frac{\zeta g''(\zeta)}{g'(\zeta)} \right) \right\} > \tau, \quad 0 \leq \nu, \tau < 1.$$

It is known that the Gaussian hypergeometric function ${}_2F_1(\gamma, \mu, \eta, \zeta)$ is one of the special functions and it is a solution of a second order linear ordinary differential equation

$$\zeta(1 - \zeta)y'' + [\eta - (\gamma + \mu + 1)\zeta]y' - \gamma\mu y = 0,$$

where γ, μ and η are constants. It is defined by

$${}_2F_1(\gamma, \mu, \eta, \zeta) = \sum_{t=0}^{\infty} \frac{(\gamma)_t (\mu)_t}{(\eta)_t t!} \zeta^t, \quad (1.3)$$

such that $\eta \neq 0, -1, -2, \dots$ and the Pochhammer symbol $(\alpha)_t$ defined by

$$(\alpha)_t = \begin{cases} \alpha(\alpha + 1) \dots (\alpha + t - 1) & \text{if } t = 1, 2, \dots \\ 1 & \text{if } t = 0. \end{cases} \quad (1.4)$$

If $\Re(\eta - \gamma - \mu) > 0$ and $\eta \neq 0, -1, -2, \dots$, then

$${}_2F_1(\gamma, \mu, \eta, 1) = \frac{\Gamma(\eta) \Gamma(\eta - \gamma - \mu)}{\Gamma(\eta - \gamma) \Gamma(\eta - \mu)}. \quad (1.5)$$

We note that

$$(\alpha)_t = \alpha(\alpha + 1)_{t-1}, \quad (1.6)$$

$$\begin{aligned} \zeta {}_2F_1(\gamma, \mu, \eta, \zeta) &= \zeta + \sum_{t=2}^{\infty} \frac{(\gamma)_{t-1} (\mu)_{t-1}}{(\eta)_{t-1} (t-1)!} \zeta^t \\ &= \zeta + \frac{\gamma\mu}{\eta} \sum_{t=2}^{\infty} \frac{(\gamma+1)_{t-2} (\mu+1)_{t-2}}{(\eta+1)_{t-2} (t-1)!} \zeta^t \\ &= \zeta - \left| \frac{\gamma\mu}{\eta} \right| \sum_{t=2}^{\infty} \frac{(\gamma+1)_{t-2} (\mu+1)_{t-2}}{(\eta+1)_{t-2} (t-1)!} \zeta^t. \end{aligned} \quad (1.7)$$

Now, let

$$\begin{aligned} F_1(\gamma, \mu, \eta, \zeta) &= \zeta ({}_2F_1(\gamma, \mu, \eta, \zeta)) \\ &= \zeta - \sum_{t=2}^{\infty} \frac{(\gamma)_{t-1} (\mu)_{t-1}}{(\eta)_{t-1} (t-1)!} \zeta^t. \end{aligned} \quad (1.8)$$

The hypergeometric function has many applications in mathematics, physics and engineering. In particular, in mathematics it has been used in different areas such as fractional calculus, geometric function theory, mathematical modeling and etc. The study of the hypergeometric function in the field of geometric function theory began to be in the researchers interest after the de Branges proof for the Bieberbach's conjecture in 1985 as he used the Gaussian hypergeometric function in his proof.

Necessary and sufficient conditions for $\zeta {}_2F_1(\gamma, \mu, \eta, \zeta)$ to be in various subclasses of analytic functions were investigated extensively by Silverman [12], Know and Cho [4], Mostafa [6] and Ramachandran et al. [10], and various geometric properties of this function were discussed in [1, 7, 9]. This paper will determine the necessary and

sufficient conditions for the Gaussian hypergeometric function to be in the classes $\overline{H}(\nu, \tau)$, $\mathcal{J}^*(\nu, \tau)$, $\mathcal{X}^*(\nu, \tau)$ and $\mathcal{R}^1(\nu, \tau)$.

2. Main Results

We need the following lemmas to prove our results.

LEMMA 1. [5] A function $g \in \mathcal{T}$ is in the class $\overline{H}(\nu, \tau)$ if and only if

$$\sum_{t=2}^{\infty} \{(t-1)(\nu t + 1) + (1-\tau)\} a_t \leq 1 - \tau, \quad 0 \leq \tau < 1.$$

LEMMA 2. [8] A function $g \in \mathcal{T}$ is in the class $\mathcal{R}^1(\nu, \tau)$ if and only if

$$\sum_{t=2}^{\infty} t[\nu(t-1) + 1] a_t \leq 1 - \tau, \quad 0 \leq \tau < 1.$$

Since $g \in \mathcal{R}^1(\nu, \tau) \Leftrightarrow \zeta g' \in \mathcal{J}^*(\nu, \tau)$, we have the following:

LEMMA 3. A function $g \in \mathcal{T}$ is in the class $\mathcal{J}^*(\nu, \tau)$ if and only if

$$\sum_{t=2}^{\infty} [\nu(t-1) + 1] a_t \leq 1 - \tau, \quad 0 \leq \tau < 1.$$

Using the same method and technique given by Silverman [11], we easily establish the following theorem, and so we omit the details.

THEOREM 1. Let $\frac{1}{3} \leq \nu < 1$. A function $g \in \mathcal{T}$ is in the class $\mathcal{X}^*(\nu, \tau)$ if and only if

$$\sum_{t=2}^{\infty} \{(t-1)[\nu(t+1) - 1] + (1-\tau)t\} a_t \leq 1 - \tau, \quad 0 \leq \tau < 1.$$

In this paper, we derive the necessary and sufficient conditions for the Gaussian hypergeometric function to be in $\overline{H}(\nu, \tau)$, $\mathcal{J}^*(\nu, \tau)$, $\mathcal{X}^*(\nu, \tau)$ and $\mathcal{R}^1(\nu, \tau)$.

THEOREM 2. If $\gamma, \mu > -1, \gamma\mu < 0$ and $\eta > \gamma + \mu + 2$, then ${}_2F_1(\gamma, \mu, \eta, \zeta) \in \overline{H}(\nu, \tau)$ if and only if

$$(2\nu + 1)\gamma\mu\eta - \gamma\mu(2\nu + 1)(\gamma + \mu + 2) + \nu(\gamma)_2(\mu)_2 + (1-\tau)(\eta - \gamma - \mu - 2)_2 \geq 0. \tag{2.1}$$

Proof. According to Lemma 1 and using (1.7), we need to show that

$$\sum_{t=2}^{\infty} \{(t-1)(\nu t + 1) + (1-\tau)\} \frac{(\gamma + 1)_{t-2}(\mu + 1)_{t-2}}{(\eta + 1)_{t-2}(t-1)!} \leq \left| \frac{\eta}{\gamma\mu} \right| (1-\tau).$$

Applying (1.5) and (1.6) , we get

$$\begin{aligned}
 & \sum_{t=2}^{\infty} \{(t-1)(\nu t+1) + (1-\tau)\} \frac{(\gamma+1)_{t-2}(\mu+1)_{t-2}}{(\eta+1)_{t-2}(t-1)!} \\
 = & \frac{\nu(\gamma+1)(\mu+1)}{(\eta+1)} \sum_{t=0}^{\infty} \frac{(\gamma+2)_t(\mu+2)_t}{(\eta+2)_t t!} + (2\nu+1) \sum_{t=0}^{\infty} \frac{(\gamma+1)_t(\mu+1)_t}{(\eta+1)_t t!} \\
 & + \frac{\eta(1-\tau)}{\gamma\mu} \sum_{t=1}^{\infty} \frac{(\gamma)_t(\mu)_t}{(\eta)_t t!} \\
 = & \frac{\nu(\gamma+1)(\mu+1)}{(\eta+1)} \frac{\Gamma(\eta+2)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} + (2\nu+1) \frac{\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-1)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} \\
 & + \frac{\eta(1-\tau)}{\gamma\mu} \left[\frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} - 1 \right] \\
 \leq & \left| \frac{\eta}{\gamma\mu} \right| (1-\tau),
 \end{aligned}$$

which is equivalent to

$$\begin{aligned}
 & \frac{\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} \left[\nu(\gamma+1)(\mu+1) + (2\nu+1)(\eta-\gamma-\mu-2) \right. \\
 & \quad \left. + \frac{(1-\tau)(\eta-\gamma-\mu-2)_2}{\gamma\mu} \right] \\
 \leq & (1-\tau) \left(\frac{\eta}{\gamma\mu} + \frac{\eta}{|\gamma\mu|} \right) = 0,
 \end{aligned}$$

which is valid if (2.1) holds. This completes the proof. □

THEOREM 3. *If $\gamma, \mu > 0$ and $\eta > \gamma + \mu + 2$, then $F_1(\gamma, \mu, \eta, \zeta) \in \overline{H}(\nu, \tau)$ if and only if*

$$\frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} \left[\frac{\nu(\gamma)_2(\mu)_2}{1-\tau} + \frac{\gamma\mu(2\nu+1)(\eta-\gamma-\mu-2)}{1-\tau} + (\eta-\gamma-\mu-2)_2 \right] \leq 2. \tag{2.2}$$

Proof. According to Lemma 1 and using (1.8), we need to show that

$$\sum_{t=2}^{\infty} \{(t-1)(\nu t+1) + (1-\tau)\} \frac{(\gamma)_{t-1}(\mu)_{t-1}}{(\eta)_{t-1}(t-1)!} \leq 1-\tau. \tag{2.3}$$

Using (1.5) and (1.6), (2.3) we get

$$\begin{aligned} & \sum_{t=2}^{\infty} \{(t-1)(\nu t+1) + (1-\tau)\} \frac{(\gamma)_{t-1}(\mu)_{t-1}}{(\eta)_{t-1}(t-1)!} \\ &= \frac{\nu\gamma\mu(\gamma+1)(\mu+1)}{\eta(\eta+1)} \sum_{t=0}^{\infty} \frac{(\gamma+2)_t(\mu+2)_t}{(\eta+2)_t t!} + \frac{(2\nu+1)\gamma\mu}{\eta} \sum_{t=0}^{\infty} \frac{(\gamma+1)_t(\mu+1)_t}{(\eta+1)_t t!} \\ & \quad + (1-\tau) \sum_{t=1}^{\infty} \frac{(\gamma)_t(\mu)_t}{(\eta)_t t!} \\ &= \frac{\nu\gamma\mu(\gamma+1)(\mu+1)}{\eta(\eta+1)} \frac{\Gamma(\eta+2)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} + \frac{(2\nu+1)\gamma\mu}{\eta} \frac{\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-1)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} \\ & \quad + (1-\tau) \left[\frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} - 1 \right] \\ & \leq 1-\tau, \end{aligned}$$

which is equivalent to

$$\begin{aligned} & \frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} [\nu(\gamma)_2(\mu)_2 + (2\nu+1)\gamma\mu(\eta-\gamma-\mu-2) + (1-\tau)(\eta-\gamma-\mu-2)_2] \\ & \leq 2(1-\tau). \end{aligned}$$

This completes the proof of Theorem 3. □

THEOREM 4. *If $\gamma, \mu > -1, \gamma\mu < 0$ and $\eta > \gamma + \mu + 1$, then ${}_2F_1(\gamma, \mu, \eta, \zeta) \in \mathcal{J}^*(\nu, \tau)$ if and only if*

$$\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-1)[\eta-\gamma-\mu-1+\nu\gamma\mu] - \tau\eta\Gamma(\eta-\gamma)\Gamma(\eta-\mu) \geq 0. \tag{2.4}$$

Proof. By using (1.7) and applying Lemma 3, it is adequate to show that

$$\sum_{t=2}^{\infty} [\nu(t-1) + 1] \frac{(\gamma+1)_{t-2}(\mu+1)_{t-2}}{(\eta+1)_{t-2}(t-1)!} \leq \left| \frac{\eta}{\gamma\mu} \right| (1-\tau). \tag{2.5}$$

It follows from (1.5), (1.6) and (2.5) that

$$\begin{aligned} & \sum_{t=2}^{\infty} [\nu(t-1) + 1] \frac{(\gamma+1)_{t-2}(\mu+1)_{t-2}}{(\eta+1)_{t-2}(t-1)!} \\ &= \nu \sum_{t=0}^{\infty} \frac{(\gamma+1)_t(\mu+1)_t}{(\eta+1)_t t!} + \frac{\eta}{\gamma\mu} \sum_{t=1}^{\infty} \frac{(\gamma)_t(\mu)_t}{(\eta)_t t!} \\ &= \nu \frac{\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-1)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} + \frac{\eta}{\gamma\mu} \left[\frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} - 1 \right] \\ & \leq \left| \frac{\eta}{\gamma\mu} \right| (1-\tau). \end{aligned}$$

Hence

$$\frac{\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-1)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} [\eta-\gamma-\mu-1+\nu\gamma\mu] \geq \tau\eta,$$

which is equivalent to (2.4). This completes the proof of Theorem 4. □

THEOREM 5. *If $\gamma, \mu > 0$ and $\eta > \gamma + \mu + 1$, then $F_1(\gamma, \mu, \eta, \zeta) \in \mathcal{J}^*(\nu, \tau)$ if and only if*

$$\frac{\Gamma(\eta)\Gamma(\eta - \gamma - \mu - 1)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} [\nu\gamma\mu + \eta - \gamma - \mu - 1] + \tau \leq 2. \tag{2.6}$$

Proof. According to Lemma 3 and using (1.8), we need to show that

$$\sum_{t=2}^{\infty} [\nu(t - 1) + 1] \frac{(\gamma)_{t-1}(\mu)_{t-1}}{(\eta)_{t-1}(t - 1)!} \leq 1 - \tau.$$

Thus,

$$\begin{aligned} & \sum_{t=2}^{\infty} [\nu(t - 1) + 1] \frac{(\gamma)_{t-1}(\mu)_{t-1}}{(\eta)_{t-1}(t - 1)!} \\ &= \frac{\nu\gamma\mu}{\eta} \sum_{t=0}^{\infty} \frac{(\gamma + 1)_t(\mu + 1)_t}{(\eta + 1)_t t!} + \sum_{t=1}^{\infty} \frac{(\gamma)_t(\mu)_t}{(\eta)_t t!} \\ &= \frac{\nu\gamma\mu}{\eta} \frac{\Gamma(\eta + 1)\Gamma(\eta - \gamma - \mu - 1)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} + \frac{\Gamma(\eta)\Gamma(\eta - \gamma - \mu)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} - 1 \\ &\leq 1 - \tau, \end{aligned}$$

if

$$\frac{\Gamma(\eta)\Gamma(\eta - \gamma - \mu - 1)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} [\nu\gamma\mu + \eta - \gamma - \mu - 1] \leq 2 - \tau,$$

which is equivalent to (2.6). This completes the proof of Theorem 5. □

THEOREM 6. *Let $\frac{1}{3} \leq \nu < 1$. If $\gamma, \mu > -1, \gamma\mu < 0$ and $\eta > \gamma + \mu + 2$, then ${}_2F_1(\gamma, \mu, \eta, \zeta) \in \mathcal{X}^*(\nu, \tau)$ if and only if*

$$\nu\gamma\mu(\gamma + 1)(\mu + 1) + \gamma\mu(3\nu - \tau)(\eta - \gamma - \mu - 2) + (1 - \tau)(\eta - \gamma - \mu - 2)_2 \geq 0. \tag{2.7}$$

Proof. According to Theorem 1 and using (1.7), we need to show that

$$\sum_{t=2}^{\infty} \left\{ (t - 1) [\nu(t + 1) - 1] + (1 - \tau)t \right\} \frac{(\gamma + 1)_{t-2}(\mu + 1)_{t-2}}{(\eta + 1)_{t-2}(t - 1)!} \leq \left| \frac{\eta}{\gamma\mu} \right| (1 - \tau).$$

Applying (1.5) and (1.6), we have

$$\begin{aligned}
 & \sum_{t=2}^{\infty} \{(t-1)[\nu(t+1)-1] + (1-\tau)t\} \frac{(\gamma+1)_{t-2}(\mu+1)_{t-2}}{(\eta+1)_{t-2}(t-1)!} \\
 = & \frac{\nu(\gamma+1)(\mu+1)}{(\eta+1)} \sum_{t=0}^{\infty} \frac{(\gamma+2)_t(\mu+2)_t}{(\eta+2)_t t!} + (3\nu-\tau) \sum_{t=0}^{\infty} \frac{(\gamma+1)_t(\mu+1)_t}{(\eta+1)_t t!} \\
 & + \frac{\eta(1-\tau)}{\gamma\mu} \sum_{t=1}^{\infty} \frac{(\gamma)_t(\mu)_t}{(\eta)_t t!} \\
 = & \frac{\nu(\gamma+1)(\mu+1)}{(\eta+1)} \frac{\Gamma(\eta+2)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} + (3\nu-\tau) \frac{\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-1)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} \\
 & + \frac{\eta(1-\tau)}{\gamma\mu} \left[\frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} - 1 \right] \\
 \leq & \left| \frac{\eta}{\gamma\mu} \right| (1-\tau),
 \end{aligned}$$

which is equivalent to

$$\begin{aligned}
 & \frac{\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} \left[\nu(\gamma+1)(\mu+1) + (3\nu-\tau)(\eta-\gamma-\mu-2) \right. \\
 & \qquad \qquad \qquad \left. + \frac{1-\tau}{\gamma\mu} (\eta-\gamma-\mu-2)_2 \right] \leq 0.
 \end{aligned}$$

This completes the proof of Theorem 6. □

THEOREM 7. *Let $\frac{1}{3} \leq \nu < 1$. If $\gamma, \mu > 0$ and $\eta > \gamma + \mu + 2$, then $F_1(\gamma, \mu, \eta, \zeta) \in \mathcal{X}^*(\nu, \tau)$ if and only if*

$$\begin{aligned}
 & \frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} \left[\frac{\nu(\gamma)_2(\mu)_2}{1-\tau} + \frac{\gamma\mu(3\nu-\tau)(\eta-\gamma-\mu-2)}{1-\tau} \right. \\
 & \qquad \qquad \qquad \left. + (\eta-\gamma-\mu-2)_2 \right] \leq 2. \tag{2.8}
 \end{aligned}$$

Proof. By using (1.8) and applying Theorem 1, we need to show that

$$\sum_{t=2}^{\infty} \{(t-1)[\nu(t+1)-1] + (1-\tau)t\} \frac{(\gamma)_{t-1}(\mu)_{t-1}}{(\eta)_{t-1}(t-1)!} \leq 1-\tau.$$

It follows from (1.5) and (1.6) that

$$\begin{aligned}
& \sum_{t=2}^{\infty} \{(t-1)[\nu(t+1)-1] + (1-\tau)t\} \frac{(\gamma)_{t-1}(\mu)_{t-1}}{(\eta)_{t-1}(t-1)!} \\
= & \frac{\nu\gamma\mu(\gamma+1)(\mu+1)}{\eta(\eta+1)} \sum_{t=0}^{\infty} \frac{(\gamma+2)_t(\mu+2)_t}{(\eta+2)_t t!} + \frac{(3\nu-\tau)\gamma\mu}{\eta} \sum_{t=0}^{\infty} \frac{(\gamma+1)_t(\mu+1)_t}{(\eta+1)_t t!} \\
& + (1-\tau) \sum_{t=1}^{\infty} \frac{(\gamma)_t(\mu)_t}{(\eta)_t t!} \\
= & \frac{\nu(\gamma)_2(\mu)_2}{(\eta)_2} \frac{\Gamma(\eta+2)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} + \frac{(3\nu-\tau)\gamma\mu}{\eta} \frac{\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-1)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} \\
& + (1-\tau) \left[\frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} - 1 \right] \\
\leq & 1-\tau,
\end{aligned}$$

which is equivalent to

$$\begin{aligned}
& \frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} [\nu(\gamma)_2(\mu)_2 + (3\nu-\tau)\gamma\mu(\eta-\gamma-\mu-2) + (1-\tau)(\eta-\gamma-\mu-2)_2] \\
& \leq 2(1-\tau).
\end{aligned}$$

This completes the proof of Theorem 7. \square

THEOREM 8. *If $\gamma, \mu > -1, \gamma\mu < 0$ and $\eta > \gamma + \mu + 2$, then $\zeta_2 F_1(\gamma, \mu, \eta, \zeta) \in \mathcal{R}^1(\nu, \tau)$ if and only if*

$$\begin{aligned}
& \Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-2) [\nu(\gamma)_2(\mu)_2 + (1+2\nu)(\eta-\gamma-\mu-2)\gamma\mu + (\eta-\gamma-\mu-2)_2] \\
& - \eta\tau\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-2) \geq 0.
\end{aligned} \tag{2.9}$$

Proof. According to Lemma 2 and using (1.7), we need to show that

$$\sum_{t=2}^{\infty} t[\nu(t-1)+1] \frac{(\gamma+1)_{t-2}(\mu+1)_{t-2}}{(\eta+1)_{t-2}(t-1)!} \leq \left| \frac{\eta}{\gamma\mu} \right| (1-\tau).$$

From (1.5) and (1.6) it follows that

$$\begin{aligned}
& \sum_{t=2}^{\infty} t[\nu(t-1)+1] \frac{(\gamma+1)_{t-2}(\mu+1)_{t-2}}{(\eta+1)_{t-2}(t-1)!} \\
= & \frac{\nu(\gamma+1)(\mu+1)}{(\eta+1)} \sum_{t=0}^{\infty} \frac{(\gamma+2)_t(\mu+2)_t}{(\eta+2)_t t!} + (1+2\nu) \sum_{t=0}^{\infty} \frac{(\gamma+1)_t(\mu+1)_t}{(\eta+1)_t t!} \\
& + \frac{\eta}{\gamma\mu} \sum_{t=1}^{\infty} \frac{(\gamma)_t(\mu)_t}{(\eta)_t t!} \\
= & \frac{\nu(\gamma+1)(\mu+1)}{(\eta+1)} \frac{\Gamma(\eta+2)\Gamma(\eta-\gamma-\mu-2)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} + (1+2\nu) \frac{\Gamma(\eta+1)\Gamma(\eta-\gamma-\mu-1)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} \\
& + \frac{\eta}{\gamma\mu} \left[\frac{\Gamma(\eta)\Gamma(\eta-\gamma-\mu)}{\Gamma(\eta-\gamma)\Gamma(\eta-\mu)} - 1 \right] \\
\leq & \left| \frac{\eta}{\gamma\mu} \right| (1-\tau),
\end{aligned}$$

which is equivalent to

$$\frac{\Gamma(\eta + 1)\Gamma(\eta - \gamma - \mu - 2)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} [\nu(\gamma)_2(\mu)_2 + (1 + 2\nu)(\eta - \gamma - \mu - 2)\gamma\mu + (\eta - \gamma - \mu - 2)_2] \geq \eta\tau.$$

This completes the proof of Theorem 8. □

THEOREM 9. *If $\gamma, \mu > 0$ and $\eta > \gamma + \mu + 2$, then $F_1(\gamma, \mu, \eta, \zeta) \in \mathcal{R}^1(\nu, \tau)$ if and only if*

$$\frac{\Gamma(\eta)\Gamma(\eta - \gamma - \mu - 2)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} [(1 + 2\nu)\gamma\mu(\eta - \gamma - \mu - 2) + \nu(\gamma)_2(\mu)_2 + (\eta - \gamma - \mu - 2)_2] + \tau \leq 2. \tag{2.10}$$

Proof. According to Lemma 2 and using (1.6) and (1.8), we need to show that

$$\sum_{t=2}^{\infty} t[\nu(t - 1) + 1] \frac{(\gamma)_{t-1}(\mu)_{t-1}}{(\eta)_{t-1}(t - 1)!} \leq 1 - \tau.$$

Thus,

$$\begin{aligned} & \sum_{t=2}^{\infty} t[\nu(t - 1) + 1] \frac{(\gamma)_{t-1}(\mu)_{t-1}}{(\eta)_{t-1}(t - 1)!} \\ &= \frac{\nu\gamma\mu(\gamma + 1)(\mu + 1)}{\eta(\eta + 1)} \sum_{t=0}^{\infty} \frac{(\gamma + 2)_t(\mu + 2)_t}{(\eta + 2)_t t!} + \frac{(1 + 2\nu)\gamma\mu}{\eta} \sum_{t=0}^{\infty} \frac{(\gamma + 1)_t(\mu + 1)_t}{(\eta + 1)_t t!} \\ & \quad + \sum_{t=1}^{\infty} \frac{(\gamma)_t(\mu)_t}{(\eta)_t t!} \\ &= \frac{\nu(\gamma)_2(\mu)_2}{(\eta)_2} \frac{\Gamma(\eta + 2)\Gamma(\eta - \gamma - \mu - 2)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} + \frac{(1 + 2\nu)\gamma\mu}{\eta} \frac{\Gamma(\eta + 1)\Gamma(\eta - \gamma - \mu - 1)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} \\ & \quad + \frac{\Gamma(\eta)\Gamma(\eta - \gamma - \mu)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} - 1 \\ &\leq 1 - \tau, \end{aligned}$$

if

$$\frac{\Gamma(\eta)\Gamma(\eta - \gamma - \mu - 2)}{\Gamma(\eta - \gamma)\Gamma(\eta - \mu)} [(1 + 2\nu)\gamma\mu(\eta - \gamma - \mu - 2) + \nu(\gamma)_2(\mu)_2 + (\eta - \gamma - \mu - 2)_2] \leq 2 - \tau,$$

which is equivalent to (2.10). This completes the proof of Theorem 9. □

3. Conclusion

Seeking for the necessary and sufficient conditions for the hypergeometric function to belong to different subclasses of analytic functions was investigated by many researchers intensively. In our paper, we investigated the necessary and sufficient conditions for the Gaussian hypergeometric function ${}_2F_1(\gamma, \mu, \eta, \zeta)$ and $F_1(\gamma, \mu, \eta, \zeta)$ to belong to the classes $\overline{H}(\nu, \tau)$, $\mathcal{J}^*(\nu, \tau)$, $\mathcal{X}^*(\nu, \tau)$ and $\mathcal{R}^1(\nu, \tau)$. For future work one can find the conditions for the Gaussian hypergeometric function to belong to different classes of analytic functions.

Acknowledgement

The authors would like to thank Prof. Abdel Moneim Y. Lashin for his support and valuable contribution in this research.

References

- [1] M. K. Aouf, A. O. Mostafa and H. M. Zayed, *Some properties of uniformly starlike and convex hypergeometric functions*, Azerbaijan J. Math. **9** (2) (2019), 3–19.
- [2] N. E. Cho, S. Y. Woo and S. Owa, *Uniform convexity properties for hypergeometric functions*, Fract. Calc. Appl. Anal. **5** (2002) (3), 303–314.
- [3] B. F. Hassan, *A0n application of certain integral and differential operators*, M. Sc. Thesis, Mansoura University, 2016.
- [4] O. S. Kwon and N. E. Cho, *Starlike and convex properties for hypergeometric functions*, Internat. J. Math. Math. Sci. **2008** (2008), Art. ID 159029, 11 Pages.
<https://doi.org/10.1155/2008/159029>
- [5] A. Y. Lashin, *On a certain subclass of starlike functions with negative coefficients*, J. Inequal. Pure Appl. Math. **10** (2009) (2), 1–8.
- [6] A. O. Mostafa, *Starlikeness and convexity results for hypergeometric functions*, Comput. Math. Appl. **59** (2010), 2821–2826.
<https://doi.org/10.1016/j.camwa.2010.01.053>
- [7] R. Nusrat, S. A. Eman and J. Abu, *On starlikeness and convexity properties of the generalized Bessel-hypergeometric function*, Indian J. Industrial Appl. Math. **10** (2019), 182–201.
<https://doi.org/10.5958/1945-919X.2019.00013.6>
- [8] H. Orhan, *On a subclass of analytic functions with negative coefficients*, Appl. Math. Comput. **142** (2003), 243–253.
[https://doi.org/10.1016/S0096-3003\(02\)00299-0](https://doi.org/10.1016/S0096-3003(02)00299-0)
- [9] G. I. Oros, *New conditions for univalence of confluent hypergeometric function*, Symmetry **13** (2021), 82.
<https://doi.org/10.3390/sym13010082>
- [10] C. Ramachandran, L. Vanitha and G. Murugusundaramoorthy, *Starlike and uniformly convex functions involving generalized hypergeometric function*, Internat. J. Pure Appl. Math. **92** (5) (2014), 691–701.
<http://dx.doi.org/10.12732/ijpam.v92i5.5>
- [11] H. Silverman, *Univalent functions with negative coefficients*, Proc. Amer. Math. Soc. **51** (1) (1975), 109–116.
- [12] H. Silverman, *Starlike and convexity properties for hypergeometric functions*, J. Math. Anal. Appl. **172** (1993) (2), 574–581.
<https://doi.org/10.1006/jmaa.1993.1044>

Abeer O. Badghaish

Department of Mathematics, Faculty of Science, King Abdulaziz, University,
Jeddah, Kingdom of Saudi Arabia.
E-mail: abadghaish@kau.edu.sa

Amani Z. Bajamal

Department of Mathematics, Faculty of Science, King Abdulaziz, University,
Jeddah, Kingdom of Saudi Arabia.
E-mail: azbajamal@kau.edu.sa