

ARTICLE

Study on the Digestion-Induced Changes in the Characteristics and Bioactivity of Korean Native and Overseas Cattle-Derived Peptides

 Jae Hyeon Kim¹, Da Young Lee¹, Seung Yun Lee², Ermie Jr. Mariano¹,
 Jae Won Jeong¹, Seung Hyeon Yun¹, Juhyun Lee¹, Jinmo Park¹, Yeongwoo Choi¹,
 Dahee Han¹, Jin Soo Kim¹, Cheorun Jo³, and Sun Jin Hur^{1,*}
¹Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea

²Division of Animal Science, Division of Applied Life Science (BK21 Four), Institute of Agriculture & Life Science, Gyeongsang National University, Jinju 52828, Korea

³Department of Agricultural Biotechnology, Center for Food and Bioconvergence, and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea

 OPEN ACCESS

Received July 14, 2023

Revised September 1, 2023

Accepted September 27, 2023

***Corresponding author** : Sun Jin Hur
 Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea
 Tel: +82-31-670-4673
 Fax: +82-31-670-3108
 E-mail: hursj@cau.ac.kr

***ORCID**

Jae Hyeon Kim
<https://orcid.org/0000-0003-1174-4737>
 Da Young Lee
<https://orcid.org/0000-0002-3172-0815>
 Seung Yun Lee
<https://orcid.org/0000-0002-8861-6517>
 Ermie Jr. Mariano
<https://orcid.org/0000-0003-2630-4603>
 Jae Won Jeong
<https://orcid.org/0000-0001-5240-1875>
 Seung Hyeon Yun
<https://orcid.org/0000-0002-9940-2960>
 Juhyun Lee
<https://orcid.org/0000-0001-6777-4447>
 Jinmo Park
<https://orcid.org/0009-0004-9626-1025>
 Yeongwoo Choi
<https://orcid.org/0009-0000-1882-4890>
 Dahee Han
<https://orcid.org/0009-0005-6423-3414>
 Jin Soo Kim
<https://orcid.org/0009-0007-7974-7885>
 Cheorun Jo
<https://orcid.org/0000-0003-2109-3798>
 Sun Jin Hur
<https://orcid.org/0000-0001-9386-5852>

Abstract This study was conducted to compare and analyze the changes in the biochemical characteristics and biological activity of peptide extracts derived from Chickso, Hanwoo, and Wagyu beef during digestion. The results of the *in vitro* digestion analysis revealed that the digestion rate, total free amino acid content, and antioxidant and antihypertensive activities of Chickso loin and shank myofibrillar proteins were significantly higher ($p < 0.05$) than those of Hanwoo and Wagyu loin and shank myofibrillar proteins. Particularly, the peptide extracts of Chickso loin and shank had a high angiotensin-converting enzyme inhibitory activity. In mice *in vivo* digestion experiment, the blood serum of mice fed with Chickso loin peptide extract (< 10 kDa) showed the highest antioxidant enzyme activity. Thus, Chickso peptide extracts were deemed to be similar or more bioactive than Hanwoo and Wagyu peptide extracts, and can be used as bioactive materials.

Keywords Korean native cattle, Chickso, peptide extracts, antioxidant activity, antihypertensive activity

Introduction

Currently, Korea is facing rapid changes in population structure and lifestyle, and the food industry is greatly affected by such changes in social structure (Cha and Rha, 2021). With the increase in per capita income, consumers are investing in more delicious and immune-boosting meals. In particular, the demand for meat, including beef, which is considered an immune-boosting food rich in protein, zinc, iron, and

vitamin D, is steadily increasing (Kim and Na, 2001; Whitnall and Pitts, 2019). Many studies on livestock have already been conducted in advanced countries to scientifically investigate the functionality of meats such as beef, chicken, and pork in response to consumers' demands; however, research on the same remains insufficient in Korea. In the future, various breeds and imported beef are expected to flow into Korea through various channels such as free trade agreements. Therefore, preparing the necessary domestic countermeasures accordingly is essential.

Hanwoo (*Bos taurus coreanae*) is the representative Korean native cattle. In general, Hanwoo is divided into five types: brown (Hanwoo, major), tiger color (Chickso, *Bos namadicus*), black (Heukwoo, *Bos namadicus Falconer*), Jeju black (Jeju Heukwoo, *Bos primigenius*), and white (Baekwoo; Jo et al., 2012; Kim et al., 2020; Utama et al., 2018).

Hanwoo gave many advantages to farming and transportation with the following strengths: not prone to disease, docile temperament, strong legs and hooves, withstanding rough breeding management, and good fertility (Lee et al., 2014). However, Hanwoo is currently being bred for meat and is raised in intensive farming to obtain beef. In addition, Korean meat quality grading system mainly evaluates marbling, and it can be said that Korean consumers' tendency to value the amount and taste of fat rather than lean meat has created today's Korean beef (Alam et al., 2013; Joo et al., 2017). Korean studies on the quality of Korean beef have revealed that Korean beef has superior muscle fat accumulation, unsaturated fatty acid composition, and sensorial properties compared to imported beef (Hwang, 2004; Park and Yoo, 1994). In addition, many studies have reported the excellent health-improving properties (antioxidant ability, anti-hypertension, anti-inflammatory, and immune-boosting effects) of Korean beef (Seol et al., 2018).

Chickso is one of the native breeds of Korean native cattle that has been bred for a long time besides with Hanwoo. It has dark brown and black vertical stripes on a yellowish-brown background, resembling a tiger pattern. Chickso has lower marbling than Hanwoo, but it has a higher oleic acid content than regular Hanwoo, which gives the meat a soft and unique taste (Lee and Joo, 2022; Utama et al., 2018). In particular, Chickso has a reddish color and strong meat flavor compared to Hanwoo, thus research and commercialization of aging, processing, cutting, and cooking methods are necessary. Furthermore, although a few comparative studies on the quality of foreign and native breeds have been conducted, no study has reported the functional properties of livestock products derived from traditional Korean cattle except for Hanwoo. Therefore, it is considered necessary to study the meat quality and functional characteristics of Chickso to establish it as a new breed different from Hanwoo and to lay the foundation for industrialization. Therefore, this study aimed to compare the characteristics of peptides derived from Chickso and Hanwoo, which are traditional domestic breeds, and Wagyu, which is a representative overseas breed.

Materials and Methods

Materials

α -Amylase from hog pancreas (CAS No. 9000-90-2), porcine bile extract (CAS No. 8008-63-7), lipase from porcine pancreas (CAS No. 9001-62-1), mucin from porcine stomach Type II (CAS No. 84082-64-4), pepsin from porcine gastric mucosa (CAS No. 9001-75-6), crystalline uric acid ($\geq 99\%$; CAS No. 69-93-2), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) diammonium salt (CAS No. 30931-67-0), 2,2-diphenyl-1-picrylhydrazyl (DPPH; CAS No. 1898-66-4), 3-(2-pyridyl)-5,6-diphenyl-1,2,4-triazine-p,p'-disulfonic acid monosodium salt hydrate (ferrozine), iron (II) chloride (CAS No. 7758-94-3), and iron (III) chloride (CAS No. 7705-08-0) were purchased from Sigma-Aldrich (St. Louis, MA, USA). EDTA-2Na dihydrate (CAS No. 6381-92-6), trichloroacetic acid (TCA; CAS No. 76-03-9), sodium phosphate dibasic anhydrous (CAS No. 7558-79-4), and sodium phosphate monobasic anhydrous (CAS No. 7558-80-7) were purchased from

Daejung Chemicals (Siheung, Korea). Potassium hexacyanoferrate (III) (CAS No. 13746-66-2) was purchased from Kanto Chemical (Tokyo, Japan) and 2,6-di-tert-butyl-4-methylphenol (99.0%) was purchased from Samchun Pure Chemical (Seoul, Korea).

Experimental animals

All animal experiments were approved by the Experimental Animal Ethics Committee of Chung-Ang University (Approval number: 2021-00038) and conducted in accordance with the rules of animal ethics. The institute of cancer research (ICR) mouse used in the experiment were eight-week-old females weighing 25–30 g purchased from Orient Bio. During the adaptation period, the mice were fed with water and pellet *ad libitum*. During the experiment, the mice were fed with defined amounts of mixed feed crumble and samples. DY-B-0061 bedding (shaving type) was purchased from Duyeol Biotech. The breeding environment was maintained at $22\pm 2^\circ\text{C}$ temperature, $60\pm 5\%$ relative humidity, and 12-hour light:dark cycle (07:00–19:00).

Acquisition of peptide extracts

Fig. 1 shows the process followed for preparing peptide extracts from the loin and shank of Chickso (*Bos namadicus*




Fig. 1. The process of preparing myofibrillar proteins and peptide extracts (<10 kDa) from Chickso, Hanwoo, and Wagyu beef. DW, distilled water; PBS, phosphate buffer.

Falconer), Hanwoo (*Bos taurus coreanae*), and Wagyu (*Bos taurus*) cattle. The Chickso loin and shank cuts used in the experiment were purchased from Woosung Ranch (1328, Chuseong-ro, Damyang-eup, Damyang-gun, Jeollanam-do) and Goseong Chickso (312-23, Sinan-ri, Ganseong-eup, Goseong-gun, Gangwon-do) in Korea. The Hanwoo loin and shank cuts were purchased from Famous Livestock (688-7, Nae-ri, Daedeok-myeon, Anseong-si, Gyeonggi-do) and Manse Livestock in Korea (15, Deokbongseowon-ro, Gongdo-eup, Anseong-si, Gyeonggi-do) in Korea. The Wagyu loin and shank cuts were purchased from Meatbox Global (22, Teheran-ro 34-gil, Gangnam-gu, Seoul) in Korea. Each sample (500 g) was completely ground using a stand mixer (SHMF-H3500TG, Hanil Electric, Bucheon, Korea). Subsequently, 10 L of distilled water (DW) was added and the samples were washed 20 times for about 30 min each time to remove as much blood and fat as much as possible. Next, the samples were homogenized using a total of 4 L of 0.04 M phosphate buffer (PBS; pH 7.4) and then centrifuged at $1,977\times g$ for 15 min at 4°C to obtain myofibrillar proteins. Thereafter, they were hydrolyzed with alkaline-AK enzyme (protein hydrolysate obtained by fermenting soybean meal with *Bacillus methylotrophicus*) at pH 11.0 for 2 h at 60°C . The protein hydrolysates were heated at 80°C for 15 min to deactivate the enzyme action and dried in a 55°C dry oven for 24 h. The dried protein hydrolysates were diluted in DW at a concentration of 25 mg/mL. Subsequently, peptides were fractionated to a size of 10 kDa or less through ultrafiltration using an Amicon filter tube (Amicon Ultra, Darmstadt, Germany) at $2,306\times g$ for 15 min and then freeze-dried at -70°C .

In vitro digestion experiment

The preparation conditions of synthetic digestive juices used in the human *in vitro* digestion model are described in Table 1 (Hur et al., 2011). Samples (120 mg) of loin and shank peptide extracts from Chickso, Hanwoo, and Wagyu were transferred into 50 mL tubes. Saliva (5 mL) was added and the mixture was shaken at 150 rpm for 5 min at 37°C in a shaking water bath to carry out the oral digestion step. Subsequently, 10 mL of gastric juice was added to the tube and reacted in the shaking water bath at 150 rpm for 2 h at 37°C to conduct the gastrointestinal digestion step. After the reaction, 10 mL of small intestinal juice and 5 mL of bile were added into the tube and reacted at 150 rpm for 2 h at 37°C to perform the small intestinal digestion step. After all the digestive steps were completed, 25 mL and 15 mL of DW were added to the tubes of the oral and gastrointestinal digestive steps, respectively. The final volume of the samples was 35 mL. In all *in vitro* digestion experiments, DW was added instead of digestive enzymes to obtain pre-digestion control samples under the same conditions. After the digestion experiment, all samples were centrifuged at $13,000\times g$ for 20 min at 4°C and then the supernatants were extracted and used as samples.

Table 1. Constituents and concentrations of synthetic digestive juices used in the human *in vitro* digestion model

Variable	Saliva (oral digestion step)	Gastric juice (gastrointestinal digestion step)	Duodenal juice (small intestinal digestion step)	Bile juice (small intestinal digestion step)
Inorganic and organic components	1.7 mL NaCl (175.3 g/L) 8 mL urea (25 g/L) 15 mg uric acid	6.5 mL HCl (37 g/L) 18 mL $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (22.2 g/L) 1 g bovine serum albumin	6.3 mL KCl (89.6 g/L) 9 mL $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (22.2 g/L) 1 g bovine serum albumin	69.3 mL NaHCO_3 (84.7 g/L) 10 mL $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (22.2 g/L) 1.8 g bovine serum albumin 30 g bile
Enzymes	290 mg α -amylase 25 mg mucin	2.5 g pepsin 3 g mucin	9 g pancreatin 1.5 g lipase	-
pH	6.8 \pm 0.2	1.50 \pm 0.02	8.0 \pm 0.2	7.0 \pm 0.2

Molecular weight distribution analysis

A gel permeation chromatography system was used to analyze the molecular weight distribution before and after *in vitro* digestion of Chickso, Hanwoo, and Wagyu loin and shank peptide extracts. Specifically, an ACQUITY APC System (Waters, Milford, MA, USA) with an Xbridge Protein BEH SEC column (Xbridge Protein, San Jose, CA, USA) was used for the analysis. All samples were filtered with a 0.2 μm syringe filter before use. Deionized water (A) and MeOH (B) were used as mobile phase solvents; the composition was 90% A and 10% B and the flow rate was 0.7 mL/min.

Analysis of changes in free amino acid composition

A Hitachi L-8900 amino acid analyzer (Hitachi High-Technologies, Tokyo, Japan) was used to analyze changes in the composition of the free amino acids before and after *in vitro* digestion of Chickso, Hanwoo, and Wagyu loin and shank peptide extracts. After diluting each sample to an appropriate concentration, 1 mL of 5% TCA solution was added to 1 mL of the sample, vortexed, and then centrifuged at $13,572\times g$ for 20 min at 20°C. Thereafter, 4 mL of n-hexane was added to 2 mL of the supernatant, agitated for 10 min, and then centrifuged at $1,977\times g$ for 20 min at 20°C. Subsequently, the lower layer solution was obtained and the centrifugation step was repeated twice to completely remove the n-hexane. Then, the samples were filtered using a 0.2 μm syringe filter and used for analysis.

In vitro physiological activity change analyses

Analysis of changes in the antioxidant activity

ABTS, DPPH, iron-chelating, and reducing power assays were used to analyze the antioxidant activity. ABTS radical scavenging activity was analyzed using Solution 1, prepared by mixing 0.038 mg of ABTS with 10 mL of DW according to the method described by Arts et al. (2004), and Solution 2, prepared by mixing 33 mg of potassium persulfate with 50 mL of DW. The ABTS solution was prepared by mixing 10 mL each of Solutions 1 and 2, covered with aluminum foil, and overnight incubated at room temperature to continue the activation of the solution. Twenty-microliter of samples were divided into a 96-well plate, followed by the addition of 180 μL working solution of ABTS reagent in MeOH. The well plate was wrapped in foil and the samples were allowed to react for 10 min. Then, the absorbance of the samples was measured at a wavelength of 734 nm using a microplate reader (Spectramax 190, Molecular Device, San Jose, CA, USA).

$$\text{ABTS scavenging activity (\%)} = \left(1 - \frac{\text{Absorbance of sample}}{\text{Absorbance of control}}\right) \times 100 \quad (1)$$

The DPPH radical scavenging activity was measured using the method described by Brand-Williams et al. (1995). DPPH (2 mg) and MeOH (25 mL) were mixed to prepare a DPPH solution, covered with aluminum foil, and refrigerated. The samples (0.1 mL) were added to a well plate and 0.1 mL of the DPPH solution was mixed with the samples. Subsequently, the well plate was wrapped in foil and reacted at room temperature for 30 min. Then, the absorbance of the samples was measured at a wavelength of 517 nm using a microplate reader (Spectramax 190, Molecular Device).

$$\text{DPPH scavenging activity (\%)} = \left(1 - \frac{\text{Absorbance of sample}}{\text{Absorbance of control}}\right) \times 100 \quad (2)$$

The iron-chelating assay was performed according to the method described by Decker and Welch (1990). EDTA-2Na

dihydrate was used as the standard solution, and DW the solvent in which the sample was dissolved was set as the control. After adding 200 μL of each prepared sample into individual 1.5 mL tubes, 200 μL of 2 mM iron (II) chloride and 40 μL of 5 mM ferrozine were added. After vortexing and reacting at room temperature for 10 min, the absorbance of the samples was measured at a wavelength of 562 nm using a microplate reader (Spectramax 190, Molecular Device).

$$\text{Iron chelating ability (\%)} = \left(1 - \frac{\text{Absorbance of sample}}{\text{Absorbance of control}}\right) \times 100 \quad (3)$$

The reducing power assay was conducted using the method described by Oyaizu (1986), with slight modifications. Briefly, 2.5 mL of 10% potassium ferricyanide and 2.5 mL of 0.2 M sodium PBS (pH 6.6) were added to 2.5 mL of the samples dissolved in ethanol and then reacted in an incubator at 50°C for 20 min. Subsequently, 2.5 mL of 10% TCA was added to terminate the reaction and the mixture was centrifuged at 2,841 \times g for 10 min. The supernatant was collected and mixed with a 0.1% ferric chloride solution in a ratio of 5:1 (v/v). Subsequently, the absorbance of the samples was measured at a wavelength of 700 nm to evaluate the reducing power.

Analysis of changes in the antihypertensive activity

To measure the antihypertensive activity using the angiotensin-converting enzyme (ACE) inhibitory assay, 50 mg/mL (w/v) of rabbit lung acetone powder (L0756, Sigma-Aldrich) was added to a 0.1 M sodium borate buffer (pH 8.3) containing 0.5 M sodium chloride. The mixture was stirred at 4°C for approximately 18–24 h for extraction. Subsequently, the supernatant was collected through centrifugation at 9,425 \times g for 30 min at 4°C to prepare the ACE solution. Next, a 0.1 M sodium borate buffer (pH 8.3) was prepared by appropriately mixing sodium tetraborate and boric acid to achieve a pH of 8.3. Hippuryl-His-Leu was added to the buffer to obtain a final concentration of 8.3 mM and the resulting solution was used as the ACE substrate.

Each sample (50 μL) was placed in a 2 mL tube, and a sample with DW only was used as the control. ACE substrate (50 μL) was added to each of the prepared samples and reacted for 10 min at 37°C. Thereafter, 50 μL of the ACE solution (25 mM/mL) was added and the mixture was incubated at 37°C for 30 min. Subsequently, 250 μL of 1 M HCl was added to a sample to stop the reaction and then 500 μL of ethyl acetate was added and the mixture was vortexed for 1 min. After centrifugation at 1,977 \times g for 10 min, 200 μL of the supernatant was taken and dried at 60°C for 30 min. Then, 1 mL of DW was added to it. The antihypertensive activity of the prepared sample was evaluated by measuring its absorbance at a wavelength of 228 nm using a UV/Vis Spectrophotometer (Agilent Technologies, Santa Clara, CA, USA).

$$\text{ACE inhibitory activity (\%)} = \left(\frac{\text{Absorbance of sample} - \text{Absorbance of sample}_{\text{HCl}}}{\text{Absorbance of control} - \text{Absorbance of control}_{\text{HCl}}}\right) \times 100 \quad (4)$$

Animal experiment

A total of 21 eight-week-old female ICR mice were randomly distributed into the control and treatment cages. The experiment was conducted for 21 d after a two-week adaptation period. The daily feed intake of general ICR mice was limited to 5 g, and the peptide extract was diluted in 100 μL DW to obtain a concentration of 800 mg/kg and then administered orally once a day using an oral injection needle.

After the peptide extract feeding experiment was completed, all experimental animals were sacrificed using CO₂ gas, and their heart blood was collected. Blood collected for antioxidant enzyme activity analysis was centrifuged at 2,000×g for 10 min at 4°C using a plasma separation gel tube (Gsmeditech, Wonju, Korea). The plasma obtained was used as a sample for the antioxidant activity analyses, and the serum obtained by centrifugation for 15 min using a serum separation gel tube (Gsmeditech) was used for ACE activity analysis.

Analysis of changes in antioxidant activity

Catalase activity of the samples were determined using an OxiTec Catalase Assay Kit (Biomax, Guri, Korea). Antioxidant activity was measured through three experiments using mice plasma samples. The prepared samples (25 µL) were added to a 96-well plate containing 25 µL of a 40 µM H₂O₂ solution. The resulting samples were reacted under dark conditions for 30 min at room temperature. After the reaction was completed, 50 µL of Oxi-Probe/HRP working solution was added to the wells and reacted at 20°C–25°C for 30 min. Subsequently, the absorbance of the samples was measured at 570 nm. The amount of enzyme that will decompose 1 µM of H₂O₂ per min at pH 7.0 at 25°C is equated to 1 unit (U) of catalase activity.

The peroxidase (POD) activity was determined by analyzing the content of H₂O₂ or peroxide present in the samples using an Oxi-Probe/HRP working solution and an OxiTec Hydrogen Peroxide/Peroxidase Assay kit (Biomax). The prepared samples (50 µL) were added to a 96-well plate with 50 µL of Oxi-Probe/H₂O₂ working solution and reacted under dark conditions for 30 min at 20°C–25°C. Then, the absorbance of the samples was measured at 560 nm using a microplate reader.

The superoxide dismutase (SOD) activity of the samples was determined by forming water-soluble tetrazolium salt (WST)-formazan complex through a redox reaction of xanthone oxidase and WST using a OxiTec SOD assay kit (Biomax). The prepared samples (20 µL) were added to a 96-well plate, and 200 µL of WST working solution was added to each well. Then, 20 µL of enzyme working solution was added to the well and mixed carefully. Thereafter, the plate was reacted at 37°C for 30 min, and the absorbance of the samples was measured at 450 nm using a microplate reader.

$$\text{SOD activity (Inhibition rate, \%)} = \left(1 - \frac{\text{Absorbance of sample} - \text{Absorbance of blank 2}}{\text{Absorbance of blank 1} - \text{Absorbance of blank 3}} \right) \quad (5)$$

Analysis of changes in antihypertensive activity

ACE activity was measured using an ACE activity assay kit (Elabscience, Houston, TX, USA). The absorbances of the sample and the blank were measured at 340 nm using a UV/Vis Spectrophotometer (Agilent Technologies). The ACE activity (U/L) was calculated using the following formula:

$$\text{ACE activity (U/L)} = \left(\frac{\text{Absorbance of sample} - \text{Absorbance of blank}}{\text{Reaction time}} \right) \times \frac{1,000}{\epsilon \times d} \times \frac{\text{Total volume}}{\text{Volume of sample}} \times \text{DF} \quad (6)$$

where, ϵ is the extinction coefficient (0.8 L/mmol/cm) of a wavelength of 340 nm at an optical distance of 1 cm, d is the optical distance of a quartz cuvette (0.5 cm), DF is the dilution factor, and the reaction time is 5 min.

Statistical analysis

For the analysis of the experimental results, a statistical analysis was conducted after obtaining the mean and SD using SPSS Statics 26 (IBM, Armonk, NY, USA). The Student's t-test was used for analyzing the difference between two variables and the

one-way analysis of variance method was used for analyzing the differences among three or more variables. The Student–Newman–Keuls post-test was conducted at a significance level of $p < 0.05$ for each average difference.

Results and Discussion

Molecular weight distribution analysis

Tables 2 and 3 present the molecular weight distributions of the peptide extracts derived from Chickso, Hanwoo, and Wagyu beef before and after *in vitro* digestion. The degree of decomposition of the peptide due to digestion could be

Table 2. The molecular weight distribution of peptide extracts (<10 kDa) from Chickso, Hanwoo, and Wagyu loin

Contents	Chickso		Hanwoo		Wagyu	
	Before digestion	After digestion	Before digestion	After digestion	Before digestion	After digestion
Mn (Da)	302±11.78 ^B	267±6.56 ^B	270±19.47 ^B	282±12.17 ^B	466±15.54 ^A	297±8.96 ^B
Mw (Da)	666±9.16 ^B	312±64.51 ^D	338±30.81 ^D	355±22.38 ^D	2,007±115.52 ^A	479±21.75 ^C
Mz (Da)	1,595±61.57 ^B	373±39.15 ^D	446±35.68 ^D	438±29.52 ^D	5,037±175.28 ^A	1,138±34.48 ^C
Polydispersity	2.20±0.08 ^B	1.17±0.12 ^C	1.25±0.06 ^C	1.26±0.11 ^C	4.30±0.35 ^A	1.61±0.15 ^C
>10,000 Da (%)	-	-	-	-	-	-
5,000–10,000 Da (%)	0.07±0.01	-	-	-	0.001±0.000	-
3,000–5,000 Da (%)	2.65±0.28	-	-	-	1.19±0.08	-
1,000–3,000 Da (%)	15.72±0.77	0.42±0.05 ^D	1.49±0.07	0.56±0.02	8.41±0.35	1.10±0.07
200–1,000 Da (%)	58.33±0.59	83.46±2.14 ^A	79.27±2.29	79.18±4.45	76.80±3.78	84.75±2.14
<200 Da (%)	23.23±0.41	16.12±0.63	19.25±0.78	20.27±0.71	13.60±0.12	14.15±1.10

Each value is a mean±SD of three replicates.

^{A–D} Different uppercase letters in the same line indicate significant differences among Chickso, Hanwoo, and Wagyu loin peptides ($p < 0.05$).

Mn, number-average molecular weight; Mw, weight-average molecular weight; Mz, Z-average molecular weight.

Table 3. The molecular weight distribution of peptide extracts (<10 kDa) from Chickso, Hanwoo, and Wagyu shank

Contents	Chickso		Hanwoo		Wagyu	
	Before digestion	After digestion	Before digestion	After digestion	Before digestion	After digestion
Mn (Da)	508±15.54 ^A	285±6.19 ^B	324±5.84 ^B	296±4.42 ^B	534±10.11 ^A	280±7.38 ^B
Mw (Da)	1,954±59.62 ^A	395±9.27 ^B	527±9.69 ^B	435±8.53 ^B	2,136±76.19 ^A	325±9.32 ^B
Mz (Da)	4,821±73.83 ^A	730±11.25 ^B	901±11.38 ^B	799±10.52 ^B	5,268±112.38 ^A	381±15.25 ^C
Polydispersity	3.85±0.12 ^A	1.39±0.08 ^B	1.63±0.11 ^B	1.47±0.11 ^B	4.00±0.20 ^A	1.16±0.09
>10,000 Da (%)	1.63±0.05	-	-	-	2.05±0.04	-
5,000–10,000 Da (%)	7.86±0.78	-	-	-	10.28±0.15	-
3,000–5,000 Da (%)	12.74±0.88	0.18±0.01	0.11±0.01	0.15±0.01	12.36±0.38	-
1,000–3,000 Da (%)	26.65±1.25	5.36±0.08	12.98±0.35	7.74±0.86	24.48±2.54	-
200–1,000 Da (%)	39.83±3.12	80.43±1.49	69.99±2.89	77.30±3.98	41.35±3.15	85.70±1.17
<200 Da (%)	11.30±0.52	14.03±0.28	16.93±1.12	14.81±0.95	9.49±0.49	14.30±0.28

Each value is a mean±SD of three replicates.

^{A–C} Different uppercase letters in the same line indicate significant differences among Chickso, Hanwoo, and Wagyu shank peptides ($p < 0.05$).

Mn, number-average molecular weight; Mw, weight-average molecular weight; Mz, Z-average molecular weight.

predicted based on the change in molecular weight of the peptide extract. Irrespective of the breeds and cuts, there was an increase in the number of low-molecular-weight peptides after digestion. In the pre-digestion samples, peptides with a molecular weight of 3,000–10,000 Da were present. However, most of them were digested into low-molecular-weight peptides of 1,000 Da or less after digestion. The proportion of 200–1,000 Da peptides among the loin and shank peptide extracts of Chickso before *in vitro* digestion was approximately 58% and 39%, respectively, but after *in vitro* digestion, it was approximately 83% and 80%, respectively, indicating a significant increase in the proportion of low-molecular-weight peptides after digestion. These results indicate that the peptide bonds were cleaved by hydrolysis during *in vitro* digestion.

According to previous studies, the biological activity of peptides depends on the molecular structure and physicochemical properties of peptides, and small peptides have higher bioavailability and bioactivity (Udenigwe and Aluko, 2012). Therefore, the peptide extracts derived from the loin and shank, which are high in low-molecular-weight peptides of less than 3,000 Da, are expected to have efficient antioxidant and antihypertensive activities.

Analysis of changes in free amino acid composition

Tables 4 and 5 present the free amino acid compositions of the peptide extracts before and after *in vitro* digestion. The total

Table 4. Changes in the amino acid compositions of Chickso, Hanwoo, and Wagyu loin peptides due to *in vitro* digestion

Amino acids (mg/100 g)	Chickso		Hanwoo		Wagyu	
	Before digestion	After digestion	Before digestion	After digestion	Before digestion	After digestion
Aspartic acid	217.2±12.5 ^B	236.7±15.1 ^A	205.3±11.2 ^B	232.8±19.8 ^A	187.5±15.5 ^C	166.7±11.5 ^D
Threonine	100.0±3.8 ^B	416.2±21.9 ^A	103.7±5.9 ^B	408.8±28.7 ^A	109.8±9.6 ^B	467.1±15.3 ^A
Serine	88.6±2.4 ^C	280.0±15.8 ^A	90.8±3.8 ^C	275.9±12.7 ^A	80.1±4.2 ^C	151.4±11.5 ^B
Asparagine	74.7±2.3 ^C	290.8±13.7 ^A	89.5±4.1 ^C	287.9±22.3 ^A	65.3±5.8 ^C	172.3±12.7 ^B
Glutamic acid	947.1±52.1 ^C	1,133.3±95.3 ^A	940.2±22.7 ^C	1,113.7±75.1 ^A	828.4±41.7 ^D	1,024.1±83.4 ^B
Glutamine	10.8±0.7 ^C	450.9±41.1 ^B	12.4±0.9 ^C	437.4±33.5 ^B	9.9±0.3 ^C	689.3±22.3 ^A
Glycine	39.4±1.5 ^C	326.9±15.8 ^A	38.6±0.8 ^C	326.2±18.9 ^A	31.5±2.1 ^C	201.6±17.2 ^B
Alanine	164.7±12.2 ^B	529.7±29.2 ^A	167.7±11.5 ^B	521.5±37.8 ^A	146.5±10.9 ^B	498.9±18.9 ^A
Valine	176.1±13.6 ^C	495.2±22.5 ^A	177.0±14.9 ^C	488.3±16.5 ^A	169.7±13.2 ^C	392.2±26.3 ^B
Cystine	80.8±5.4 ^B	140.4±13.1 ^A	82.6±4.3 ^B	145.4±11.2 ^A	51.0±2.3 ^B	136.8±11.2 ^A
Methionine	259.5±12.9 ^B	805.8±42.3 ^A	260.1±15.7 ^B	790.1±38.9 ^A	211.2±15.8 ^B	673.1±45.7 ^A
Isoleucine	291.3±15.8 ^C	690.0±44.8 ^A	288.4±24.4 ^C	683.6±101.2 ^A	278.3±21.2 ^C	549.6±15.4 ^B
Leucine	1,067.9±98.5 ^C	3,279.8±239.8 ^A	1,067.0±82.3 ^C	3,212.2±284.5 ^A	854.8±33.6 ^C	2,366.4±115.7 ^B
Tyrosine	158.7±10.6 ^C	1,754.5±112.7 ^B	138.4±8.8 ^C	1,717.1±211.7 ^B	164.2±17.7 ^C	2,041.8±132.4 ^A
Phenylalanine	182.5±11.2 ^C	1,815.2±89.5 ^A	189.2±13.1 ^C	1,781.7±189.4 ^A	178.0±11.5 ^C	1,551.2±109.8 ^B
Tryptophan	89.5±8.3 ^C	293.3±19.8 ^A	81.4±3.8 ^C	274.2±42.1 ^A	151.3±8.0 ^B	292.8±12.1 ^A
Proline	44.1±6.9 ^B	93.0±5.3 ^A	37.0±2.6 ^B	91.5±15.3 ^A	24.4±3.2 ^C	95.7±3.8 ^A
Lysine	394.1±27.3 ^C	2,651.1±127.7 ^A	391.5±24.3 ^C	2,591.4±306.2 ^A	498.7±11.7 ^B	2,606.6±201.7 ^A
Histidine	28.3±2.2 ^C	278.3±12.6 ^B	28.3±1.6 ^C	278.4±17.6 ^B	33.8±1.9 ^C	356.3±16.9 ^A
Arginine	291.2±11.8 ^C	2,886.6±227.9 ^A	292.8±17.5 ^C	2,838.4±165.9 ^A	250.7±12.1 ^C	1,560.8±137.0 ^B

Each value is a mean±SD of three replicates.

^{A-D} Different uppercase letters in the same line indicate significant differences among Chickso, Hanwoo, and Wagyu loin peptides ($p < 0.05$).

Table 5. Changes in the amino acid compositions of Chickso, Hanwoo, and Wagyu shank peptides due to *in vitro* digestion

Amino acids (mg/100 g)	Chickso		Hanwoo		Wagyu	
	Before digestion	After digestion	Before digestion	After digestion	Before digestion	After digestion
Aspartic acid	126.0±9.8 ^C	319.8±26.8 ^B	94.9±7.8 ^C	336.3±18.5 ^B	131.2±10.2 ^C	415.8±11.7 ^A
Threonine	478.7±13.7 ^B	776.7±24.1 ^A	497.1±14.0 ^B	884.6±27.4 ^A	383.3±23.7 ^B	764.0±27.3 ^A
Serine	173.5±11.2 ^B	407.9±30.0 ^A	95.7±5.9 ^B	355.1±21.1 ^A	124.5±5.5 ^B	419.2±13.8 ^A
Asparagine	20.0±1.8 ^D	190.3±10.9 ^A	15.3±0.4 ^D	58.4±0.9 ^C	18.9±1.2 ^D	135.5±4.9 ^B
Glutamic acid	322.3±21.5 ^C	712.3±41.4 ^B	316.9±14.5 ^C	1,001.3±20.4 ^A	414.8±20.4 ^C	1,174.2±75.5 ^A
Glutamine	0.0±0.0	367.5±22.1 ^A	0.0±0.0	72.5±3.6 ^C	0.0±0.0	210.3±13.5 ^B
Glycine	109.0±8.4 ^B	412.6±20.8 ^A	157.9±9.6 ^B	437.5±14.7 ^A	96.6±3.9 ^B	409.1±17.9 ^A
Alanine	443.2±13.7 ^C	855.60±44.9 ^A	267.3±11.5 ^D	603.7±16.9 ^B	359.2±14.8 ^C	832.8±36.1 ^A
Valine	771.1±62.2 ^B	1,165.4±107.4 ^A	739.9±43.3 ^B	1,160.2±25.0 ^A	564.6±22.5 ^C	1,059.7±55.4 ^A
Cystine	5.4±0.4 ^B	34.8±0.9 ^A	7.4±0.4 ^B	36.1±0.8 ^A	5.2±0.3 ^B	40.4±1.5 ^A
Methionine	947.0±31.6 ^B	1,227.0±41.7 ^A	990.8±63.7 ^B	1,220.7±62.5 ^A	763.0±33.4 ^C	1,207.7±62.2 ^A
Isoleucine	561.6±91.7 ^B	1,070.2±32.5 ^A	631.9±39.9 ^B	1,174.2±43.7 ^A	447.9±31.5 ^B	1,047.6±44.3 ^A
Leucine	2,008.5±118.4 ^C	3,596.9±270.3 ^A	1,687.0±182.1 ^D	3,185.9±243.8 ^B	1,343.8±116.9 ^D	3,463.2±311.5 ^A
Tyrosine	665.5±24.8	1,892.0±98.9	861.9±47.2	2,031.3±78.8	512.2±23.9	2,073.5±101.6
Phenylalanine	629.9±41.9	1,933.2±121.6	788.0±29.6	1,922.7±62.6	455.5±17.4	1,999.5±74.8
Tryptophan	130.5±10.3	393.3±21.5	111.4±6.3	374.2±10.4	53.2±2.6	385.4±11.2
Proline	62.8±3.8	107.0±4.9	44.1±4.0	94.1±4.9	41.7±2.3	108.6±2.9
Lysine	1,153.1±78.8	1,432.7±15.7	1,374.7±68.9	2,250.8±66.9	704.9±33.4	1,749.1±46.8
Histidine	211.0±15.1	420.1±26.5	147.7±9.5	310.3±15.4	86.0±0.7	348.1±16.4
Arginine	188.9±12.4	1,574.1±104.2	171.1±4.3	1,665.7±51.2	190.0±11.2	1,654.5±44.7

Each value is a mean±SD of three replicates.

^{A-D} Different uppercase letters in the same line indicate significant differences among Chickso, Hanwoo, and Wagyu shank peptides ($p < 0.05$).

free amino acid contents detected after the *in vitro* digestion of Chickso, Hanwoo, and Wagyu shank were 9,315.42 mg/100 g, 9,153.71 mg/100 g, and 6,864.45 mg/100 g, respectively, which were higher than those detected for the loin peptide extracts of each breed. Regardless of the cuts, the main free amino acids of the peptide extracts before *in vitro* digestion were glutamic acid, leucine, lysine, isoleucine, and arginine, and the main free amino acids detected while leucine, tyrosine, lysine, arginine, and phenylalanine were dominant after *in vitro* digestion.

Considering that the content of cysteine and methionine sulfur-containing amino acids known to have antioxidant activity (Atmaca, 2004) increased significantly compared to that before digestion, the peptide extracts can be expected to have higher antioxidant activities after the digestion process. According to Houglund et al. (2013), histidine, tryptophan, and tyrosine are also representative reactive amino acids that can exhibit effective antioxidant activity by reacting with active oxygen species. In addition, peptides with hydrophobic amino acids, such as valine and isoleucine, and aromatic amino acids, such as phenylalanine and tryptophan, contribute to inhibiting ACE activity (Aluko, 2015). Thus, the peptide extracts can also be expected to have higher antihypertensive activities after the digestion process.

Changes in physiological activity based on the *in vitro* digestion experiment

Changes in antioxidant activity

The results of the antioxidant analysis based on the *in vitro* digestion of the loin and shank-derived peptide extracts of Chickso, Hanwoo, and Wagyu beef are shown in Figs. 2, 3, 4, and 5. Approximately 19% of ABTS radical scavenging ability (Fig. 2), 76% of DPPH radical scavenging ability (Fig. 3), 13% of iron chelating ability (Fig. 4), and 6% or more of reducing power (Fig. 5) increased (detail data are not shown). Comparison of the ABTS radical elimination ability of the peptide extracts before digestion showed that the loin had significantly higher antioxidant activity than the shank, and Chickso (75.01%) and Hanwoo (76.06%) loins had higher antioxidant activity than Wagyu loin (71.13%). Furthermore, the DPPH




Fig. 2. Changes in the 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) radical scavenging activity of Chickso, Hanwoo, and Wagyu loin and shank peptide extracts due to *in vitro* digestion. A comparison of the ABTS radical scavenging activity of the peptides before and after digestion. Data are presented as mean \pm SD. * $p < 0.05$ and ** $p < 0.001$ indicate a significant difference according to digestion.




Fig. 3. Changes in the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity of Chickso, Hanwoo, and Wagyu loin and shank peptide extracts due to *in vitro* digestion. A comparison of the DPPH radical scavenging activity of the peptides before and after digestion. Data are presented as mean \pm SD. ** $p < 0.001$ indicates a significant difference according to digestion.




Fig. 4. Changes in the iron chelating activity of Chickso, Hanwoo, and Wagyu loin and shank peptide extracts due to *in vitro* digestion. A comparison of the iron chelating activity of the peptides before and after digestion. * $p < 0.05$ indicates a significant difference according to digestion.




Fig. 5. Changes in the reducing power of Chickso, Hanwoo, and Wagyu loin and shank peptide extracts due to *in vitro* digestion. A comparison of the reducing power of the peptides before and after digestion. * $p < 0.05$ indicates a significant difference according to digestion.

radical scavenging ability of Chickso loin (36.79%) before digestion was approximately 2.9 times higher than that of other breeds, and Chickso loin (46.34%) and shank (43.59%) had the highest DPPH radical scavenging abilities among the digested samples. There was no significant difference in the iron chelating ability between breeds and cuts, and Wagyu samples had the highest iron chelating ability among all pre- and post-digestion samples. Furthermore, Chickso shank (1.04) had the highest reducing power among all pre-digestion samples, and Hanwoo loin (1.07) had the highest reducing power among all post-digestion samples. Among the post-digestion samples, the reducing power of the Hanwoo samples was significantly higher than that of the Chickso and Wagyu samples. A comparison of the molecular weight distribution analysis presented in Tables 2 and 3 revealed that proteins have already been digested enough during peptide extraction to fully express their physiological activity, making it difficult to increase the antioxidant effect through additional decomposition during the

digestion process. Chen et al. (1998) stated that low-molecular-weight peptides of 1 kDa or less have antioxidant activity. Similarly, Olagunju et al. (2018) reported that peptides with a molecular weight of 1–3 kDa have a higher antioxidant activity than peptides with other molecular weights. Lee and Hur (2017) also revealed that protein hydrolysates derived from beef source fiber proteins with a molecular weight of 3 kDa or less exhibit strong free radical scavenging ability. The differences in antioxidant activity between breeds and cuts may have been affected by the degree of hydrolysis according to the characteristics of the proteases, the size of peptide molecules, hydrophobicity, amino acid composition, and structural properties of the peptides produced (Pihlanto, 2006).

The current study revealed that the antioxidant activity increased slightly when the peptide extracts were hydrolyzed using the *in vitro* digestion model, and overall, the antioxidant activity of the Chickso and Hanwoo samples was higher than that of the Wagyu samples. These results are likely due to the high content of low-molecular-weight peptides such as glutathione (a crystalline peptide composed of glutamic acid, cysteine, and glycine), which are known to have excellent antioxidant effects (Pompella et al., 2003).

Changes in antihypertensive activity

Fig. 6 shows the antihypertensive activities of Hanwoo, Chickso, and Wagyu peptide extracts based on the ACE inhibitor assay. All pre-digestion samples showed a high ACE inhibitory ability of more than 70%, and the peptide extracts of Chickso loin and shank had the highest antihypertensive activities. However, after the digestion process, the ACE inhibition ability in all samples decreased sharply to less than 30%. This was probably because of the enzymatic decomposition of peptides with antihypertensive activity or the loss of activity due to the pH and temperature conditions during the *in vitro* digestion process. The results of this experiment were similar to those of a previous study by Jang and Lee (2005), who reported that excessive hydrolysis lowers the antihypertensive activity of peptides. Because all samples showed high activity before digestion, it was decided that additional experiments using animal models were needed to determine whether the antihypertensive activity was retained during *in vivo* digestion.




Fig. 6. Changes in the angiotensin-converting enzyme (ACE) inhibitory activity of Chickso, Hanwoo, and Wagyu loin and shank peptide extracts due to *in vitro* digestion. A comparison of the ACE inhibitory activity of the peptides before and after digestion. Data are presented as mean±SD. ** $p < 0.001$ indicates a significant difference according to digestion.

Changes in physiological activity based on the animal experiments

Changes in antioxidant activity

Fig. 7 shows the antioxidant activities observed in mice fed with the peptide extracts derived from the loin and shank of Chickso, Hanwoo, and Wagyu beef. All treatment groups showed catalase activity of more than 340 mU/mL, which was higher than that of the control, and the highest activity was observed in the treatment groups fed with peptide extracts derived from Chickso loin. The POD activity of all treatment groups, especially the groups fed with Chickso and Wagyu loin peptide extracts, was higher than that of the control by 0.38 mU/mL or more. The SOD activity of all treatment groups was similar to or higher than that of the control. In addition, the production rate of methionine, a sulfur-containing amino acid known to have antioxidant activity in proteins, and histidine and tyrosine, reactive amino acids, in Chickso and Wagyu loin/shank peptides were higher than that in the Hanwoo loin/shank peptide samples after digestion (Tables 4 and 5). In general, histidine-containing peptides exhibit antioxidant activity through a single electron transfer mechanism and histidine-containing peptides through a hydrogen atom transfer mechanism (Esfandi et al., 2019).

According to previous studies, peptide extracts derived from milk, fish, chicken breasts, and various animal materials are known to have excellent antioxidant activity in the body (Liu et al., 2016; Power et al., 2013; Sakanaka et al., 2005; Sun et




Fig. 7. The antioxidant enzyme activities of mice plasma which fed with the peptide extracts derived from the loin and shank of Chickso, Hanwoo, and Wagyu. (A) Catalase activity, (B) peroxidase activity, (C) superoxide dismutase (SOD) activity. Data are presented as mean \pm SD. ^{A-F} Different uppercase letters indicate a significant difference based on breeds and cuts ($p < 0.05$). CTL, Control.

al., 2012). In addition, the antioxidant activity of shank was higher than that of loin from Hanwoo (Oh et al., 2016). The study of Kim et al. (2018) demonstrated that the antioxidant octapeptide (Cys-Cys-Cys-Cys-Ser-Val-Gln-Lys; 888.60 Da) isolated and identified from Korean beef extract has high oxygen absorbance capacity. Also, free amino acids detected through the digestion process are known to have a positive effect on enhancing the antioxidant activity of enzymes. Most glutathione is synthesized between cells, and the reduced form of glutathione protects cells by reducing active oxygen (Dringen and Hamprecht, 1999; Gaucher et al., 2018; Levy et al., 1993; Piste, 2013). Lysine can increase the ability to free radical scavenging to protect against oxidative damage by upregulating the expression of antioxidant enzyme genes (Li, 2016). Methionine can act as an endogenous antioxidant, and is an efficient remover of almost all oxidizing molecules under physiological conditions such as H₂O₂ and hydroxyl radicals (Atmaca, 2004; Levine et al., 1996). The results of the present study also confirm that the consumption of beef-derived peptide extracts can have antioxidant effects on the body. Thus, beef-derived peptide extracts can be used as a pharmaceutical product to prevent cell oxidation.

Analysis of changes in antihypertensive activity

Fig. 8 shows the antihypertensive activities in mice fed with the peptide extracts derived from the loin and shank of Chickso, Hanwoo, and Wagyu beef. Compared to the control group, which was only fed with general feed, there was no significant activity in any of the treatment groups. These results could be similar to the significant reduction in the ACE inhibitory ability of peptide extracts after digestion to less than the before digestion activity in *in vitro* digestion experiment (Fig. 6). It was judged that the peptide extracts with antihypertensive activity were digested into smaller peptides due to the mouse intestinal digestive enzymes and microorganisms, resulting in loss of antihypertensive activity (Escudero et al., 2014). The food-derived peptides produced after digestion have been shown to have an antihypertensive effect; however, this effect is reduced when peptides of already active size are consumed. In fact, peptides are known to have low bioavailability because of their weak intestinal mucosal permeability when administered orally (Hamman et al., 2005). Accordingly, the amount of bioactive peptides absorbed after digestion may not have been enough to increase the ACE inhibitory ability detected in the mouse serum, resulting in the non-significant difference between the control and the treatments (Mora et al., 2018).




Fig. 8. The angiotensin-converting enzyme (ACE) activity of Chickso, Hanwoo, and Wagyu loin and shank peptide extracts (<10 kDa) after *in vivo* digestion in mice. Data are presented as mean±SD. CTL, Control.

Therefore, additional research is needed to prevent peptides from losing their activity in the body. Furthermore, a previous study showed that peptides must be absorbed in the intestine in an undamaged form to reach the target site even after oral administration (Gianfranceschi et al., 2018). Therefore, research on small-sized peptides (formed from 2–5 amino acids) that can directly pass through the intestinal membrane without digestion and reach the target site (Vermeirssen et al., 2002) is needed.

Conclusion

This study was conducted to analyze the changes in the physicochemical properties and physiological activities of peptide extracts derived from the loin and shank of Chickso, Hanwoo, and Wagyu beef after *in vitro* and *in vivo* (in mice) digestion. Specifically, the changes in the molecular weight, amino acid composition, and antioxidant and antihypertensive abilities of the peptides due to the digestion process were analyzed.

The results revealed that there was no significant difference in the digestion rates of Chickso loin and shank peptide extracts during the *in vitro* digestion process, but the total free amino acid content and antioxidant activity of the Chickso and Hanwoo samples were higher than those of the Wagyu samples. Chickso loin and shank peptide extracts showed a high ACE inhibitory activity. Furthermore, *in vivo* experiments in mice showed high antioxidant activity in the treatment group fed with Chickso-derived peptide extracts.

Therefore, the peptide extract of Chickso, one of the traditional cattle breeds in Korea, is considered to have better bioactivity than the peptide extracts of Hanwoo and Wagyu. In the future, if research is conducted to maximize the activity of such peptide extracts in the body, it is believed that the genetic resources of Chickso will be valuable and should be preserved. In addition, if a composition with antioxidant and antihypertensive effects is established using conventional beef-derived peptides, it is expected to have applications in preventing, treating, or improving various diseases caused by high blood pressure or free radicals.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This work was carried out with the financial support of the “Cooperative Research Program for Agriculture Science and Technology Development (Project No. PJ016201),” Rural Development Administration, Korea. This work was also supported by the Chung-Ang University, Korea.

Author Contributions

Data curation: Kim JH, Lee DY, Lee SY, Mariano EJ, Jeong JW, Yun SH, Lee J, Hur SJ. Formal analysis: Kim JH. Methodology: Kim JH, Lee DY, Lee SY. Software: Kim JH. Validation: Lee DY, Lee SY, Jo C. Investigation: Kim JH, Lee DY, Lee SY, Mariano EJ, Jeong JW, Yun SH, Lee J, Park J, Choi Y, Han D, Kim JS, Hur SJ. Writing - original draft: Kim JH, Hur SJ. Writing - review & editing: Kim JH, Lee DY, Lee SY, Mariano EJ, Jeong JW, Yun SH, Lee J, Park J, Choi Y, Han D,

Kim JS, Jo C, Hur SJ.

Ethics Approval

All animal experiments were approved by the Experimental Animal Ethics Committee of Chung-Ang University (Approval number: 202-00038) and conducted in accordance with the rules of animal ethics.

References

- Alam M, Cho KH, Lee SS, Choy YH, Kim HS, Cho CI, Choi TJ. 2013. Effect of carcass traits on carcass prices of Holstein steers in Korea. *Asian-Australas J Anim Sci* 26:1388-1398.
- Aluko RE. 2015. Antihypertensive peptides from food proteins. *Annu Rev Food Sci Technol* 6:235-262.
- Arts MJTJ, Haenen GRMM, Voss HP, Bast A. 2004. Antioxidant capacity of reaction products limits the applicability of the trolox equivalent antioxidant capacity (TEAC) assay. *Food Chem Toxicol* 42:45-49.
- Atmaca G. 2004. Antioxidant effects of sulfur-containing amino acids. *Yonsei Med J* 45:776-788.
- Brand-Williams W, Cuvelier ME, Berset CL. 1995. Use of a free radical method to evaluate antioxidant activity. *LWT-Food Sci Technol* 28:25-30.
- Cha SS, Rha YA. 2021. A study on the customers eating out behaviors in food consumption patterns. *Korean J Food Health Converg* 7:7-15.
- Chen HM, Muramoto K, Yamauchi F, Fujimoto K, Nokihara K. 1998. Antioxidative properties of histidine-containing peptides designed from peptide fragments found in the digests of a soybean protein. *J Agric Food Chem* 46:49-53.
- Decker EA, Welch B. 1990. Role of ferritin as a lipid oxidation catalyst in muscle food. *J Agric Food Chem* 38:674-677.
- Dringen R, Hamprecht B. 1999. *N*-acetylcysteine, but not methionine or 2-oxothiazolidine-4-carboxylate, serves as cysteine donor for the synthesis of glutathione in cultured neurons derived from embryonal rat brain. *Neurosci Lett* 259:79-82.
- Escudero E, Mora L, Toldrá F. 2014. Stability of ACE inhibitory ham peptides against heat treatment and *in vitro* digestion. *Food Chem* 161:305-311.
- Esfandi R, Walters ME, Tsopmo A. 2019. Antioxidant properties and potential mechanisms of hydrolyzed proteins and peptides from cereals. *Heliyon* 5:e01538.
- Gaucher C, Boudier A, Bonetti J, Clarot I, Leroy P, Parent M. 2018. Glutathione: Antioxidant properties dedicated to nanotechnologies. *Antioxidants* 7:62.
- Gianfranceschi GL, Gianfranceschi G, Quassinti L, Bramucci M. 2018. Biochemical requirements of bioactive peptides for nutraceutical efficacy. *J Funct Foods* 47:252-263.
- Hamman JH, Enslin GM, Kotzé AF. 2005. Oral delivery of peptide drugs: Barriers and developments. *BioDrugs* 19:165-177.
- Houglund JL, Darling J, Flynn S. 2013. Protein posttranslational modification. In *Molecular basis of oxidative stress: Chemistry, mechanisms, and disease pathogenesis*. Villamena FA (ed). Wiley, Hoboken, NJ, USA. pp 71-92.
- Hur SJ, Lim BO, Decker EA, McClements DJ. 2011. *In vitro* human digestion models for food applications. *Food Chem* 125:1-12.
- Hwang IH. 2004. Effect of palatability traits on satisfactory level of Korean beef consumers. *Korean J Food Sci Anim Resour* 24:310-318.
- Jang A, Lee M. 2005. Purification and identification of angiotensin converting enzyme inhibitory peptides from beef

- hydrolysates. *Meat Sci* 69:653-661.
- Jo C, Cho SH, Chang J, Nam KC. 2012. Keys to production and processing of Hanwoo beef: A perspective of tradition and science. *Anim Front* 2:32-38.
- Joo ST, Hwang YH, Frank D. 2017. Characteristics of Hanwoo cattle and health implications of consuming highly marbled Hanwoo beef. *Meat Sci* 132:45-51.
- Kim HJ, Yang SR, Jang A. 2018. Anti-proliferative effect of a novel anti-oxidative peptide in Hanwoo beef on human colorectal carcinoma cells. *Korean J Food Sci Anim Resour* 38:1168-1178.
- Kim S, Kim K, Roh H, Kim D, Kim S, Kim C, Lee S, Ko Y, Cho C. 2020. Identification of white Hanwoo breed using single nucleotide polymorphism markers. *J Korean Acad Ind Soc* 21:240-246.
- Kim Y, Na HJ. 2001. Food sources of thiamin, riboflavin and niacin based on food composition table and national annual food supply data in food balance sheet. *Korean J Nutr* 34:809-820.
- Lee SH, Park BH, Sharma A, Dang CG, Lee SS, Choi TJ, Choy YH, Kim HC, Jeon KJ, Kim SD, Yeon SH, Park SB, Kang HS. 2014. Hanwoo cattle: Origin, domestication, breeding strategies and genomic selection. *J Anim Sci Technol* 56:2.
- Lee SY, Hur SJ. 2017. Angiotensin converting enzyme inhibitory and antioxidant activities of enzymatic hydrolysates of Korean native cattle (Hanwoo) myofibrillar protein. *BioMed Res Int* 2017:5274637.
- Lee T, Joo N. 2022. Comparison of fatty acid, amino acid, and sensory properties of Chikhanwoo (Korean brindle cattle) and Hanwoo (Korean native cattle). *Culin Sci Hosp Res* 28:18-27.
- Levine RL, Mosoni L, Berlett BS, Stadtman ER. 1996. Methionine residues as endogenous antioxidants in proteins. *Proc Natl Acad Sci USA* 93:15036-15040.
- Levy EJ, Anderson ME, Meister A. 1993. Transport of glutathione diethyl ester into human cells. *Proc Natl Acad Sci USA* 90:9171-9175.
- Li X. 2016. Effects of lysine on the immune and structural barrier functions and the related mechanisms of intestines in fish. Sichuan Agricultural University, Sichuan, China.
- Liu R, Xing L, Fu Q, Zhou GH, Zhang WG. 2016. A review of antioxidant peptides derived from meat muscle and by-products. *Antioxidants* 5:32.
- Mora L, Gallego M, Toldrá F. 2018. ACEI-inhibitory peptides naturally generated in meat and meat products and their health relevance. *Nutrients* 10:1259.
- Oh M, Kim EK, Jeon BT, Tang Y, Kim MS, Seong HJ, Moon SH. 2016. Chemical compositions, free amino acid contents and antioxidant activities of Hanwoo (*Bos taurus coreanae*) beef by cut. *Meat Sci* 119:16-21.
- Olagunju AI, Omoba OS, Enujiugha VN, Alashi AM, Aluko RE. 2018. Pigeon pea enzymatic protein hydrolysates and ultrafiltration peptide fractions as potential sources of antioxidant peptides: An *in vitro* study. *LWT-Food Sci Technol* 97:269-278.
- Oyaizu M. 1986. Studies on products of Browning reactions: Antioxidative activities of product of Browning reaction prepared from glucosamine. *Jpn J Nutr Diet* 44:307-315.
- Park BS, Yoo IJ. 1994. Animal products and processing: Comparison of fatty acid composition among imported beef, Holstein steer beef and Hanwoo beef. *J Anim Sci Technol* 36:69-75.
- Pihlanto A. 2006. Antioxidative peptides derived from milk proteins. *Int Dairy J* 16:1306-1314.
- Piste P. 2013. Cysteine –master antioxidant. *Int J Pharm Chem Biol Sci* 3:143-149.
- Pompella A, Visvikis A, Paolicchi A, De Tata, V, Casini AF. 2003. The changing faces of glutathione, a cellular protagonist.

Biochem Pharmacol 66:1499-1503.

- Power O, Jakeman P, FitzGerald RJ. 2013. Antioxidative peptides: Enzymatic production, *in vitro* and *in vivo* antioxidant activity and potential applications of milk-derived antioxidative peptides. *Amino Acids* 44:797-820.
- Sakanaka S, Tachibana Y, Ishihara N, Juneja LR. 2005. Antioxidant properties of casein calcium peptides and their effects on lipid oxidation in beef homogenates. *J Agric Food Chem* 53:464-468.
- Seol KH, Choe J, Kim HJ, Hwang JT, Lee M, Jo C. 2018. Bioactivities of peptide fractions derived from proteolytic enzyme-injected Hanwoo *longissimus* muscle in a model system. *Int J Food Prop* 21:205-212.
- Sun Y, Pan D, Guo Y, Li J. 2012. Purification of chicken breast protein hydrolysate and analysis of its antioxidant activity. *Food Chem Toxicol* 50:3397-3404.
- Udenigwe CC, Aluko RE. 2012. Food protein-derived bioactive peptides: Production, processing, and potential health benefits. *J Food Sci* 77:R11-R24.
- Utama DT, Lee CW, Park YS, Jang A, Lee SK. 2018. Comparison of meat quality, fatty acid composition and aroma volatiles of Chikso and Hanwoo beef. *Asian-Australas J Anim Sci* 31:1500-1506.
- Vermeirssen V, Deplancke B, Tappenden KA, Van Camp J, Gaskins HR, Verstraete W. 2002. Intestinal transport of the lactokinin Ala-Leu-Pro-Met-His-Ile-Arg through a Caco-2 Bbe monolayer. *J Pept Sci* 8:95-100.
- Whitnall T, Pitts N. 2019. Global trends in meat consumption. *Agric Commod* 9:96-99.