

# Regularity and Normality in Soft Bitopological Ordered Spaces

S. A. El-Sheikh<sup>1</sup>, S. A. Kandil<sup>2</sup>, S. Hussien<sup>3 1</sup>

[slamma\\_elarabi@yahoo.com](mailto:slamma_elarabi@yahoo.com)

<sup>1</sup>Mathematics Department, Faculty of Education, Ain Shams University, Cairo, Egypt.

<sup>2</sup>Mathematics Department, Canadian International college, Cairo, Egypt.

<sup>3</sup>Mathematics Department, Faculty of science, Helwan University, Cairo, Egypt.<sup>1</sup>

Corresponding author. Name : Salama Hussien Ali

## Abstract

This paper examines regularity and normality in soft separation axioms for soft bitopological ordered spaces and their relationships with other properties. The findings expand our understanding of bitopological ordered spaces. Previous research, such as Al-Shami's work [3], has established distinctions between separation axioms in topological ordered spaces, which are more effective in describing these spaces' properties.

### Keywords:

*Soft topology, soft bitopological ordered space and bi-ordered soft regularity and normality.*

## 1. Introduction

Nachbin [17] (1965) introduced the concept of a topological ordered space by adding a partial order to the structure of a topological space. McCartan [13] (1968) used monotone neighborhoods to study ordered separation axioms in these spaces. To address vagueness and uncertainty in real-life problems, mathematical tools like fuzzy sets, intuitionistic fuzzy sets, rough sets, vague sets, and soft sets have been developed. Molodtsov [16] (1999) introduced soft sets as a mathematical tool to handle vagueness and uncertainty. Maji et al. [15] and [14] (2003, 2002) and Aktas and Cagman [1] (2007) further developed soft set theory and its application in decision making problems and algebraic structures. Shabir and Naz [20] (2011) investigated soft separation axioms for crisp points, and Hussain and Ahmad [7] (2011) examined properties of soft interior, closure, and boundary. Nazmul and Samanta [18] (2013) studied neighborhood properties of soft topological spaces, and four different types of separation axioms were defined and discussed in a series of papers (Gocur [10] 2015, Hussain [8] 2015, Shabir 2011 [20], and Tantawy [22] 2016). Singh and Noorie [21] (2017) extended the understanding of soft topological spaces and their properties. Ittanagi [9] (2014) introduced soft bitopological spaces and soft separation axioms, and Kandil et al. [11] (2016) further studied their structures and defined

basic concepts. El-Shafei et al. [3], [4] (2019, 2018) developed two new types of soft relations and introduced the concept of soft topological ordered spaces and ordered soft separation axioms. El-Sheikh et al. [5], [6] (2023, 2022) introduced the concept of soft bitopological ordered spaces and studied the relationships between increasing (decreasing, balancing) pairwise open (closed) soft sets and neighborhoods, and increasing (decreasing) pairwise soft neighbourhoods. This paper discusses the use of soft sets and soft bitopologies in ordered spaces. Section 2 provides definitions and properties of soft sets and soft topologies. Section 3 introduces the concept of "Bi-ordered soft regularity and normality." Examples are provided to demonstrate the connections between these concepts and highlight their characteristics.

## 2. Preliminaries

References and resources, such as [3], [5], [6], [12], [20], can aid in further exploration and comprehension of specialized mathematical concepts, including soft set theory, soft points, soft topological space, soft topological ordered space, and soft bitopological ordered space. From now on, we will use the notation  $X$  to refer to the universe set,  $E$  to denote a fixed set of parameters, and  $2^X$  to represent the power set of  $X$ .

**Definition 2.1** [12] A binary relation  $\leq$  on a set  $X$  is a partial order relation if it is reflexive, anti-symmetric, and transitive. The equality relation on  $X$ , denoted by  $N$ , is defined as  $\{(a,a) : a \in X\}$ .

**Definition 2.2** [17] A triple  $(X, \tau, \leq)$  is called a topological ordered space when  $(X, \tau)$  is a topological space and  $(X, \leq)$  is a partially ordered set.

**Definition 2.3** [2, 15, 16, 20] A soft set is defined as a pair  $(M, E)$ , where  $M : E \rightarrow 2^X$ . The notation  $M_E$  is used instead of  $(M, E)$  for brevity. A soft set can also be represented as a set of ordered pairs, where  $M_E = \{(e, M(e)) | e \in E, M(e) \subseteq 2^X\}$ .

The collection of all soft sets over  $X$  is denoted by  $P(X)^E$ .

A null soft set, denoted by  $\emptyset$ , is one where  $M(e) = \emptyset$  for all  $e \in E$ . An absolute soft set, denoted by  $X_E$ , is one where  $M(e) = X$  for all  $e \in E$ .

Two soft sets,  $M_E, N_E \in P(X)^E$ , are considered a soft subset, denoted by  $N_E \vee M_E$ , if  $N(e) \subseteq M(e)$  for all  $e \in E$ . They are considered equal, denoted by  $N_E = M_E$ , if  $N_E \vee M_E$  and  $M_E \vee N_E$ . The union and intersection of two soft sets,  $N_E$  and  $M_E$ , are represented by  $N_E \cup M_E$  and  $N_E \cap M_E$ , respectively. The difference of two soft sets,  $N_E$  and  $M_E$ , is denoted by  $N_E - M_E$ , and the complement of a soft set  $N_E$  is denoted by  $N_E^c$ .

**Definition 2.4** [18, 19] A soft set  $N_E : E \rightarrow 2^X$  defined as  $N(\alpha) = \alpha$  if  $\alpha = e$  and  $N(\alpha) = \emptyset$  if  $\alpha \in E - \{e\}$  is called a soft point and denoted by  $a^e$ . The collection of all soft points over  $X$  is denoted by  $Sp(X)^E$ . A soft point  $a^e$  is said to be belonging to a soft set  $N_E$ , denoted by  $a^e \in N_E$ , if for the member  $e \in E$ ,  $a^e \subseteq N(e)$ .

**Definition 2.5** [3, 16] For a soft set  $N_E$  over  $X$  and an element  $a \in X$ , we say  $a \in N_E$  if  $a \in N(\alpha)$  for every  $\alpha \in E$  and  $a \in N_E$  if  $a \in N(\alpha)$  for some  $\alpha \in E$ .

We say  $a \in N_E$  if  $a \in N(\alpha)$  for some  $\alpha \in E$  and  $a \notin N_E$  if  $a \notin N(\alpha)$  for every  $\alpha \in E$ . The notations  $\in, \notin, \in$  and  $\notin$  are respectively read as belong, non-belong, partial belong and total non-belong relations.

**Definition 2.6** [20] A soft topology on a set  $X$  is a collection of soft sets over  $X$  with a certain set of properties. Specifically, the collection must contain

the null soft set and the absolute soft set, and it must be closed under arbitrary unions and finite intersections.

A soft topological space is a triple  $(X, \tau, E)$  where  $\tau$  is a soft topology on  $X$ . The members of  $\tau$  are called soft open sets, and their complements are called soft closed sets.

**Definition 2.7** [18] Let  $P(X)^E$  and  $P(Y)^K$  be families of soft sets over  $X$  and  $Y$ , respectively. Let  $\phi : X \rightarrow Y$  and  $\psi : E \rightarrow K$  be two mappings. The mapping  $\phi_\psi : P(X)^E \rightarrow P(Y)^K$  is a soft mapping from  $X$  to  $Y$ , denoted by  $\phi_\psi$ , defined as follows:

1. For  $G_E \in P(X)^E$ ,  $\phi_\psi(G_E)(k) = \bigcup_{e \in \psi^{-1}(k)} G(e)$  if  $\psi^{-1}(k) \neq \emptyset$ , and  $\phi_\psi(G_E)(k) = \emptyset$  otherwise, for all  $k \in K$ . The soft set  $\phi_\psi(G_E)$  is called the soft image of  $G_E$ .
2. For  $F_K \in P(Y)^K$ ,  $\phi_\psi^{-1}(F_K)(e) = \phi^{-1}(F(\psi(e)))$ , for all  $e \in E$ . The soft set  $\phi_\psi^{-1}(F_K)$  is called the soft inverse image of  $F_K$ .

**Definition 2.8** [23] Let  $P(X)^E$  and  $P(Y)^K$  be two families of soft sets over  $X$  and  $Y$ , respectively. A soft mapping  $\phi_\psi : P(X)^E \rightarrow P(Y)^K$  is called soft surjective( injective) mapping if  $\phi, \psi$  are surjective( injective) mappings, respectively. A soft mapping which is a soft surjective and soft injective mapping is called a soft bijection mapping.

**Proposition 2.1** [18] Consider  $\phi_\psi : P(X)^E \rightarrow P(Y)^K$  is a soft map and let  $G_E$  and  $H_K$  be two soft subsets of  $P(X)^E$  and  $P(Y)^K$ , respectively. Then we have the following results:

1.  $G_E \subseteq \phi_\psi^{-1}(\phi_\psi(G_E))$  and the equality relation holds if  $\phi_\psi$  is injective.
2.  $\phi_\psi(\phi_\psi^{-1}(H_K)) \vee H_K$  and the equality relation holds if  $\phi_\psi$  is surjective.

**Definition 2.9** [18] A soft map  $\phi_\psi : (X, \tau, E) \rightarrow (Y, \eta, K)$  is said to be:

1. Soft continuous if the inverse image of each soft open subset of  $(Y, \eta, K)$  is a soft open subset of  $(X, \tau, E)$ .
2. Soft open ( resp. soft closed ) if the image of each soft open ( resp. soft closed ) subset of  $(X, \tau, E)$  is a soft open ( resp. soft closed ) subset of  $(Y, \eta, K)$ .
3. Soft homeomorphism if it is bijective, soft continuous and soft open.

**Definition 2.10** [9, 11] A quadrable system  $(X, \tau_1, \tau_2, E)$  is called a soft bitopological space when  $\tau_1$  and  $\tau_2$  are soft topologies on the set  $X$  with a fixed set of parameters  $E$ . A soft set  $N_E$  in a soft bitopological space  $(X, \tau_1, \tau_2, E)$  is called pairwise open soft (PO-soft) if there exists a  $\tau_1$ -open soft set  $N_E^1$  and a  $\tau_2$ -open soft set  $N_E^2$  such that  $N_E = N_E^1 \cup N_E^2$ , and pairwise closed soft (PC-soft) if the complement of  $N_E$  is a PO-soft set. The family of all PO-soft sets, denoted by  $\tau_{12}$ , is a supra soft topological space associated with the soft bitopological space  $(X, \tau_1, \tau_2, E)$ .

**Definition 2.11** [3] A triple  $(X, E, \leq)$  is called a partially ordered soft space when  $\leq$  is a partial order relation on the set  $X$ . An increasing soft operator  $i : (P(X)^E, \leq) \rightarrow (P(X)^E, \leq)$  and a decreasing soft operator  $d : (P(X)^E, \leq) \rightarrow (P(X)^E, \leq)$  are defined for each soft set  $N_E$  in  $P(X)^E$  by  $i(N_E)(e) = iN(e) = \{a \in X : b \leq a, \text{ for some } b \in N(e)\}$  and  $d(N_E)(e) = dN(e) = \{a \in X : a \leq b, \text{ for some } b \in N(e)\}$  respectively. A soft subset  $N_E$  of the partially ordered soft space  $(X, E, \leq)$  is called increasing if  $N_E = i(N_E)$ , decreasing if  $N_E = d(N_E)$ .

**Proposition 2.2** [3] The following two results hold for a soft map  $\phi_\psi : P(X)^E \rightarrow P(Y)^K$ .

1. The image of each soft point is soft point.

2. If  $\phi_\psi$  is bijective, then the inverse image of each soft point is soft point.

**Definition 2.12** [3] Let  $x^e$  and  $y^e$  be two soft points in a partially ordered soft set  $(X, E, \leq)$ . Then  $x^e \leq y^e$  if  $x \leq y$ .

**Definition 2.13** [3] A soft map  $\phi_\psi : (P(X)^E, \leq_1) \rightarrow (P(Y)^K, \leq_2)$  is said to be:

1. Increasing if  $x^e \leq_1 y^e$ , then  $\phi_\psi(x^e) \leq_2 \phi_\psi(y^e)$ .
2. Decreasing if  $x^e \leq_1 y^e$ , then  $\phi_\psi(y^e) \leq_2 \phi_\psi(x^e)$ .
3. Ordered embedding if  $x^e \leq_1 y^e$  if and only if  $\phi_\psi(x^e) \leq_2 \phi_\psi(y^e)$ .

**Theorem 2.1** [3] The following two results hold for a soft map  $\phi_\psi : (P(X)^E, \leq_1) \rightarrow (P(Y)^K, \leq_2)$ .

1. If  $\phi_\psi$  is increasing, then the inverse image of each increasing ( resp. decreasing ) soft subset of  $Y_K$  is an increasing ( resp. decreasing ) soft subset of  $X_E$ .
2. If  $\phi_\psi$  is decreasing, then the inverse image of each increasing ( resp. decreasing ) soft subset of  $Y_K$  is an decreasing ( resp. increasing ) soft subset of  $X_E$ .

**Theorem 2.2** [3] Let  $\phi_\psi : (P(X)^E, \leq_1) \rightarrow (P(Y)^K, \leq_2)$  be a bijective ordered embedding soft map. Then the image of each increasing ( resp. decreasing ) soft subset of  $X_E$  is an increasing ( resp. decreasing ) soft subset of  $Y_K$ .

**Definition 2.14** [3] A quadrable system  $(X, \tau, E, \leq)$  can be rephrased as a soft topological ordered space (STOS) if  $(X, \tau, E)$  is a soft topological space and  $(X, E, \leq)$  is a partially ordered soft space.

**Definition 2.15** [3] A soft subset  $W_E$  of an STOS  $(X, \tau, E, \leq)$  is called increasing (resp. decreasing) soft neighborhood of  $x \in X$  if  $W_E$  is soft neighborhood of  $x$  and increasing (resp. decreasing).

**Definition 2.16** [5] The system  $(X, \tau_1, \tau_2, E, \leq)$  is said to be a soft bitopological ordered space (SBTOS), if

$(X, \tau_1, \tau_2, E)$  is a soft bitopological space and  $(X, E, \leq)$  is a partially ordered soft space.

**Definition 2.17** [5] A soft set  $W_E$  in a SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is called:

1. ITPS (IPPS)- nbd of  $a \in X$  if  $W_E$  is a total ( partial ) pairwise soft neighborhood of  $a$  and increasing.
2. DTPS (DPPS)- nbd of  $a \in X$  if  $W_E$  is a total ( partial ) pairwise soft neighborhood of  $a$  and decreasing.

**Definition 2.18** [6] An SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is said to be:

1.  $LPST_1 (LPST_1^{**})$ - ordered if for every distinct points  $x, y$  in  $X$  such that  $x \not\leq y$  there exists an ITPS (IPPS)- nbd  $W_E$  of  $x$  such that  $y \in W_E$ .
2.  $LPST_1^\bullet (LPST_1^*)$ - ordered if for every distinct points  $x, y$  in  $X$  such that  $x \not\leq y$  there exists an ITPS (IPPS)- nbd  $W_E$  of  $x$  such that  $y \in W_E$ .
3.  $UPST_1 (UPST_1^{**})$ - ordered if for every distinct points  $x, y$  in  $X$  such that  $x \not\leq y$  there exists a DTPS (DPPS)- nbd  $W_E$  of  $y$  such that  $x \in W_E$ .
4.  $UPST_1^\bullet (UPST_1^*)$ - ordered if for every distinct points  $x, y$  in  $X$  such that  $x \not\leq y$  there exists a DTPS (DPPS)- nbd  $W_E$  of  $y$  such that  $x \in W_E$ .
5.  $PST_0 (PST_0^\bullet)$ - ordered space if it is  $LPST_1 (LPST_1^\bullet)$ - ordered or  $UPST_1 (UPST_1^\bullet)$ - ordered.
6.  $PST_0^* (PST_0^{**})$ - ordered space if it is  $LPST_1^* (LPST_1^{**})$ - ordered or  $UPST_1^* (UPST_1^{**})$ - ordered.
7.  $PST_1 (PST_1^\bullet)$ - ordered space if it is  $LPST_1 (LPST_1^\bullet)$ - ordered and  $UPST_1 (UPST_1^\bullet)$ - ordered.

8.  $PST_1^* (PST_1^{**})$ - ordered space if it is  $LPST_1^* (LPST_1^{**})$ - ordered and  $UPST_1^* (UPST_1^{**})$ - ordered.

9.  $PST_2 (PST_2^*)$ - ordered space if for every distinct points  $x, y$  in  $X$  such that  $x \not\leq y$  there exist disjoint total ( partial ) pairwise soft neighborhoods  $W_E$  and  $V_E$  of  $x$  and  $y$ , respectively, such that  $W_E$  is increasing and  $V_E$  is decreasing.

10.  $PST_2^\bullet (PST_2^{**})$ - ordered space if for every distinct points  $x, y$  in  $X$  such that  $x \not\leq y$  there exist disjoint

total ( partial ) pairwise soft neighborhood  $W_E$  of  $x$  and partial ( total ) pairwise soft neighborhood  $V_E$  of  $y$  such that  $W_E$  is increasing and  $V_E$  is decreasing.

### 3. Bi-Ordered Soft Regularity and Normality

In this section, a new concept called "Bi-ordered soft regularity and normality" is introduced and its properties are examined. Various examples are provided to demonstrate the relationships among these concepts and to illustrate the results obtained.

**Definition 3.1** For two soft subsets  $G_E$  and  $H_E$  of an SBTOS  $(X, \tau_1, \tau_2, E, \leq)$ , we say that  $G_E$  is pairwise soft neighborhood of  $H_E$  provided that there exists a pairwise open soft set  $F_E$  such that  $H_E \vee F_E \vee G_E$ .

**Definition 3.2** An SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is said to be:

1. Lower ( resp. upper ) PT- soft regularly ordered if for every decreasing ( resp. increasing ) pairwise closed soft set  $H_E$  and  $x \in X$  such that  $x \in H_E$  there exist disjoint pairwise soft neighborhood  $W_E$  of  $H_E$  and increasing ( resp. decreasing ) total pairwise soft neighborhood  $V_E$  of  $x$  such that  $W_E$  is decreasing ( resp. increasing ).
2. Lower ( resp. upper ) PP- soft regularly ordered if for every decreasing ( resp.

- increasing ) pairwise closed soft set  $H_E$  and  $x \in X$  such that  $x \notin H_E$  there exist disjoint pairwise soft neighborhood  $W_E$  of  $H_E$  and increasing ( resp. decreasing ) partial pairwise soft neighborhood  $V_E$  of  $x$  such that  $W_E$  is decreasing ( resp. increasing ).
3. Lower ( resp. upper )  $P^*T$ - soft regularly ordered if for every decreasing ( resp. increasing ) pairwise closed soft set  $H_E$  and  $x \in X$  such that  $x \notin H_E$  there exist disjoint pairwise soft neighborhood  $W_E$  of  $H_E$  and increasing ( resp. decreasing ) total pairwise soft neighborhood  $V_E$  of  $x$  such that  $W_E$  is decreasing ( resp. increasing ).
  4. Lower ( resp. upper )  $P^*P$ - soft regularly ordered if for every decreasing ( resp. increasing ) pairwise closed soft set  $H_E$  and  $x \in X$  such that  $x \notin H_E$  there exist disjoint pairwise soft neighborhood  $W_E$  of  $H_E$  and increasing ( resp. decreasing ) partial pairwise soft neighborhood  $V_E$  of  $x$  such that  $W_E$  is decreasing ( resp. increasing ).
  5. Total  $P$ - soft regularly ordered if it is both Lower  $PT$ - soft regularly ordered and upper  $PT$ - soft regularly ordered.
  6. Partial  $P$ - soft regularly ordered if it is both Lower  $PP$ - soft regularly ordered and upper  $PP$ - soft regularly ordered.
  7. Total  $P^*$ - soft regularly ordered if it is both Lower  $P^*T$ - soft regularly ordered and upper  $P^*T$ - soft regularly ordered.
  8. Partial  $P^*$ - soft regularly ordered if it is both Lower  $P^*P$ - soft regularly ordered and upper  $P^*P$ - soft regularly ordered.
  9. Lower ( resp. upper )  $TP$ - soft  $T_3$  ordered if it is both  $LPST_1$ -ordered ( resp.

$UPST_1$ -ordered ) and lower ( resp. upper )  $PT$ - soft regularly ordered.

10. Lower ( resp. upper )  $PP$ - soft  $T_3$  ordered if it is both  $LPST_1^{**}$ -ordered ( resp.  $UPST_1^{**}$ -ordered ) and lower ( resp. upper )  $PP$ - soft regularly ordered.
11. Lower ( resp. upper )  $TP^*$ - soft  $T_3$  ordered if it is both  $LPST_1^\bullet$ -ordered ( resp.  $UPST_1^\bullet$ -ordered ) and lower ( resp. upper )  $P^*T$ - soft regularly ordered.
12. Lower ( resp. upper )  $PP^*$ - soft  $T_3$  ordered if it is both  $LPST_1^*$ -ordered ( resp.  $UPST_1^*$ -ordered ) and lower ( resp. upper )  $P^*P$ - soft regularly ordered.
13.  $TP$ - soft  $T_3$  ordered if it is both lower  $TP$ - soft  $T_3$  ordered and upper  $TP$ - soft  $T_3$  ordered.
14.  $PP$ - soft  $T_3$  ordered if it is both lower  $PP$ - soft  $T_3$  ordered and upper  $PP$ - soft  $T_3$  ordered.
15.  $TP^*$ - soft  $T_3$  ordered if it is both lower  $TP^*$ - soft  $T_3$  ordered and upper  $TP^*$ - soft  $T_3$  ordered.
16.  $PP^*$ - soft  $T_3$  ordered if it is both lower  $PP^*$ - soft  $T_3$  ordered and upper  $PP^*$ - soft  $T_3$  ordered.

**Theorem 3.1** An SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is lower ( resp. upper )  $PT$ -soft regularly ordered if and only if for all  $x \in X$  and every increasing ( resp. decreasing ) pairwise open soft set  $U_E$  containing  $x$ , there is an increasing ( resp. decreasing ) total pairwise soft neighbourhood  $V_E$  of  $x$  satisfies that  $cl_{12}^s(V_E) \sqsubseteq U_E$ .

**Proof.** Necessity: Let  $x \in X$  and  $U_E$  be an increasing pairwise open soft set partially containing  $x$ . Then,  $U_E^c$  is decreasing pairwise closed soft such that  $x \notin U_E^c$ . By hypothesis, there exist disjoint decreasing pairwise soft neighbourhood  $W_E$  of  $U_E^c$  and increasing total pairwise soft neighbourhood  $V_E$  of  $x$ . So there is a pairwise open soft set  $G_E$  such that  $U_E^c \sqsubseteq G_E \sqsubseteq W_E$ . Since  $V_E \sqsubseteq W_E^c$ , then  $V_E \sqsubseteq W_E^c \sqsubseteq G_E^c \sqsubseteq U_E$  and since  $G_E^c$  is soft closed, then  $cl_{12}^s(V_E) \sqsubseteq G_E^c \sqsubseteq U_E$ .

Sufficiency: Let  $x \in X$  and  $H_E$  be a decreasing pairwise closed soft set such that  $x \in H_E$ . Then  $H_E^c$  be an increasing pairwise open soft set containing  $x$ . So that, by hypothesis, there is an increasing total pairwise soft neighbourhood  $V_E$  of  $x$  such that  $cl_{12}^s(V_E) \subseteq H_E^c$ . Consequently,  $(cl_{12}^s(V_E))^c$  is a pairwise open soft set containing  $H_E$ . Thus  $d((cl_{12}^s(V_E))^c)$  is a decreasing pairwise soft neighbourhood of  $H_E$ . Suppose that  $V_E \cap d((cl_{12}^s(V_E))^c) \neq \hat{\phi}$ .

Then there exists  $z \in X$  such that  $z \in V_E$  and  $z \in d((cl_{12}^s(V_E))^c)$ . So there exists  $y \in ((cl_{12}^s(V_E))^c)(e)$  satisfies that  $z \leq y$ . This means that  $y \in V(e)$ . But this contradicts the disjointness between  $V_E$  and  $(cl_{12}^s(V_E))^c$ . Thus  $V_E \cap d((cl_{12}^s(V_E))^c) = \hat{\phi}$ . This completes the proof.

A similar proof can be given for the case between parentheses.

**Theorem 3.2** An SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is lower ( resp. upper )  $P^*T$ -soft regularly ordered if and only if for all  $x \in X$  and every increasing ( resp. decreasing ) pairwise open soft set  $U_E$  containing  $x$ , there is an increasing ( resp. decreasing ) total pairwise soft neighbourhood  $V_E$  of  $x$  satisfies that  $cl_{12}^s(V_E) \subseteq U_E$ .  
 Proof. The proof is similar to the proof of Theorem 3.1.

**Theorem 3.3** An SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is lower ( resp. upper )  $PT$  (  $PP, P^*P$  )-soft regularly ordered if and only if for all  $x \in X$  and every increasing ( resp. decreasing ) pairwise open soft set  $U_E$  containing  $x$ , there is an increasing ( resp. decreasing ) partial pairwise soft neighbourhood  $V_E$  of  $x$  satisfies that  $cl_{12}^s(V_E) \subseteq U_E$ .  
 Proof. The proof is similar to the proof of Theorem 3.1.

**Proposition 3.1** Every  $TP$ -soft  $T_3$ -ordered space  $(X, \tau_1, \tau_2, E, \leq)$  is  $PP$ -soft  $T_3$ -ordered.  
 Proof. The proposition's proof establishes that the belong relation, denoted by  $\in$ , can be extended to a partial belong relation denoted by  $b$ .

**Example 3.1** Let  $E = \{e_\alpha, e_\beta, e_\gamma\}$  be a set of parameters,  $\leq = N \cup \{(1,2)\}$  be a partial order relation on the set of natural numbers  $N$ . Define  $\tau_1 = \{G_E \vee N_E \text{ such that}$

$1 \in G_E \text{ or } [1 \in G(e_\beta) \text{ and } G_E^c \text{ is finite }]\}$  and  $\tau_2 = \{F_E \vee N_E \text{ such that } 3 \in F(e_\alpha) \text{ and } F_E^c \text{ is finite}\}$ . Obviously,  $(N, \tau_1, \tau_2, E, \leq)$  is  $PP$ -soft  $T_3$ -ordered [3]. A soft subset  $H_E$  of  $(N, \tau_1, \tau_2, E, \leq)$  is a decreasing pairwise closed soft set if  $[1 \in H_E \text{ and } H_E \text{ is infinite }]$  or  $[1 \in H(e_\beta), 3 \in H(e_\alpha) \text{ and } H_E \text{ is finite }]$ . To illustrate that  $(N, \tau_1, \tau_2, E, \leq)$  is not lower  $PT$ -soft regularly ordered, we define a decreasing soft closed set  $H_E$  as follows:  $H_E = \{(e_\alpha, \{1,2\}), (e_\beta, \{3\}), (e_\gamma, \{1,2\})\}$ .

Since  $1 \in H_E$  and there do not exist disjoint soft neighbourhoods  $W_E$  and  $V_E$  containing  $H_E$  and  $1$ , respectively, then  $(N, \tau_1, \tau_2, E, \leq)$  is not lower  $PT$ -soft regularly ordered. Hence  $(N, \tau_1, \tau_2, E, \leq)$  is not  $TP$ -soft  $T_3$ -ordered.

**Proposition 3.2** An SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is  $TP$ -soft  $T_3$ -ordered if and only if  $TP^*$ -soft  $T_3$ -ordered.

Proof. On the one hand,  $x \in G_E$  implies that  $x \in G_E$ , then  $TP$ -soft  $T_3$ -ordered implies  $PP$ -soft  $T_3$ -ordered. On the other hand, the definition of  $PP$ -soft  $T_3$ -ordered implies that for every decreasing ( resp. increasing ) pairwise closed soft set  $H_E$  and  $x \in X$  such that  $x \in H_E$ , there exist disjoint pairwise soft neighbourhood  $W_E$  of  $H_E$  and increasing ( resp. decreasing ) total pairwise soft neighbourhood  $V_E$  of  $x$ , such that  $W_E$  is decreasing ( resp. increasing ). Since  $W_E$  and  $V_E$  are disjoint, then  $x \in H_E$  and  $\forall x \in H_E$ , there exist an  $ITPS$   $N_E$  of  $x$  such that  $y \in N_E$ . Hence the definitions of  $TP$ -soft  $T_3$ -ordered and  $PP$ -soft  $T_3$ -ordered are equivalent.

**Corollary 3.1** An SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is  $PP$ -soft  $T_3$ -ordered if and only if  $PP^*$ -soft  $T_3$ -ordered.

**Proposition 3.3** Every  $TP^*$ -soft  $T_3$ -ordered space  $(X, \tau_1, \tau_2, E, \leq)$  is  $PP^*$ -soft  $T_3$ -ordered.

**Proof.** The proof for the proposition states that the belong relation  $\in$  implies a partial belong relation  $b$ . The proposition is demonstrated by proving that the belong relation, represented by  $\in$ , can be fully extended to a relation of partial belong, denoted by  $b$ .

**Example 3.2** From Example 3.1, an SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is  $PP^*$ -soft  $T_3$ -ordered but it is not  $TP^*$ -soft  $T_3$ -ordered.

**Proposition 3.4** The following three properties are equivalent if  $(X, \tau_1, \tau_2, E, \leq)$  is  $TP^*$ -soft regularly ordered:

1.  $(X, \tau_1, \tau_2, E, \leq)$  is  $PST_2$ -ordered;
2.  $(X, \tau_1, \tau_2, E, \leq)$  is  $PST_1$ -ordered;
3.  $(X, \tau_1, \tau_2, E, \leq)$  is  $PST_0$ -ordered.

**Proof.** The direction 1)  $\rightarrow$  2)  $\rightarrow$  3) is obvious from Propositions 3.1, 3.2 and 3.3 in [6].

To prove 3)  $\rightarrow$  1), let  $x, y \in X$  such that  $x \leq y$ . Since  $(X, \tau_1, \tau_2, E, \leq)$  is  $PST_0$ -ordered, then it is lower pairwise soft  $T_1$ -ordered or upper pairwise soft  $T_1$ -ordered. Say it is upper pairwise soft  $T_1$ -ordered. From Theorem 3.1 in [6], we have that  $(i(x))_E$  is pairwise soft closed. Obviously,  $(i(x))_E$  is increasing and  $y \leq (i(x))_E$ . Since  $(X, \tau_1, \tau_2, E, \leq)$  is  $TP^*$ -soft regularly ordered, then there exist disjoint decreasing total pairwise soft neighbourhood  $W_E$  of  $y$  and increasing pairwise soft neighbourhood  $V_E$  of  $(i(x))_E$  so  $V_E$  is increasing total pairwise soft neighbourhood of  $x$ . Thus  $(X, \tau_1, \tau_2, E, \leq)$  is  $PST_2$ -ordered.

**Corollary 3.2** The following three properties are equivalent if  $(X, \tau_1, \tau_2, E, \leq)$  is lower (upper)  $P^*T$ -soft regularly ordered:

1.  $(X, \tau_1, \tau_2, E, \leq)$  is  $PST_2$ -ordered;
2.  $(X, \tau_1, \tau_2, E, \leq)$  is  $PST_1$ -ordered;
3.  $(X, \tau_1, \tau_2, E, \leq)$  is  $LPST_1$  (resp.  $UPST_1$ )-ordered.

**Proposition 3.5** Every  $TP^*$ -soft  $T_3$ -ordered space  $(X, \tau_1, \tau_2, E, \leq)$  is also  $PST_2$ -ordered.

**Proof.** Proposition 3.4 implies that any  $TP^*$ -soft  $T_3$ -ordered space is also  $PST_2$ -ordered.

Here is an illustration that shows that the converse of Proposition 3.5 is not necessarily true.

**Example 3.3** Let  $E = \{e_\alpha, e_\beta\}$  be a set of parameters,  $\leq = N \cup \{(1,2)\}$  be a partial order relation on the set of natural numbers  $N$ . Define  $\tau_1 = \{G_E \vee N_E \text{ such that } 1 \in G_E \text{ and } G_E^c \text{ is infinite}\}$  and  $\tau_2 = \{F_E \vee N_E \text{ such that } 1 \in F_E^c\} \cup N_E$ . Then  $(N, \tau_1, \tau_2, E, \leq)$  is a soft bitopological ordered space. We have the following 6 cases:

For  $x, y \in N - \{1,2\}$ ,  $x \neq y$ :

1. Either  $1 \leq x$ . Then we define two soft sets  $W_E$  and  $V_E$  as follows  $W_E = \{(e_\alpha, \{1,2\}), (e_\beta, \{1,2\})\}$  and  $V_E = \{(e_\alpha, \{x\}), (e_\beta, \{x\})\}$ . So  $W_E$  is an increasing total pairwise soft neighbourhood of  $1, V_E$  is a decreasing total pairwise soft neighbourhood of  $x$  and  $W_E \cup V_E = \emptyset$ .
2. Or  $x \leq 1$ . Then we define two soft sets  $W_E$  and  $V_E$  as follows  $W_E = \{(e_\alpha, \{x\}), (e_\beta, \{x\})\}$  and  $V_E = \{(e_\alpha, \{1\}), (e_\beta, \{1\})\}$ . So  $W_E$  is an increasing total pairwise soft neighbourhood of  $x, V_E$  is a decreasing total pairwise soft neighbourhood of  $1$  and  $W_E \cup V_E = \emptyset$ .
3. Or  $2 \leq x$ . Then we define two soft sets  $W_E$  and  $V_E$  as follows  $W_E = \{(e_\alpha, \{2\}), (e_\beta, \{2\})\}$  and  $V_E = \{(e_\alpha, \{x\}), (e_\beta, \{x\})\}$ . So  $W_E$  is an increasing total pairwise soft neighbourhood of  $2, V_E$  is a decreasing total pairwise soft neighbourhood of  $x$  and  $W_E \cup V_E = \emptyset$ .
4. Or  $2 \leq 1$ . Then we define two soft sets  $W_E$  and  $V_E$  as follows  $W_E = \{(e_\alpha, \{2\}), (e_\beta, \{2\})\}$  and  $V_E = \{(e_\alpha, \{1\}), (e_\beta, \{1\})\}$ . So  $W_E$  is an increasing total pairwise soft neighbourhood of  $2, V_E$  is a decreasing total pairwise soft neighbourhood of  $1$  and  $W_E \cup V_E = \emptyset$ .
5. Or  $x \leq 2$ . Then we define two soft sets  $W_E$  and  $V_E$  as follows  $W_E = \{(e_\alpha, \{x\}), (e_\beta, \{x\})\}$  and  $V_E = \{(e_\alpha, \{1,2\}), (e_\beta, \{1,2\})\}$ . So  $W_E$  is an increasing total pairwise soft neighbourhood of

$x, V_E$  is a decreasing total pairwise soft neighbourhood of 2 and  $W_E \cup V_E = \phi$ .

6. Or  $x \leq y$ . Then we define two soft sets  $W_E$  and  $V_E$  as follows  $W_E = \{(e_\alpha, \{x\}), (e_\beta, \{x\})\}$  and  $V_E = \{(e_\alpha, \{y\}), (e_\beta, \{y\})\}$ . So  $W_E$  is an increasing total pairwise soft neighbourhood of  $x, V_E$  is a decreasing total pairwise soft neighbourhood of  $y$  and  $W_E \cup V_E = \phi$ .

To illustrate that  $(X, \tau_1, \tau_2, E, \leq)$  is not lower PT-soft regularly ordered, we define a decreasing pairwise soft set  $H_E$  as follows:  $H_E = \{(e_\alpha, \{1,2,4,5,\dots\}), (e_\beta, \{1,2,4,5,\dots\})\}$ . Since  $3 \in 6$   $H_E$  and there do not exist disjoint pairwise soft neighbourhoods  $W_E$  and  $V_E$  of  $H_E$  and 3, respectively, then  $(X, \tau_1, \tau_2, E, \leq)$  is not lower PT-soft regularly ordered.

**Definition 3.3** An SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is said to be:

1. Soft pairwise normally ordered if for each disjoint pairwise closed soft sets  $F_E$  and  $H_E$  such that  $F_E$  is increasing and  $H_E$  is decreasing, there exist disjoint pairwise soft neighborhoods  $W_E$  of  $F_E$  and  $V_E$  of  $H_E$  such that  $W_E$  is increasing and  $V_E$  is decreasing.
2. TP-soft  $T_4$ -ordered if it is soft pairwise normally ordered and  $PST_1^*$ -ordered.

**Theorem 3.4** An SBTOS  $(X, \tau_1, \tau_2, E, \leq)$  is soft pairwise normally ordered if and only if for every decreasing ( resp. increasing ) pairwise closed soft set  $F_E$  and every decreasing ( resp. increasing ) pairwise soft neighbourhood  $U_E$  of  $F_E$ , there is a decreasing ( resp. increasing ) pairwise soft neighbourhood  $V_E$  of  $F_E$ , satisfies that  $cl_{12}^s(V_E) \subseteq U_E$ .

**Proof.** Necessity: Let  $F_E$  be a decreasing pairwise closed soft set and  $U_E$  be a decreasing pairwise soft neighbourhood of  $F_E$ . Then,  $U_E^c$  is an increasing pairwise closed soft set and  $F_E \cup U_E^c = \phi$ . Since  $(X, \tau_1, \tau_2, E, \leq)$  is soft pairwise normally ordered, then there exist disjoint decreasing pairwise soft neighbourhood  $V_E$  of  $F_E$  and increasing pairwise soft neighbourhood  $W_E$  of  $U_E^c$ . Since  $W_E$  is a pairwise soft neighbourhood of  $U_E^c$ , then there exists a pairwise

closed soft set  $H_E$  such that  $U_E^c \subseteq H_E \subseteq W_E$ . Consequently,  $W_E^c \subseteq H_E^c \subseteq U_E$  and  $V_E \subseteq W_E^c$ .

So it follows that  $cl_{12}^s(V_E) \subseteq cl_{12}^s(W_E^c) \subseteq H_E^c \subseteq U_E$ . Thus  $F_E \subseteq cl_{12}^s(V_E) \subseteq cl_{12}^s(W_E^c) \subseteq H_E^c \subseteq U_E$ . Hence the necessity part holds.

Sufficiency: Let  $F_E^1$  and  $F_E^2$  be two disjoint pairwise closed soft sets such that  $F_E^1$  is decreasing and  $F_E^2$  is increasing. Then  $F_E^{2c}$  is a decreasing pairwise open soft set containing  $F_E^1$ . By hypothesis, there exists a decreasing pairwise soft neighbourhood  $V_E$  of  $F_E^1$  such that  $cl_{12}^s(V_E) \subseteq F_E^{2c}$ . Setting  $H_E = X_E - cl_{12}^s(V_E)$ . This means that  $H_E$  is a pairwise open soft set containing  $F_E^2$ . Obviously,  $F_E^2 \cap H_E, F_E^1 \cap V_E$  and  $H_E \cup V_E = \phi$ . Now,  $i(H_E)$  is an increasing pairwise soft neighbourhood of  $F_E^2$ . Suppose that  $i(H_E) \cup V_E \neq \phi$ . Then there exists  $e \in E$  and  $x \in X$  such that  $x \in i(H_E)$  and  $x \in V(e) = d(V(e))$ . This implies that there exist  $a \in H(e)$  and  $b \in V(e)$  such that  $a \leq x$  and  $x \leq b$ . As  $\leq$  is transitive, then  $a \leq b$ . Therefore  $b \in H_E \cup V_E$ . This contradicts the disjointness between  $H_E$  and  $V_E$ . Thus  $i(H_E) \cup V_E = \phi$ . Hence the proof is completed.

A proof similar can be given for the statement inside the parentheses.

**Proposition 3.6** Every TP- soft  $T_4$ -ordered space  $(X, \tau_1, \tau_2, E, \leq)$  is also TP\*- soft  $T_3$ -ordered.

**Proof.** Let  $a \in X$  and  $F_E$  be a decreasing pairwise closed soft set such that  $a \in F_E$ . Since  $(X, \tau_1, \tau_2, E, \leq)$  is PST\*-ordered, then  $(i(a))_E$  is an increasing pairwise closed soft set and since  $(X, \tau_1, \tau_2, E, \leq)$  is soft pairwise normally ordered, then there exist disjoint pairwise soft neighbourhood  $W_E$  and  $V_E$  of  $(i(a))_E$  and  $F_E$  respectively, such that  $W_E$  is increasing and  $V_E$  is decreasing. Therefore,  $(X, \tau_1, \tau_2, E, \leq)$  is lower PT-soft regularly ordered. If  $F_E$  is an increasing pairwise soft set, then we prove similarly that  $(X, \tau_1, \tau_2, E, \leq)$  is upper

PT-soft regularly ordered. Thus  $(X, \tau_1, \tau_2, E, \leq)$  is TP\*-soft regularly ordered. Hence  $(X, \tau_1, \tau_2, E, \leq)$  is TP\*-soft  $T_3$ -ordered. The converse of the above proposition is not always true as illustrated in the following example.

**Example 3.4** From Example (4.28) in [3], if we take  $\tau_1 = \{G_E \vee N_E \text{ such that } 1 \in G_E\}$  and  $\tau_2 = \{F_E \vee N_E \text{ such that } 1 \in F(e_2) \text{ and } F_E^c \text{ is finite}\}$ . Then we have  $(N, \tau_1, \tau_2, E, \leq)$  is TP\*-soft  $T_3$ -ordered, but it is not TP-soft  $T_4$ -ordered.

**Theorem 3.5** The property of being a  $PST_i (PST_i^\bullet, PST_i^*, PST_i^{**})$ -ordered space is soft bitopological ordered property, for  $i = 0, 1, 2$ .

**Proof.** We prove the theorem in case of  $PST_2$  and the other follow similar lines.

Suppose that  $\phi_\psi$  is an ordered embedding soft homeomorphism map of a  $PST_2$ -ordered space  $(X, \tau_1, \tau_2, E, \leq_1)$  on to an SBTOS  $(Y, \eta_1, \eta_2, K, \leq_2)$  and let  $x, y \in Y$  such that  $x \leq_2 y$ . Then  $x^c \leq_2 y^c, \forall e \in E$ .

Since  $\phi_\psi$  is bijective, then there exist  $a^a$  and  $b^a$  in  $X_E$  such that  $\phi_\psi(a^a) = x^c$  and  $\phi_\psi(b^a) = y^c$  and since  $\phi_\psi$  is ordered embedding, then  $a^a \leq_1 b^a$ . So  $a \leq_1 b$ . By hypothesis, there exist disjoint pairwise soft neighbourhoods  $W_E$  and  $V_E$  of  $a$  and  $b$ , respectively, such that  $W_E$  is increasing and  $V_E$  is decreasing. Since  $\phi_\psi$  is bijective soft open, then  $\phi_\psi(W_E)$  and  $\phi_\psi(V_E)$  are disjoint soft neighbourhoods of  $x$  and  $y$ , respectively. It follows by Theorem 2.2, that  $\phi_\psi(W_E)$  is increasing and  $\phi_\psi(V_E)$  is decreasing. This completes the proof.

**Theorem 3.6** The property of being a TP\*(PP\*)-soft  $T_3$ -ordered space is soft bitopological ordered property.

**Proof.** The proof is similar to the previous theorem.

**Theorem 3.7** The property of being a TP-soft  $T_4$ -ordered space is soft bitopological ordered property.

**Proof.** Suppose that  $\phi_\psi$  is an ordered embedding soft homeomorphism map of a soft pairwise normally ordered space  $(X, \tau_1, \tau_2, E, \leq_1)$  on to an SBTOS  $(Y, \eta_1, \eta_2, K, \leq_2)$  and let  $H_E$  and  $F_E$  be two disjoint pairwise closed soft sets such that  $H_E$  is increasing and

$F_E$  is decreasing. Since  $\phi_\psi$  is bijective soft continuous, then  $\phi_\psi^{-1}(H_E)$  and  $\phi_\psi^{-1}(F_E)$  are disjoint pairwise closed soft sets and since  $\phi_\psi$  is ordered embedding, then  $\phi_\psi^{-1}(H_E)$  is increasing and  $\phi_\psi^{-1}(F_E)$  is decreasing. By hypothesis, there exist disjoint pairwise soft neighbourhoods  $W_E$  and  $V_E$  of  $\phi_\psi^{-1}(H_E)$  and  $\phi_\psi^{-1}(F_E)$ , respectively, such that  $W_E$  is increasing and  $V_E$  is decreasing. So  $H_E \vee \phi_\psi(W_E)$  and  $F_E \vee \phi_\psi(V_E)$ . The disjointness of the soft neighbourhoods  $\phi_\psi(W_E)$  and  $\phi_\psi(V_E)$  completes the proof.

#### 4. Conclusion

This research introduces a new concept called "Bi-ordered soft regularity and normality" and investigates its properties. The study considers the notions of belong, non-belong, partial belong, and total non-belong in order to understand their interrelationships. Several examples are presented to facilitate comprehension.

In future research, the goal is to explore novel bi-ordered soft separation axioms by leveraging these concepts on supra soft topological spaces. The hope is that this work will inspire further research and lead to advancements in the field of soft topology.

#### References

- [1] H. Aktas and N. Cagman, Soft sets and soft groups, Inf. Sci., 177(13) (2007) 2726 - 2735
- [2] M. I. Ali, F. Feng, X. Liu, W. K. Min, M. Shabir, On some new operations in soft set theory, Comput. Math. Appl. 57 (2009) 1547-1553.
- [3] T. M. Al-shami, M. E. El-Shafei, and M. Abo-Elhamayel. On soft topological ordered spaces. Journal of King Saud University-Science, 31(4) (2019) 556-566.
- [4] T. M. Al-shami, M. E. El-Shafei and M. Abo-Elhamayel, Partial soft separation axioms and soft compact spaces, Filomat, 32 (13) (2018) 4755-4771.

- [5] S. A. El-Sheikh, S. A. Kandil and S. Hussien, a new approach to soft bitopological ordered spaces, submitted.
- [6] S. A. El-Sheikh, S. A. Kandil and S. Hussien, separation axioms in soft bitopological ordered spaces, submitted.
- [7] S. Hussain and B. Ahmed, Some properties of soft topological spaces, *Comput. Math. Appl.*, 62 (2011) 4058-4067.
- [8] S. Hussain and B. Ahmad, Soft separation axioms in soft topological spaces, *Hacet. J. Math. Stat.*, 44(3) (2015) 559 - 568.
- [9] B. M. Ittanagi, Soft bitopological spaces, *International Journal of Computer Applications* 107(7) (2014) 1-4.
- [10] O. Gocur and A. Kopuzlu, Some new properties of soft separation axioms, *Ann. Fuzzy Math. Inform.*, 9(3) (2015) 421 - 429.
- [11] A. Kandil, O. A. E. Tantawy, S. A. El-Sheikh and Shawqi. A. Hazza, Pairwise open (closed) soft sets in soft bitopological spaces, *Ann. Fuzzy Math. Inform.*, 11 (4) (2016) 571-588.
- [12] J. C. Kelly 1975. *General Topology*. Springer Verlag.
- [13] McCartan, S.D. Separation axioms for topological ordered spaces. *Math. Proc. Cambridge Philos*, 1968. *Soc.* 64, 965 - 973.
- [14] P.K. Maji, A.R. Roy and R. Biswas, An application of soft sets in a decision making problems, *Comput. Math. Appl.*, 44 (2002) 1077 - 1083.
- [15] P. K. Maji, R. Biswas and A. R. Roy, Soft set theory, *Comput. Math. Appl.*, 45 (2003) 555-562.
- [16] D. A. Molodtsov, Soft set theory-firs tresults, *Comput. Math. Appl.*, 37 (1999) 19-31.
- [17] Nachbin, L., *Topology and ordered*. D. Van Nostrand Inc., Princeton, New Jersey 1965.
- [18] S. Nazmul and S. K. Samanta, Neighborhood properties of soft topological spaces, *Ann. Fuzzy Math. Inform.* 6 (1) (2013) 1-15.
- [19] S. Nazmul and S. K. Samanta, Some properties of soft topologies and group soft topologies, *Ann. Fuzzy Math. Inform.* 8 (4) (2014) 645-661.
- [20] M. Shabir and M. Naz, On soft topological spaces, *Comput. Math. Appl.*, 61 (2011) 1786-1799.
- [21] A. Singh and N.S. Noorie, Remarks on soft axioms, *Ann. Fuzzy Math. Inform.*, 14(5) (2017) 503 - 513.
- [22] O. Tantawy, S.A. El-Sheikh and S. Hamde, Seperation axioms on soft topological spaces, *Ann. Fuzzy Math. Inform.*, 11(4) (2016) 511 - 525.
- [23] I. Zorlutuna, M. Akdag, W. K. Min and S. Atmaca, Remarks on soft topological spaces, *Ann. Fuzzy Math. Inform.*, 3 (2) (2012) 171-185.