

**REVIEW**

# Assessing Individual Muscle Characteristics to Enhance Frozen-Thawed Meat Quality

 Choeun Im<sup>1</sup>, Sumin Song<sup>1</sup>, Huilin Cheng<sup>1</sup>, Junyoung Park<sup>1,2</sup>, and Gap-Don Kim<sup>1,3,\*</sup>
<sup>1</sup>Graduate School of International Agricultural Technology, Seoul National University, Pyeongchang 25354, Korea

<sup>2</sup>Mgenic Bio, Anseong 17529, Korea

<sup>3</sup>Institutes of Green Bio Science & Technology, Seoul National University, Pyeongchang 25354, Korea

 OPEN ACCESS

**Received** March 26, 2024

**Revised** May 6, 2024

**Accepted** May 7, 2024

\*Corresponding author : Gap-Don Kim  
 Graduate School of International Agricultural Technology, Seoul National University, Pyeongchang 25354, Korea  
 Tel: +82-33-339-5778  
 Fax: +82-33-339-5779  
 E-mail: gapdonkim@snu.ac.kr

\*ORCID  
 Choeun Im  
<https://orcid.org/0000-0003-3564-7069>  
 Sumin Song  
<https://orcid.org/0000-0001-7115-2253>  
 Huilin Cheng  
<https://orcid.org/0000-0003-0628-3358>  
 Junyoung Park  
<https://orcid.org/0000-0003-2569-6422>  
 Gap-Don Kim  
<https://orcid.org/0000-0001-5870-8990>

**Abstract** This study assessed previous research aimed at mitigating the adverse effects of freeze-thawing on meat quality. Specifically, it focuses on assessing the physicochemical alterations in meat resulting from freezing, freeze-thawing, or technologies to minimize these alterations. Recent studies have focused on conventional freeze-thaw technology applicable across various livestock species and muscle types. However, recent research has indicated the necessity for developing freeze-thaw technology considering the unique characteristics of individual muscles. In this review, we summarize previous studies that have compared alterations in the physicochemical properties of primary muscles owing to freezing or freeze-thawing. Despite the introduction of various technologies to significantly reduce the adverse effects on meat quality resulting from freeze-thawing, it is essential to consider the unique characteristics (proximate composition, pH, and muscle fiber characteristics) of individual muscles or cuts to develop enhanced the freeze-thaw processing technology.

**Keywords** freeze-thaw technology, meat quality, muscle type, muscle characteristics

## Introduction

Freezing meat enables long-term storage and distribution, inhibits the growth of microbes, and minimizes alterations in quality (Leygonie et al., 2012). However, thawing frozen meat leads to adverse alterations in quality, such as excessive exudation of meat juice, discoloration, and accelerated oxidation of proteins and lipids (Kim et al., 2013; Zhang et al., 2017).

Persistent academic and industrial initiatives aim to develop technologies to mitigate adverse alterations in meat quality by minimizing physicochemical changes during the freeze-thaw process (Zhang et al., 2023). Various physical and chemical phenomena that occur in meat during freezing or freeze-thaw processes have been observed, and their effects on meat quality characteristics have been assessed (Leygonie et al., 2012).

Additionally, methods such as controlling freezing and thawing speed, brine injection, vitamin E treatment, high-pressure, ohmic and electrostatic field treatments, and aging technology have been introduced to enhance frozen-thawed meat quality (Cevik and Icier, 2020; He et al., 2014; Hou et al., 2020; Kim et al., 2018; Zhu et al., 2004). Numerous studies have summarized and reviewed the technologies developed over the last two decades (Zhang et al., 2023). Most of these studies have focused on freeze-thaw technology, a widely applied method, regardless of the livestock species and muscle type. However, meat exhibits characteristics (proximate components, pH, and muscle fiber composition) unique to each muscle and displays varying physicochemical characteristics before freeze-thawing (Park et al., 2022). Therefore, these inherent variations in characteristics are anticipated to change depending on the muscle used during meat processing, including freezing and thawing.


However, previous studies have not adequately assessed the variations in muscle or meat cut characteristic-based freezing susceptibility and its influencing factors. Therefore, this review summarized the research results regarding the physical and chemical alterations in significant muscles owing to freezing or freeze-thawing. Moreover, we emphasize the necessity of developing technology to minimize adverse alterations in meat quality owing to freeze-thawing, depending on the characteristics of each muscle. To achieve this, we assessed previous studies related to the freezing and freeze-thawing of meat, specifically focusing on the results of muscles or comparative studies between various muscles or cuts.

## Formation of Ice Crystals through Freezing and Its Effect on Muscle Tissues

Freezing meat effectively prevents spoilage by inhibiting the growth of microorganisms (Coombs et al., 2017). However, thawing meat following freezing gradually eliminates its latent heat, resulting in altered physicochemical attributes, including reduced juiciness and water-holding capacity (WHC), discoloration, increased rancidity, and texture alterations (Cheng et al., 2020; Cheng et al., 2021; Park et al., 2012). This is because of the effects of ice crystals that form between and within myofibrils during the freezing process (Dang et al., 2021; Schudel et al., 2021). Meat contains approximately 75% water depending on the species, muscle type, and fat content (Huff-Lonergan and Lonergan, 2005). Among the different water types (bound, entrapped, and free) within meat, the entrapped and free water, crucial for chemical and biochemical reactions in meat, are susceptible to freezing due to ice crystal formation; however, bound water remains non-frozen in meat. Approximately 88% of the total water content in meat is freezable (Xanthakis et al., 2013).

Ice crystal formation in meat typically begins at approximately  $-1^{\circ}\text{C}$ , with approximately 75% of the moisture of the forming ice crystals at  $-5^{\circ}\text{C}$  (Cooke and Wien, 1971; Huff-Lonergan and Lonergan, 2005). When the core temperature of the meat drops to  $-12^{\circ}\text{C}$ , approximately 92% of the moisture forms ice crystals, and the residual moisture maintains the ingredients in the meat, including proteins, in an unfrozen state (Cooke and Wien, 1971; Huff-Lonergan and Lonergan, 2005). While meat is frozen, water initially creates ice crystal nuclei both inside and outside the muscle fibers (myocytes), which then gradually expand in the form of branches, producing large or small crystals (Xanthakis et al., 2013). These ice crystals can physically damage muscle microstructure, resulting in Z-line destruction, I-band weakening, and increased intermyofibrillar space (Añón and Calvelo, 1980; Cheng et al., 2020). The degree of physical destruction within the muscle varies depending on the size of ice crystals (Dang et al., 2021; Schudel et al., 2021).

The freezing rate of meat determines the size and number of ice crystals, which affect the muscle fiber structure (Fig. 1). Rapid freezing leads to the formation of smaller ice crystals outside muscle fibers, whereas slow freezing leads to the formation of larger ice crystals (Cheftel et al., 2002; Fernández et al., 2006; Su et al., 2014). The freezing rate is linked to the



**Fig. 1.** Schematic diagram of the physical destruction of meat tissue due to differences in freezing speed. Rapid freezing creates relatively small ice crystals within muscle, resulting in less tissue destruction, while slow freezing creates large ice crystals within muscle, resulting in severe tissue destruction.

heat transfer rate from the outside to core of the meat and is closely related to the uniformity and non-uniformity of the histological and physicochemical properties of frozen or frozen-thawed meat (Choi et al., 2016; Sanz et al., 1999). Drip loss inevitably occurs when the frozen meat is thawed. During thawing, ice crystal formation varies based on the freezing speed of meat, which determines the degree of muscle tissue destruction and drip loss (Dang et al., 2021; Schudel et al., 2021). Additionally, the formation of ice crystals alters the solute concentration both within and outside the muscle fiber, resulting in an imbalance, thereby leading to biochemical and physical alterations in multiple meat components (Bevilacqua et al., 1979; Huff-Lonergan and Lonergan, 2005; Love and Haraldsson, 1961). Therefore, ice crystal formation and physicochemical alterations within the muscle tissue owing to freezing are significant factors that directly or indirectly affect muscle food quality characteristics such as WHC, meat color, flavor, and texture (Cheng et al., 2020; Cheng et al., 2021; Dang et al., 2021; Wang et al., 2020).

## Effects of Freezing and Freeze-Thawing on Meat Physicochemical Properties

Freezing is an effective method to extend the shelf life of meat. However, the freezing and thawing processes have adverse effects on meat quality characteristics, such as discoloration, reduction in WHC, and alterations in sensory properties (flavor, taste, and texture; Dang et al., 2021; Wang et al., 2020). A detailed overview of the alteration in color, WHC, texture, and shelf life of meat owing to freezing and freeze-thaw cycles is described below.

### Water-holding capacity

The shelf life of meat is usually determined by its appearance, color, flavor, nutritional value, and microbial activity. The

various ingredients of meat are crucial factors that affect its quality characteristics and alterations (McMillin, 2008). The moisture of meat is primarily retained in the interstices between the thick and thin myofibril filaments. This occurs within and between myofibrils, and a small amount of water within the muscle is retained through the electrostatic attraction between proteins (Bond et al., 2004; Cheng and Sun, 2008). Additionally, the fat present in meat contains water, which contributes to maintaining its moisture content (Joo et al., 2002). When meat releases moisture naturally or owing to certain factors (cooking, freeze-thawing, and pressing), the exudate contains nutrients and flavor components, such as vitamins, minerals, and amino acids (Añón and Calvelo, 1980; Leygonie et al., 2012; Ngapo et al., 1999). Therefore, WHC (the ability of meat to retain moisture) is synonymous with the effectiveness of retaining various useful ingredients in meat. Additionally, WHC is commonly assessed by measuring the degree of drip loss, thawing, cooking, and purging (Honikel, 1998; Honikel and Hamm, 1994; Huff-Lonergan and Lonergan, 2005).

Freezing significantly reduces the WHC of meat. Thawing of the frozen meat results in excessive exudation of its juices (Leygonie et al., 2012). Therefore, basic and applied studies were conducted to minimize the amount of meat exudate, as listed in Table 1. The primary results of previous studies on the loss of exudates and alterations in WHC in meat based on freezing or freeze-thaw processes are summarized below.

Regardless of the animal species, increasing the freezing rate can reduce the size of ice crystals and the thawing loss (Kim et al., 2020; Yun et al., 2021). Repeated freeze-thaw cycles result in a significant deterioration of meat quality; additionally, a substantial alteration in freezing temperature results in unfavorable WHC compared to a minor temperature alteration (Wang et al., 2020). The WHC varies based on the order of aging (cold storage) and freezing. When aged for approximately three weeks following freeze-thawing, more exudates were lost compared to frozen-thawed following aging (Kim et al., 2018). In studies using various cuts (muscles) of pork and beef, the loss of exudates owing to freeze-thaw may vary based on the muscle type. Cooking loss of pork loin (*M. longissimus thoracis et lumborum*) increased significantly following freeze thawing compared to that of other muscles (*M. psoas major*, *M. semimembranosus*, and *M. semitendinosus*). However, in beef, tenderloin (*M. psoas major*) exhibited significant WHC following freeze thawing compared to other muscles (*M. longissimus lumborum*, *M. semimembranosus*, and *M. semitendinosus*; Cheng et al., 2020; Cheng et al., 2021). Additionally, it has been confirmed that cooking loss in beef significantly increases when the freezing period increases, regardless of muscle type (Cho et al., 2017). Freezing solutions consisting of sodium chloride, ethyl alcohol, and chitosan reduce the size of ice crystals and the thawing-related loss in pork (Hou et al., 2020).

During thawing, meat undergoes protein breakdown, lipid oxidation, color changes, and ice crystal melting, reducing its WHC (Gan et al., 2022; Min et al., 2016). Poudyal et al. (2023) observed that shorter thawing times increased drip loss in porcine *M. semimembranosus*, whereas longer thawing times enhanced the reabsorption of free water from ice crystals, thereby reducing the final moisture loss. However, shorter thawing times may result in inadequate water reabsorption, resulting in excessive water loss (Gonzalez-Sanguinetti et al., 1985). Min et al. (2016) observed that thawing meat using pressure ohmic thawing of up to 200 MPa with an electric field strength of 40 V/cm and pressure-assisted thawing of up to 200 MPa reduced drip loss in beef *psoas major* compared to conventional thawing methods.

### **Meat color**

Consumers prioritize price and color when purchasing meat (Carpenter et al., 2001). However, meat discoloration occurs during storage or retail displays. Based on previous research, 15%–20% of discoloration occurs during retail displays, resulting in an economic loss of 50% or more (Hood and Riordan, 1973; Mancini and Hunt, 2005). The primary factors

**Table 1. Previous studies on changes in physicochemical properties of meat due to freeze-thawing**

Species	Muscles	Freezing/thawing/storage conditions	Outcome	References
Porcine	<i>M. longissimus dorsi</i> (LTL)	Sequence of freezing and aging: Freezing without aging (FT), aging prior to freezing (AFT), and freezing and aging (FTA) Slow-freezing: -20°C for 8 weeks and thawing 1°C for 2 days Fast-freezing: -80 °C and stored -20°C for 8 weeks and thawing 1°C for 2 days Aging: 1°C for 19 days	pH: FTA≤FT<AFT ↓ WBSF in FTA with slow-freezing ↑ Purge/thaw loss in FTA unrelated with freeze rate ↑ CIE L* in FTA with slow-freezing ↓ CIE a* in slow-freezing ↓ CIE b* in FTA with slow-freezing	Kim et al. (2018)
Porcine	<i>M. longissimus dorsi</i> <i>M. psoas major</i> (PM) <i>M. semimembranosus</i> (SM) <i>M. semitendinosus</i> (ST)	Fresh: 4°C for 7 days Freeze-thawing: -20°C for 5 days and thawing at 4°C for 2 days	- pH in all muscle cut and freeze- thawing - Purge loss ↓ CIE L* in LTL and ST ↑ CIE a* in LTL and SM ↑ CIE b* in LTL and ↓ in PM and ST ↑ WBSF in SM and ST ↑ Cooking loss in LTL ↓ Drip loss in SM and ST ↓ WHC and toughness in SM and ST ↑ Muscle type I and IIA size in ST ↑ Muscle type IIXB amount in SM	Cheng et al. (2021)
Bovine	<i>M. longissimus thoracis et lumborum</i>	Fresh: Aging at 4°C for 2, 7, and 14 days Freeze-thawing: Aging for 2, 7, and 14 days at 4°C, freezing at -20°C for 2 months, and overnight thawing at 4°C	↓ WBSF over time ↓ WHC in freeze-thawing ↑ Tenderness, juiciness and meat taste in fresh - Panel preference	Lagerstedt et al. (2008)
Bovine	<i>M. longissimus thoracis et lumborum</i> (LT) <i>M. psoas major</i> <i>M. semimembranosus</i> <i>M. semitendinosus</i>	Fresh: 4°C for 5 days Freeze-thawing: -20°C for 3 days and thawing at 2°C for 2 days	- pH in all muscle cut and freeze-thawing ↑ CIE L* in LT with freeze-thawing ↓ CIE a* in LT, ↑ CIE a* in SM and ST with freeze-thawing ↓ CIE b* in SM with freeze-thawing ↓ WBSF in LT with freeze-thawing ↑ Purge losses in LT, SM and ST with freeze-thawing ↓ Drip loss in all cuts except PM with freeze-thawing ↓ Cooking loss in LT and PM, ↑ cooking loss in SM with freeze- thawing ↑ WHC and ↓ discoloration in PM than SM ↑ Muscle type I amount in PM than SM	Cheng et al. (2020)
Porcine Chicken	Porcine neck Chicken leg	Freeze-thawing: Frozen at -18°C, -50°C, and -60°C for 0.5, 1, 2, 3, 4, 5, and 6 months and then thawed at 2°C	↓ Thawing loss at -60°C, ↑ thawing loss at -18°C during 6 months in porcine neck and chicken leg with freeze-thawing - Cooking loss in pork neck and chicken leg with different temperature ↑ CIE L*, CIE a*, and CIE b* in pork neck frozen at -18°C	Kim et al. (2020)

**Table 1. Previous studies on changes in physicochemical properties of meat due to freeze-thawing (continued)**

Species	Muscles	Freezing/thawing/storage conditions	Outcome	References
Bovine	<i>M. longissimus thoracis et lumborum</i>	A4: Ageing at $-1.5^{\circ}\text{C}$ for 4 weeks A4F2: Ageing at $-1.5^{\circ}\text{C}$ for 4 weeks and freezing at $-18^{\circ}\text{C}$ for 2 weeks with fast freezing in freezing solution or slow freezing in air-blast at $-18^{\circ}\text{C}$ A3F2: Ageing at $-1.5^{\circ}\text{C}$ for 3 weeks and freezing at $-18^{\circ}\text{C}$ for 2 weeks with fast freezing in freezing solution or slow freezing in air-blast at $-18^{\circ}\text{C}$ F2: Fast freezing in freezing solution or slow freezing in air-blast at $-18^{\circ}\text{C}$ for 2 weeks	- pH, cooking loss, drip loss, purge loss and WBSF by freeze rate pH: A4F2>F2>A4>A3F2 Purge loss and drip loss: F2>A3F2>A4F2>A4 Cooking loss: A4>A4F2>A3F2>F2 WBSF: F2>A4>A3F2=A4F2 CIE L*, CIE a*, and C: A4>A4F2, A3F2, and F2 Hue angle: A4<A4F2, A3F2, and F2	Kim et al. (2015)
Bovine	Hind legs	A: Freezing at $-18^{\circ}\text{C}$ with freezing fluctuation cycles repeat for 3 days B: Freezing at $-18^{\circ}\text{C}$ and $-17^{\circ}\text{C}$ ; stored at $-18^{\circ}\text{C}$ for 4 hours, raised to $-17^{\circ}\text{C}$ , then returned to $-18^{\circ}\text{C}$ for another 4 hours with freezing fluctuation cycles repeat for 3 days C: Freezing at $-18^{\circ}\text{C}$ and $-15^{\circ}\text{C}$ ; stored at $-18^{\circ}\text{C}$ for 4 hours, raised to $-15^{\circ}\text{C}$ , then returned to $-18^{\circ}\text{C}$ for another 4 hours with freezing fluctuation cycles repeat for 3 days D: Cycled between $-18^{\circ}\text{C}$ and $-13^{\circ}\text{C}$ ; stored at $-18^{\circ}\text{C}$ for 4 hours, raised to $-13^{\circ}\text{C}$ , then returned to $-18^{\circ}\text{C}$ for another 4 hours with freezing fluctuation cycles repeat for 3 days	Free drip loss and cooking loss: A<B<C<D Centrifuge drip loss: A=B<C<D CIE L*: A>C>B>D CIE a*: A>B>C>D CIE b*: A<B<C<D	Wang et al. (2020)
Bovine	<i>M. longissimus thoracis</i> <i>M. longissimus lumborum</i> (LL) <i>M. psoas major</i> <i>M. semimembranosus</i>	Fresh: Aging at $2^{\circ}\text{C}$ for 0, 7, and 14 days Freeze-thawing: Frozen at $-18^{\circ}\text{C}$ for 0, 3, 6, and 9 months	↓ CIE L* in PM with increase of freezing period - CIE a* and CIE b* by aging and freezing period ↑ WHC in LL, PM, and SM by freezing for 3 months and in LT by freezing for 3 and 6 months ↑ Cooking loss in LL, LT, PM, and SM with an increase of freezing period ↓ WBSF in LL, LT, and SM with aging period	Cho et al. (2017)
Porcine	<i>M. longissimus dorsi</i>	Freeze-thawing: $-20^{\circ}\text{C}$ , $-30^{\circ}\text{C}$ , and $-40^{\circ}\text{C}$ in airflow velocities of 0, 1.5, and 3 m/s and thawing at $2^{\circ}\text{C}$ over night Ice crystal size Category I: Slow freezing in $<0.4$ cm/h Category II: Intermediate freezing in 0.6–0.7 cm/h Category III: Rapid in $>0.96$ cm/h	Ice crystal size: Slow freezing ( $<0.5$ cm/h)>rapid freezing ( $>0.5$ cm/h) ↓ Thawing loss in category III>II>I ↓ Drip loss and cooking loss in category III>II>I ↓ CIE a* and CIE b* in category I ↓ WBSF in category II and III than at $-20^{\circ}\text{C}$ with no air	Yun et al. (2021)

**Table 1. Previous studies on changes in physicochemical properties of meat due to freeze-thawing (continued)**

Species	Muscles	Freezing/thawing/storage conditions	Outcome	References
Porcine	<i>M. longissimus dorsi</i>	ISF: Fast freezing in freezing solution at $-22^{\circ}\text{C}$ consisting of 16% sodium chloride, 25% ethyl alcohol, 1.2% chitosan, 0.8% antifreeze protein, and 57% water and thawing at 1, 31, 61 and 91 days at $4^{\circ}\text{C}$ for 12 h AF: Slow freezing in air-blast at $-22^{\circ}\text{C}$ and thawing at 1, 31, 61 and 91 days at $4^{\circ}\text{C}$ for 12 h	↓ Ice crystal formation in ISF ↓ WBSF in AF ↑ Myofibrillar damage and lipid oxidation in AF - CIE L*, CIE a*, and CIE b* in ISF and AF with thawing for 91 days ↑ pH in AF at 91 days, - pH in ISF ↑ Thawing loss in AF for 1, 31, and 91 days ≐ Cooking loss in AF and ISF	Hou et al. (2020)
Bovine	<i>M. longissimus thoracis</i>	Fresh: Aged at $2^{\circ}\text{C}$ for 2, 3, 4, 5, 6, 7, 10, 14, 21, and 35 days Freeze-thawing: Aged for 3, 4, 5, 6, 7, 10, 14, 21, and 35 days at $2^{\circ}\text{C}$ , frozen at $-16^{\circ}\text{C}$ for 2 months and thawed at $1^{\circ}\text{C}$ for 24 h	Cooking loss: Fresh<frozen-thawed for 1, 3, 4, 5, 6, 7, and 10 days WBSF: Fresh>frozen-thawed for all days ≐ WBSF for 3, 6, and 7 days aged-frozen meat with aged for 14 to 21 days fresh meat	Shanks et al. (2002)
Bovine	<i>M. longissimus thoracis et lumborum</i>	Pelvic suspension: Carcasses hung in pelvic bone Achilles suspension: Carcasses hung in Achilles tendon Both suspensions treated at $4^{\circ}\text{C}$ , aged for 2, 4, 7, 14, and 21 days and frozen at $-20^{\circ}\text{C}$ for 7 days, and thawed at $4^{\circ}\text{C}$ overnight	- Tenderness in both suspensions ↓ WBSF in pelvic suspension for 2 days ageing ≐ WBSF between suspensions after ageing for 14 and 21 days ≐ pH in both groups during ageing ↓ WHC in Achilles suspension	Enfält et al. (2004)
Bovine	<i>M. longissimus thoracis</i>	Fresh: Aged at $4^{\circ}\text{C}$ for 3 and 10 days Freeze-thawing: Aged at $4^{\circ}\text{C}$ for 3 and 10 days and frozen at $-20^{\circ}\text{C}$ and $-80^{\circ}\text{C}$ for 30, 75, and 90 days, and thawed at $4^{\circ}\text{C}$ for 48 h	↓ CIE L*, CIE a*, and chroma in fresh aged for 3 and 10 days ↓ CIE a*, CIE b*, and chroma, and ↑ hue in freeze-thawing for 90 days ↑ Freezing loss and press loss in fresh aged for 3 and 10 days - Cooking loss in freeze-thawing for storage periods ↑ Press loss in freeze-thawing for 90 days ↑ Cooking loss in freeze-thawing for 30, 60 and 90 days ↓ WBSF in fresh aged for 3 days and freeze-thawing for 75 and 90 days - WBSF in Fresh aged for 10 days ↑ Tenderness in freeze-thawing aged for 3 days and frozen for 30, 75 and 90 days	Vieira et al. (2009)
Porcine	<i>M. psoas major</i>	Freezing Fresh: Stored in $4^{\circ}\text{C}$ for further analysis SR: $-10^{\circ}\text{C}$ supercooled rapid freezing in static air freezer at $-80^{\circ}\text{C}$ NR: Non-supercooled rapid freezing at $-80^{\circ}\text{C}$ SS: $-10^{\circ}\text{C}$ supercooled slow freezing at $-20^{\circ}\text{C}$ NS: Non-supercooled slow freezing at $-20^{\circ}\text{C}$ Thawing RT: Room temperature thawing in $20^{\circ}\text{C}$ IW: Ice water thawing at $0\pm 0.5^{\circ}\text{C}$ RF: Refrigerator thawing at $4^{\circ}\text{C}$ TW: Tap water thawing under $12\pm 2^{\circ}\text{C}$ running water	- Hardness in all freeze-thawing condition Drip loss: IW and RF<TW and RT ↑ Drip loss in NS ↓ Size ice crystal in SR	Poudyal et al. (2023)

**Table 1. Previous studies on changes in physicochemical properties of meat due to freeze-thawing (continued)**

Species	Muscles	Freezing/thawing/storage conditions	Outcome	References
Bovine	<i>M. psoas major</i>	Freezing Fresh: Stored at 4°C for 2 days Freezing: Frozen at -40°C for overnight Thawing POT: Pressure ohmic thawing up to 200 MPa and electric field strength of 40 V/cm PAT: Pressure-assisted thawing up to 200 MPa Ohmic thawing: Ohmically thawed to 8°C under electric field of 40 V/cm to 4°C at 0.1 MPa Conventional thawing: Thawed to 4°C by immersion thawing	↓ Drip loss in PAT, POT and ohmic thawing Cooking loss: POT<PAT<ohmic thawing<conventional thawing ↑ WBSF in PAT - CIE L* in all condition ↓ CIE a* and CIE b* in PAT, ohmic thawing and conventional thawing	Min et al. (2016)
Bovine Porcine Lamb	<i>M. psoas major</i>	Freezing Fresh: Refrigeration temperature Freezing: At -80°C for 2 days Thawing MT: Microwave thawing UT: Ultrasonic thawing IT: Infrared thawing at 60°C and the power of the infrared tube to 12 W RTT: Thawed at room temperature SWT: Thawed in still water at 25°C	↓ Thawing loss and cooking loss in UT ↑ Cooking loss in RTT ↑ CIE L* in RRT ↓ CIE L* in UT ↑ CIE b* in UT ↓ Lipid oxidation in UT	Gan et al. (2022)

≈, approximately equal; -, no significantly different; ↑, increase; ↓, decrease; >, higher; <, lower; =, equal; ≥, higher or equal. WBSF, Warner-Bratzler shear force; WHC, water-holding capacity.

influencing meat color are the myoglobin content and chemical state (Suman and Joseph, 2013). Myoglobin alters meat color based on redox phenomena (King et al., 2023). In its unoxidized state containing iron molecules, myoglobin exists as deoxy-myoglobin (deoxyMb), and the meat color appears purple. When deoxyMb reacts with oxygen in the air, known as blooming, it transforms into oxy-myoglobin (oxyMb), which displays a bright cherry-red color. When deoxyMb and oxyMb lose electrons or both electrons and oxygen, they are converted into oxidized myoglobin (metmyoglobin; metMb), resulting in meat with a brownish tint.

Therefore, meat color changes based on the chemical state (oxidation or reduction states, bonding with oxygen or water molecules) of the heme iron molecule in myoglobin. Additionally, the role of oxidation-reduction enzymes in altering the chemical state is crucial (Tang et al., 2005a; Tang et al., 2005b; Tang et al., 2005c). External factors that affect the oxidation-reduction states of myoglobin, such as packaging conditions (aerobic vs. anaerobic), storage temperature (refrigerated vs. frozen), and storage period, directly alter the oxidation, reduction, and oxygenation states (King et al., 2023). Moreover, it also affects the activity of enzymes involved in the oxidation-reduction of myoglobin (Tang et al., 2005a; Tang et al., 2005b; Tang et al., 2005c). Freeze thawing is one of the factors responsible for these intricate alterations in meat color (Jeong et al., 2011).

As demonstrated in Table 1, a slow freezing speed correlated with an increase in the CIE L\* of the meat. Moreover, with an extended frozen storage period, the CIE L\* of beef tenderloin reduces (Cho et al., 2017; Hou et al., 2020; Kim et al., 2020). However, the CIE a\* and CIE b\* of pork and beef do not vary during the freezing period (Cho et al., 2017; Hou et al., 2020). Additionally, repeated freeze-thaw with significant temperature alteration results in a reduction in the CIE L\* and CIE a\* of beef, whereas the CIE b\* increases (Wang et al., 2020).



Meanwhile, during cold storage following freeze-thaw, a slower freezing speed results in an increase in the CIE L\* of pork loin and a reduction in CIE b\* (Kim et al., 2018). Studies on the effects of freeze-thaw on pork and beef muscles have shown contrasting results for the loin muscles of the two species. The CIE L\* of beef was reduced and the CIE a\* increased following freeze-thawing, whereas CIE a\* increased following freeze-thawing (Cheng et al., 2020; Cheng et al., 2021). However, the CIE L\* of pork loin tends to be increased and CIE a\* is reduced following freeze-thawing (Cheng et al., 2021). In contrast, *semimembranosus* exhibited an increase in CIE a\* owing to freeze-thawing, regardless of the species. Moreover, *psaos major* exhibited a tendency to reduce CIE b\* owing to freeze-thawing of beef. However, *psaos major* exhibited relatively smaller alterations in meat color owing to freeze-thawing in both species compared to other muscles, such as *M. longissimus thoracis et lumborum*, *M. semimembranosus*, and *M. semitendinosus* (Cheng et al., 2020; Cheng et al., 2021). Furthermore, the meat color changes during thawing. Min et al. (2016) demonstrated that thawing beef *psaos major* did not alter the CIE L\* of meat. However, the CIE a\* and CIE b\* reduced in all thawing treatments, such as pressure-assisted thawing up to 200 MPa, pressure ohmic thawing up to 200 MPa, electric field strength of 40 V/cm, and immersion thawing. Gan et al. (2022) observed that ultrasonic thawing for beef, pork, and lamb *M. psaos major* resulted in minimal alterations in meat color compared to other thawing methods, such as microwave, room temperature, and 25°C water thawing. This indicates that ultrasonic thawing is significantly effective in preserving muscle quality and reducing myoglobin and lipid oxidation in meat.

## Tenderness

Meat tenderness is a crucial sensory quality, and its variations are caused by alterations in the chemical composition and structural properties of muscle fibers and connective tissues, which are influenced by the animal species, breed, slaughter method, and postmortem processing of the meat (Pogorzelski et al., 2022; Zhang et al., 2023). Freezing affects meat tenderness with a positive correlation between freezing and tenderness (Lagerstedt et al., 2008). Ice crystals generated through freezing exert pressure on muscle tissues, resulting in physical destruction and protein decomposition (Añón and Calvelo, 1980; Cheng et al., 2020). When meat is frozen, larger ice crystals cause significant physical destruction of the muscle tissue and meat tenderness (Leygonie et al., 2012; Zhang et al., 2023). Thawing technique also affects meat tenderness. Min et al. (2016) demonstrated that the pressure ohmic thawing technique results in minimal texture alteration, with shear forces closely resembling those of fresh meat.

Aging the meat adequately before freezing negates the tenderizing effect of freezing (Vieira et al., 2009). Enfält et al. (2004) discovered that beef subjected to initial aging for seven days before freezing exhibited similar shear force to beef aged for 21 days in refrigeration. Shanks et al. (2002) observed that frozen-thawed *longissimus dorsi* exhibited a lower shear force than chilled meat over the same period. Freezing enhances tenderness (reduction in shear force); however, the degree of enhancement varies based on the freezing rate. Rapid freezing forms smaller ice crystals compared to slow freezing, resulting in relatively less enhancement of tenderness (Kim et al., 2015; Kim et al., 2018; Yun et al., 2021). Additionally, prolonged frozen storage enhances meat tenderness (Cho et al., 2017). Furthermore, distinct tenderness patterns following freeze-thawing were observed among various muscle types, with a reduction in the tenderness of porcine *M. semimembranosus* and *M. semitendinosus*, and an enhancement in the tenderness of bovine *M. longissimus lumborum* (Cheng et al., 2020; Cheng et al., 2021). Cheng et al. (2020) and Cheng et al. (2021) reported that varied characteristics of the muscle fibers in each muscle may contribute to the varying susceptibilities of muscle types owing to freezing. Additionally, freezing can physically damage muscle tissue and enhance tenderness but weakens the WHC of meat. This leads to excessive meat exudation upon

thawing, resulting in a dry or tough meat texture. Because excessive exudation of meat juice negatively affects tenderness, it can be enhanced or reduced through freeze-thawing based on the characteristics of each muscle. In order to reduce the negative changes in meat texture caused by the reduction in WHC, it is necessary to minimize the destruction of the tissue through rapid freezing and high pressure treatments (Choi et al., 2016; Min et al., 2016; Sanz et al., 1999).

### Microbial growth and oxidation in meat

Freezing is a microbiologically safer and long-term meat storage method compared to refrigeration. This is because microbial growth is inhibited at approximately  $-12^{\circ}\text{C}$ , cell metabolism in animal tissue is inhibited at approximately  $-18^{\circ}\text{C}$ , and alterations in meat quality are minimal at  $-55^{\circ}\text{C}$  (Dave and Ghaly, 2011; Hansen et al., 2004). Beef and lamb can be stored at  $-18^{\circ}\text{C}$  for 12 months, pork for six months, and poultry meat for 9–12 months (Valero Díaz et al., 2023). Freezing inhibits microbial growth and kills approximately 60% of the microbial population. However, surviving microbes proliferate upon thawing, and fat oxidation persists even during freezing (Dave and Ghaly, 2011; Zhou et al., 2010). Therefore, freezing does not ensure the safety of meat. Lipid oxidation and meat spoilage bacteria are significant factors that shorten the shelf life of meat. Meat products undergo oxidation during processing, storage, and light exposure. This process deteriorates the nutritional and organoleptic properties of meat and results in the formation of toxic compounds that are harmful to humans (Papuc et al., 2017).

Thawing results in the conversion of solid ice crystals to liquid water, which raises the surface temperature and can facilitate microbial reactivation by providing appropriate temperature and humidity conditions (Marriott et al., 1980). Additionally, juices exuded during freezing and thawing provide a nutritious environment and suitable medium for microbes (Leygonie et al., 2012). Rapid thawing techniques are crucial for minimizing microbial growth (Min et al., 2016) and reducing damage caused by thawing.

Bacteria significantly associated with meat spoilage include *Carnobacterium* spp., *Enterobacteriaceae*, *Lactobacillus* spp., *Leuconostoc* spp., *Shewanella putrefaciens*, *Pseudomonas* spp., and *Brochothrix thermosphacta* (Borch et al., 1996). *Pseudomonas* species are the primary cause of meat spoilage at temperatures between  $-1^{\circ}\text{C}$  and  $25^{\circ}\text{C}$  in air. *Pseudomonas* can penetrate meat because of its proteolytic activity (Gill and Penney, 1977; Gupta and Nagamohini, 1992). Therefore, bacterial spoilage of meat results in odors, off-flavors, discoloration, gas production, slime formation, and pH reduction, thereby causing significant economic losses (Papuc et al., 2017).

The primary factor that causes the deterioration of stored meat is lipid oxidation, which is caused by enzymes produced by microbes and meat enzymes or autoxidation (Zhang et al., 2010). Among microbes, certain isolates produce lipid-oxidizing enzymes, such as lipoxygenase, linoleate, oxygen oxidoreductase, and hydroperoxide lyase (Brash, 1999). Meat oxidation is a significant factor in the deterioration of meat quality, affecting flavor, texture, nutritional value, and color. Lipid oxidation in fresh meat, known as autoxidation, occurs during cold or frozen storage and is induced by oxygen and lipid free radicals (Soyer et al., 2010). Free radicals are atoms or molecules characterized by unpaired electrons that are inherently unstable and reactive in nature. When they attract electrons from stable compounds, they become free radicals, creating a continuous cycle (Fang et al., 2002; Škrovánková et al., 2012). The resulting hydroperoxides cause biological damage to lipids, hexanes, enzymes, and proteins, thereby resulting in the production of detrimental compounds, such as malondialdehyde and cholesterol oxidation products (Morrissey et al., 1998). Numerous factors influence lipid oxidation in meat, including heat, light, antioxidants, oxygen, phospholipid, and unsaturated fatty acid contents (Guyon et al., 2016). Gan et al. (2022) observed that freeze-thawing of psoas major from bovine, porcine, and lamb increased lipid oxidation in meat across all animal species.

However, ultrasonic thawing resulted in significant stable alterations. The degree of lipid oxidation increases with longer thawing times (Fioramonti et al., 2017; Gan et al., 2022). Microwave thawing has a shorter thawing time; however, because of the energy and heat generated, it is excessive and facilitates lipid oxidation (Gan et al., 2022; Lorentzen et al., 2020). Phospholipids present in cell membranes are highly sensitive to oxidation because of their higher unsaturation compared with other fats in meat (Domínguez et al., 2019). Lean meat contains a relatively high proportion of easily oxidized phospholipids. Therefore, phospholipids are the primary cause of fat oxidation in lean meats (Domínguez et al., 2019). Additionally, phospholipid fat oxidation results in noticeable alterations in the meat quality. Lipase-mediated phospholipid degradation has also been observed in frozen meat. These reactions occur at various freezing temperatures and storage periods (Pikul et al., 1985). Therefore, the degree to which packaging reduces air exposure during frozen storage plays a crucial role in determining fatty acid oxidation. Protein oxidation, assessed through carbonyl and sulfhydryl content, is significantly correlated with lipid oxidation (Mercier et al., 1998; Renner et al., 1999). Meat, specifically chicken, has a high unsaturated fatty acid content. Therefore, lipid and protein oxidation are more prevalent during frozen storage. Moreover, the degree of oxidation is highly significant in chicken legs than in chicken breasts (Soyer et al., 2010).

## Assessing Individual Muscle Characteristics to Enhance Meat Freezing/Thawing Technologies

The quality of meat subjected to freezing and thawing depends on pretreatment, freezing, storage, and thawing conditions. Freezing is the primary process that significantly influences the overall quality of frozen muscle food (Nakazawa and Okazaki, 2020). Rapid freezing and the formation of small ice crystals are crucial for meat freezing because they reduce muscle tissue damage and drip loss during thawing (Li and Sun, 2002). Kim et al. (2018) demonstrated that total exudate loss was reduced in pork loin muscle subjected to rapid freezing ( $-80^{\circ}\text{C}$ ) compared to slow freezing ( $-20^{\circ}\text{C}$ ). Moreover, aging before freezing and rapid freezing contributed to reduced deterioration. Additionally, rapid freezing reduces the purge and drip loss in beef loins (Kim et al., 2015). Kim et al. (2020) discovered that preserving pork neck and chicken leg meat at varying storage temperatures ( $-18^{\circ}\text{C}$ ,  $-50^{\circ}\text{C}$ , and  $-60^{\circ}\text{C}$ ) for six months revealed that maintaining the quality of fresh meat without any degradation for the complete duration was achieved through freezing at  $-60^{\circ}\text{C}$ . High-pressure freezing is anticipated to achieve significant supercooling effects that can form uniform ice crystals within meat products (Cheftel et al., 2002). Upon pressure release, an increase in pressure results in the transformation of type-I ice crystals (less dense than liquid water) to type-IV ice crystals (Cheftel et al., 2000). Type IV ice crystals, which are smaller and denser than water, exhibit no expansion upon freezing, in contrast with the 9%–13% typical expansion observed in type I crystals. They can produce high-quality freeze-thawed meat with minimal cell structure damage. However, high-pressure freezing is expensive and has limitations on meat size (Chevalier et al., 2000; Fernández et al., 2007). Additionally, static electric field freezing (Fallah-Joshaqani et al., 2019) and ultrasonic technology (Patist and Bates, 2008) have been proposed, with ultrasonic technology reducing the thawing time and structural damage to muscle fibers owing to freezing (Guo et al., 2021).

The degree of alteration in meat quality owing to freezing and freeze-thaw varies based on the moisture and fat content and muscle fiber characteristics (Huff-Lonergan and Lonergan, 2005; Song et al., 2020). Among the muscle fiber types, type I is less susceptible to freezing than type II. Therefore, muscles with a higher type I distribution exhibit less deterioration in meat quality when subjected to freezing and thawing (Cheng et al., 2020; Cheng et al., 2021; Song et al., 2020). Moreover, various cuts/muscles exhibit distinct muscle fiber characteristics, including muscle fiber composition, resulting in distinct freezing

susceptibilities. Variations in freezing susceptibility based on muscle fiber type have been confirmed in major cuts (loin or strip loin, tenderloin, round, and eye of round) of beef and pork (Cheng et al., 2020; Cheng et al., 2021). Beef tenderloin (*M. psoas major*) with a high proportion of type I muscle fibers exhibited a relatively lower degree of reduction in WHC and discoloration owing to freeze-thawing compared to round (*M. semimembranosus*) and eye of round (*M. semitendinosus*) with a high proportion of type II muscle fibers. This pattern was also observed in the WHC and tenderness of pork. Therefore, these results emphasize the significance of freeze-thaw that considers the components or muscle fiber characteristics of each muscle/cut, regardless of the animal species.

Thawing meat results in ice crystal melting, thereby reducing WHC. Therefore, it is crucial to use appropriate thawing techniques to enhance the quality of thawed meat (Alonso et al., 2016), prevent damage to the muscle structure, minimize fat and protein oxidation, and reduce water retention (Zhang et al., 2023). Generally, meat is thawed using air or water at refrigeration or room temperature, which provides a cost-effective approach without physical treatment (Gan et al., 2022; Lan et al., 2021). Advanced thawing technologies, including high-pressure, microwave, ohmic, ultrasonic, electrostatic field, and radiofrequency field and combinations of these techniques (Zhang et al., 2023), can effectively reduce the thawing time and minimize the deterioration of meat quality compared to traditional thawing techniques. Gan et al. (2022) reported that ultrasonic thawing, a physical technique, can effectively prevent muscle structure deformation in meat. Although shorter thawing times are known to reduce meat deterioration, Lan et al. (2021) demonstrated that microwave thawing results in serious muscle damage owing to regional heating. However, a study discovered that radiofrequency thawing can reduce alterations in the structure and properties of proteins, thereby increasing the thawing efficiency.

Each skeletal muscle has its own morphological, physiological, kinematic, and functional characteristics. However, muscle cells (fibers) exhibit varying physicochemical characteristics (Park et al., 2022). As demonstrated in Table 2, representative porcine skeletal muscles were identified based on their varying compositions of muscle fiber types (I, IIA, IIX, and IIB), resulting in varied pH, meat color, WHC, and tenderness (Chang et al., 2003; Park et al., 2022; Ruusunen and Puolanne, 2004). Moreover, various bovine skeletal muscles have been profiled and their fiber characteristics and physicochemical properties were differentiated (Joo et al., 2017; Lang et al., 2020; Von Seggern et al., 2005). As reported by Cheng et al. (2020) and Cheng et al. (2021), various levels of quality alterations through freeze-thaw process are anticipated between various muscle types exhibiting varying muscle fiber characteristics and meat quality. Specifically, porcine skeletal muscles with a higher type II fiber composition and lower type I fiber composition, such as *M. longissimus dorsi*, *M. gluteus superficialis*, *M. semimembranosus*, and *M. semitendinosus* should avoid freezing because type II fibers are highly susceptible to freezing than type I fibers (Chang et al., 2003; Park et al., 2022; Ruusunen and Puolanne, 2004; Song et al., 2020). For bovine skeletal muscles, *M. psoas major* and *M. superficialis flexors* are anticipated to remain stable when frozen owing to their higher type I fiber composition than that of type II fibers. In contrast, bovine *M. semitendinosus*, *M. semimembranosus*, and *M. gluteus medius*, which consist predominantly of type IIB fibers, are anticipated to be susceptible to freezing (Joo et al., 2017; Lang et al., 2020). The muscles are prone to instability during freezing, specifically in pork loin and round cuts, in contrast to the highly stable muscles during freezing, such as tenderloin and a part of the shank. Until recently, with only limited studies, alterations in meat quality owing to freezing and thawing, considering the muscle fiber composition of each cut, have not been adequately assessed. Furthermore, no freezing and thawing technologies were introduced.

However, the effects of meat components, such as moisture, fat, protein, and collagen on alterations in muscle tissue and

**Table 2. Previous studies on muscle fiber characteristics and physicochemical properties in various porcine and bovine skeletal muscles**

Species	Breed	Sex/age/carcaass weight	Muscles	Outcome	References
Porcine	Berkshire Tamworth Duroc Large White	Male/NP/NP	<i>M. longissimus dorsi</i> <i>M. psoas major</i>	<p>↑ Type I, muscle fiber proportion in <i>M. psoas major</i> of the Berkshire</p> <p>↑ Type I, muscle fiber proportion in <i>M. longissimus dorsi</i> of the Large White</p> <p>↑ Type IIA, muscle fiber proportion in <i>M. psoas major</i> than in <i>M. longissimus dorsi</i></p> <p>↑ Type IIB and ↓ type IIA, muscle fiber amount in <i>M. longissimus dorsi</i> of Berkshire, Duroc and Tamworth</p> <p>↓ Type I, muscle fiber amount in Large White and ↑ type I amount in both muscles of Duroc</p> <p>↑ Type IIA, muscle fiber amount in Duroc and ↓ type IIA amount in Large White and Tamworth</p> <p>↓ Type IIB and ↓ type IIX, muscle fiber amount in <i>M. psoas major</i> of Large White</p> <p>↓ pH at 45 min, drip loss, CIE L*, and hue value in <i>M. psoas major</i> than <i>M. longissimus dorsi</i></p> <p>↑ pH at 24 h, CIE a* and CIE b* in <i>M. psoas major</i> than <i>M. longissimus dorsi</i></p>	Chang et al. (2003)
Porcine	Wild Domestic	Male and female/ 165±2 days/ 105.1±8.8 kg	<i>M. longissimus dorsi</i> <i>M. gluteus superficialis</i> <i>M. semimembranosus</i> <i>M. infra spinam</i> <i>M. masseter</i>	<p>↓ Type IIB and ↑ type IIA, muscle fiber area % in <i>M. longissimus dorsi</i>, <i>M. gluteus superficialis</i>, and <i>M. semimembranosus</i> of wild pig</p> <p>↑ Type IIB, muscle fiber area % in <i>M. longissimus dorsi</i>, <i>M. gluteus superficialis</i>, and <i>M. semimembranosus</i> in domestic pig</p> <p>↑ Type IIA and ↓ type IIB, muscle fiber area % in <i>M. infra spinam</i> of wild pig</p> <p>↑ Type I, muscle fiber area % in <i>M. masseter</i> of wild pig</p>	Ruusunen and Puolanne (2004)
Porcine	LYD	Castrated/NP/ 79.2±3.5 kg	<i>M. infrahyoid</i> <i>M. rectus abdominis</i> <i>M. gracilis</i> <i>M. psoas major</i> <i>M. semitendinosus</i> <i>M. semimembranosus</i> <i>M. vastus</i> <i>M. diaphragm</i> <i>M. longissimus dorsi</i> <i>M. biceps brachii</i> <i>M. biceps femoris</i> <i>M. rectus femoris</i> <i>M. subscapularis</i> <i>M. superficialis digital flexor</i>	<p>↓ Moisture content, and high CIE L*, cooking loss, and WBSF in <i>M. semitendinosus</i> than in <i>M. psoas major</i> and <i>M. vastus</i></p> <p>↑ Moisture, crude protein, and WBSF but ↓ fat content, pH, CIE a*, and CIE b* in <i>M. longissimus dorsi</i> than in <i>M. diaphragm</i></p> <p>↓ Moisture content, higher crude fat content, and ↑ CIE b* in <i>M. biceps femoris</i> than in <i>M. biceps brachii</i> and <i>M. rectus femoris</i></p> <p>↑ pH and WBSF, and ↓ drip loss in <i>M. subscapularis</i> than in <i>M. superficialis digital flexor</i></p> <p>↑ All muscle fiber area in <i>M. psoas major</i> than in <i>M. semimembranosus</i>, <i>M. vastus</i>, and <i>M. semitendinosus</i></p> <p>↓ Type I, relative fiber area in <i>M. semitendinosus</i></p> <p>↓ Type IIA and IIXB, relative fiber area in <i>M. semimembranosus</i></p> <p>↑ Type IIA, relative fiber area in <i>M. diaphragm</i> than in <i>M. longissimus dorsi</i></p> <p>↑ Type IIX and ↓ type IIB in <i>M. biceps brachii</i> than in <i>M. biceps femoris</i> and <i>M. rectus femoris</i></p> <p>↓ Type IIA, muscle fiber density in <i>M. biceps femoris</i></p> <p>↑ Type IIX and ↓ type IIB in <i>M. biceps brachii</i> than in <i>M. biceps femoris</i> and <i>M. rectus femoris</i></p> <p>↓ Type IIA, muscle fiber density in <i>M. biceps femoris</i></p>	Park et al. (2022)

**Table 2.** Previous studies on muscle fiber characteristics and physicochemical properties in various porcine and bovine skeletal muscles (continued)

Species	Breed	Sex/age/carcass weight	Muscles	Outcome	References
Bovine	Chinese Simmental cattle	Bull/ 26 months/ 378±30 kg	<i>M. longissimus thoracis</i> <i>M. psoas major</i> <i>M. semitendinosus</i>	<p>↑ Type I, IIA, and IIB muscle fiber density in <i>M. longissimus thoracis</i> and <i>M. semitendinosus</i> than in <i>M. psoas major</i></p> <p>↑ Type I, muscle fiber number % in <i>M. psoas major</i>.</p> <p>↓ Type I, muscle fiber number % in <i>M. semitendinosus</i></p> <p>↑ Type IIB, muscle fiber number % in <i>M. semitendinosus</i>.</p> <p>↓ Type IIB, muscle fiber number % in <i>M. psoas major</i></p> <p>↑ WBSF in <i>M. semitendinosus</i> and <i>M. longissimus thoracis</i> than in <i>M. psoas major</i></p> <p>↓ WBSF, hardness, and chewiness values in <i>M. psoas major</i> than in <i>M. semitendinosus</i> and <i>M. longissimus thoracis</i></p> <p>↓ Pressing loss in <i>M. semitendinosus</i> and <i>M. longissimus thoracis</i> at 1, 3, and 21 days of aging</p> <p>↑ Cooking loss in all muscles at 1 and 3 days of aging</p> <p>↑ CIE L* in <i>M. semitendinosus</i> at 1 and 21 days of aging</p> <p>↑ CIE a* in <i>M. psoas major</i> at 1 and 3 days of aging</p> <p>↑ CIE b* in <i>M. semitendinosus</i> than in <i>M. psoas major</i> at 7, 14, and 21 days of aging</p>	Lang et al. (2020)
Bovine	Hanwoo	Steer/NP/NP	<i>M. longissimus lumborum</i> <i>M. psoas major</i> <i>M. semimembranosus</i> <i>M. semitendinosus</i> <i>M. gluteus medius</i> <i>M. triceps brachii</i> <i>M. rectus abdominis</i> <i>M. superficialis flexor</i>	<p>↑ Type I, muscle fiber number % in <i>M. superficialis flexor</i></p> <p>↑ Type IIA, muscle fiber number % in <i>M. psoas major</i> and <i>M. triceps brachii</i></p> <p>↓ Type IIA, muscle fiber number % in <i>M. rectus abdominis</i></p> <p>Type IIB fiber area %: <i>M. semitendinosus</i>&gt;<i>M. semimembranosus</i>&gt;<i>M. longissimus lumborum</i></p> <p>↓ Type IIB, muscle fiber number % in <i>M. superficialis flexor</i></p> <p>↑ Type I, muscle fiber area % in <i>M. superficialis flexor</i> and ↓ Type I, muscle fiber area % in <i>M. semitendinosus</i></p> <p>↑ Type IIA, muscle fiber area % in <i>M. gluteus medius</i> and ↓ Type IIA, muscle fiber area % in <i>M. longissimus lumborum</i></p> <p>↑ Moisture and collagen contents in <i>M. superficialis flexor</i> and ↓ moisture and collagen contents in <i>M. longissimus lumborum</i></p> <p>↑ CIE a* in <i>M. psoas major</i> and <i>M. superficialis flexor</i> and ↓ CIE a* in <i>M. semimembranosus</i></p> <p>↑ CIE L* in <i>M. longissimus lumborum</i> and ↓ CIE L* in <i>M. superficialis flexor</i></p> <p>↑ Drip loss and cooking loss in <i>M. semimembranosus</i></p> <p>↓ Drip loss and cooking loss in <i>M. longissimus lumborum</i> and <i>M. psoas major</i></p> <p>↑ WBSF in <i>M. superficialis flexor</i> and ↓ WBSF in <i>M. psoas major</i></p>	Joo et al. (2017)

**Table 2.** Previous studies on muscle fiber characteristics and physicochemical properties in various porcine and bovine skeletal muscles (continued)

Species	Breed	Sex/age/carcass weight	Muscles	Outcome	References
Bovine	NP	NP/NP/ 250–431 kg	Chuck 27 muscles Round 12 muscles	↑ Fat content in <i>M. cutaneous omo brachialis</i> , <i>M. longissimus costarum</i> , <i>M. multifidus</i> and <i>spinalis dorsi</i> , <i>M. serratus ventralis</i> , and <i>M. superficial pectoral</i> ↑ WHC in <i>M. deltoideus</i> and <i>M. supraspinatus</i> ↑ WBSF in <i>M. biceps brachii</i> , <i>M. infraspinatus</i> , <i>M. levatores costarum</i> , <i>M. multifidus</i> and <i>spinalis dorsi</i> , <i>M. serratus ventralis</i> , <i>M. subscapularis</i> , and <i>M. teres major</i> ↑ pH in <i>M. dorsalis oblique</i> , <i>M. longissimus costarum</i> , <i>M. levatores costarum</i> , <i>M. multifidus</i> and <i>spinalis dorsi</i> , <i>M. rhomboidus</i> , <i>M. serratus ventralis</i> , <i>M. subscapularis</i> , <i>M. supraspinatus</i> , and <i>M. trapezius</i>	Von Seggern et al. (2005)

↑ and >, higher or increase; ↓, lower or decrease.

NP, not provided; WBSF, Warner-Bratzler shear force; LYD, Yorkshire×Landrace×Duroc; WHC, water-holding capacity.

quality characteristics during freezing and thawing have not been assessed. In pork and beef, various muscles exhibit distinct characteristics, resulting in diverse proximate compositions and meat quality properties (pH, color, tenderness, and WHC; Table 2). For example, the porcine *M. psoas major* has a higher pH than *M. longissimus dorsi*, whereas biceps femoris has a lower moisture content and higher fat content compared to *M. biceps brachii* and *M. rectus femoris* (Park et al., 2022; Ruusunen and Puolanne, 2004). Additionally, *M. superficialis flexor* exhibits higher moisture and collagen content, whereas *M. longissimus lumborum* exhibits lower moisture and collagen content compared to the major bovine muscles (Table 2; Joo et al., 2017). Among the various muscles of beef chuck and round cuts, a higher fat content was observed in *M. cutaneous omo brachialis*, *M. longissimus costarum*, *M. multifidus*, *M. spinalis dorsi*, *M. serratus ventralis*, and *M. superficial pectoral* (Table 2; Von Seggern et al., 2005). Moreover, pH, moisture, fat, collagen, WHC, and shear force are anticipated to influence the freeze-thawed meat quality characteristics.

## Conclusion

Methods such as rapid freezing, high-pressure treatment freezing and thawing, electric field, ultrasonic treatment thawing, and adjusting the sequence of aging and freezing reduce the degradation of the quality of meat subjected to freezing or freeze-thawing. Regardless of the type of species and cuts (muscle type), improved quality can be expected in the frozen-thawed meat by applying these technologies. Additionally, susceptibility to freezing and quality alterations in meat vary based on the characteristics of each muscle (meat cut). Proper freezing or thawing treatment considering the unique characteristics of each muscle (specifically the muscle fiber characteristics) is expected to further reduce the deterioration in meat caused by freezing. Therefore, additional research is required to assess the effects of unique muscle characteristics (proximate composition, pH, WHC, tenderness, muscle fiber characteristics, etc.) on alterations in the quality of frozen-thawed meat. In conclusion, freezing is a hygienic and safe method to extend the shelf life of meat. However, it is essential to consider unique meat characteristics when implementing technical enhancements to minimize the adverse effects of freeze-thawing on meat quality.

## Conflicts of Interest

The authors declare no potential conflicts of interest.

## Acknowledgements

This work was supported by the National Research Foundation of Korea (NRF) grant funded by Korea government (MSIT) (NRF-2022R1A2C4002360).

## Author Contributions

Conceptualization: Kim GD. Data curation: Im C, Kim GD. Formal analysis: Im C, Song S. Methodology: Im C, Cheng H, Park J, Kim GD. Writing - original draft: Im C. Writing - review & editing: Im C, Song S, Cheng H, Park J, Kim GD.

## Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

## References

- Alonso V, Muela E, Tenas J, Calanche JB, Roncalés P, Beltrán JA. 2016. Changes in physicochemical properties and fatty acid composition of pork following long-term frozen storage. *Eur Food Res Technol* 242:2119-2127.
- Añón MC, Calvelo A. 1980. Freezing rate effects on the drip loss of frozen beef. *Meat Sci* 4:1-14.
- Bevilacqua A, Zaritzky NE, Calvelo A. 1979. Histological measurements of ice in frozen beef. *Int J Food Sci Technol* 14:237-251.
- Bond JJ, Can LA, Warner RD. 2004. The effect of exercise stress, adrenaline injection and electrical stimulation on changes in quality attributes and proteins in *semimembranosus* muscle of lamb. *Meat Sci* 68:469-477.
- Borch E, Kant-Muermans ML, Blixt Y. 1996. Bacterial spoilage of meat and cured meat products. *Int J Food Microbiol* 33:103-120.
- Brash AR. 1999. Lipoxygenases: Occurrence, functions, catalysis, and acquisition of substrate. *J Biol Chem* 274:23679-23682.
- Carpenter CE, Cornforth DP, Whittier D. 2001. Consumer preferences for beef color and packaging did not affect eating satisfaction. *Meat Sci* 57:359-363.
- Cevik M, Icier F. 2020. Characterization of viscoelastic properties of minced beef meat thawed by ohmic and conventional methods. *Food Sci Technol Int* 26:277-290.
- Chang KC, da Costa N, Blackley R, Southwood O, Evans G, Plastow G, Wood JD, Richardson RI. 2003. Relationships of myosin heavy chain fibre types to meat quality traits in traditional and modern pigs. *Meat Sci* 64:93-103.
- Cheftel JC, Lévy J, Dumay E. 2000. Pressure-assisted freezing and thawing: Principles and potential applications. *Food Rev Int* 16:453-483.
- Cheftel JC, Thiebaud M, Dumay E. 2002. Pressure-assisted freezing and thawing of foods: A review of recent studies. *High*



Press Res 22:601-611.

- Cheng H, Song S, Jung EY, Jeong JY, Joo ST, Kim GD. 2020. Comparison of beef quality influenced by freeze-thawing among different beef cuts having different muscle fiber characteristics. *Meat Sci* 169:108206.
- Cheng H, Song S, Kim GD. 2021. Frozen/thawed meat quality associated with muscle fiber characteristics of porcine *longissimus thoracis et lumborum*, *psaos major*, *semimembranosus*, and *semitendinosus* muscles. *Sci Rep* 11:13354.
- Cheng Q, Sun DW. 2008. Factors affecting the water holding capacity of red meat products: A review of recent research advances. *Crit Rev Food Sci Nutr* 48:137-159.
- Chevalier D, Sequeira-Munoz A, Le Bail A, Simpson BK, Ghouil M. 2000. Effect of freezing conditions and storage on ice crystal and drip volume in turbot (*Scophthalmus maximus*): Evaluation of pressure shift freezing vs. air-blast freezing. *Innov Food Sci Emerg Technol* 1:193-201.
- Cho S, Kang SM, Seong P, Kang G, Kim Y, Kim J, Chang S, Park B. 2017. Effect of aging and freezing conditions on meat quality and storage stability of 1<sup>++</sup> grade Hanwoo steer beef: Implications for shelf life. *Korean J Food Sci Anim Resour* 37:440-448.
- Choi MJ, Min SG, Hong GP. 2016. Effects of pressure-shift freezing conditions on the quality characteristics and histological changes of pork. *LWT-Food Sci Technol* 67:194-199.
- Cooke R, Wien R. 1971. The state of water in muscle tissue as determined by proton nuclear magnetic resonance. *Biophys J* 11:1002-1017.
- Coombs CEO, Holman BWB, Friend MA, Hopkins DL. 2017. Long-term red meat preservation using chilled and frozen storage combinations: A review. *Meat Sci* 125:84-94.
- Dang DS, Bastarrachea LJ, Martini S, Matarneh SK. 2021. Crystallization behavior and quality of frozen meat. *Foods* 10:2707.
- Dave D, Ghaly AE. 2011. Meat spoilage mechanisms and preservation techniques: A critical review. *Am J Agric Biol Sci* 6:486-510.
- Domínguez R, Pateiro M, Gagaoua M, Barba FJ, Zhang W, Lorenzo JM. 2019. A comprehensive review on lipid oxidation in meat and meat products. *Antioxidants* 8:429.
- Enfält AC, Lundesjö Ahnström M, Svensson K, Hansson I, Lundström K. 2004. Tenderness in *M. longissimus dorsi* from cows: Effect of pelvic suspension and ageing time. Proceedings of the 50<sup>th</sup> International Congress of Meat Science and Technology, Helsinki, Finland.
- Fallah-Joshaqani S, Hamdami N, Keshavarzi E, Keramat J, Dalvi-Isfahan M. 2019. Evaluation of the static electric field effects on freezing parameters of some food systems. *Int J Refrig* 99:30-36.
- Fang YZ, Yang S, Wu G. 2002. Free radicals, antioxidants, and nutrition. *Nutrition* 18:872-879.
- Fernández PP, Otero L, Guignon B, Sanz PD. 2006. High-pressure shift freezing versus high-pressure assisted freezing: Effects on the microstructure of a food model. *Food Hydrocoll* 20:510-522.
- Fernández PP, Sanz PD, Molina-García AD, Otero L, Guignon B, Vaudagna SR. 2007. Conventional freezing plus high pressure-low temperature treatment: Physical properties, microbial quality and storage stability of beef meat. *Meat Sci* 77:616-625.
- Fioramonti SA, Rubiolo AC, Santiago LG. 2017. Characterisation of freeze-dried flaxseed oil microcapsules obtained by multilayer emulsions. *Powder Technol* 319:238-244.
- Gan S, Zhang M, Mujumdar AS, Jiang Q. 2022. Effects of different thawing methods on quality of unfrozen meats. *Int J*

Refrigeration 134:168-175.

- Gill CO, Penney N. 1977. Penetration of bacteria into meat. *Appl Environ Microbiol* 33:1284-1286.
- Gonzalez-Sanguinetti S, Añón MC, Calvelo A. 1985. Effect of thawing rate on the exudate production of frozen beef. *J Food Sci* 50:697-700.
- Guo Z, Ge X, Yang L, Ma G, Ma J, Yu Q, Han L. 2021. Ultrasound-assisted thawing of frozen white yak meat: Effects on thawing rate, meat quality, nutrients, and microstructure. *Ultrason Sonochem* 70:105345.
- Gupta LK, Nagamohini Y. 1992. Penetration of poultry meat by *Pseudomonas* and *Lactobacillus* spp. *World J Microbiol Biotechnol* 8:212-213.
- Guyon C, Meynier A, de Lamballerie M. 2016. Protein and lipid oxidation in meat: A review with emphasis on high-pressure treatments. *Trends Food Sci Technol* 50:131-143.
- Hansen E, Juncher D, Henckel P, Karlsson A, Bertelsen G, Skibsted LH. 2004. Oxidative stability of chilled pork chops following long term freeze storage. *Meat Sci* 68:479-484.
- He X, Liu R, Tatsumi E, Nirasawa S, Liu H. 2014. Factors affecting the thawing characteristics and energy consumption of frozen pork tenderloin meat using high-voltage electrostatic field. *Innov Food Sci Emerg Technol* 22:110-115.
- Honikel KO. 1998. Reference methods for the assessment of physical characteristics of meat. *Meat Sci* 49:447-457.
- Honikel KO, Hamm R. 1994. Measurement of water-holding capacity and juiciness. In *Quality attributes and their measurement in meat, poultry and fish products*. Pearson AM, Dutson TR (ed). Springer, New York, NY, USA. pp 125-161.
- Hood DE, Riordan EB. 1973. Discolouration in pre-packaged beef: Measurement by reflectance spectrophotometry and shopper discrimination. *J Food Sci Technol* 8:333-343.
- Hou Q, Cheng Y, Kang D, Zhang W, Zhou G. 2020. Quality changes of pork during frozen storage: Comparison of immersion solution freezing and air blast freezing. *Int J Food Sci Technol* 55:109-118.
- Huff-Lonergan E, Lonergan SM. 2005. Mechanisms of water-holding capacity of meat: The role of postmortem biochemical and structural changes. *Meat Sci* 71:194-204.
- Jeong JY, Kim GD, Yang HS, Joo ST. 2011. Effect of freeze-thaw cycles on physicochemical properties and color stability of beef *semimembranosus* muscle. *Food Res Int* 44:3222-3228.
- Joo SH, Lee KW, Hwang YH, Joo ST. 2017. Histochemical characteristics in relation to meat quality traits of eight major muscles from Hanwoo steers. *Korean J Food Sci Anim Resour* 37:716-725.
- Joo ST, Lee JI, Ha YL, Park GB. 2002. Effects of dietary conjugated linoleic acid on fatty acid composition, lipid oxidation, color, and water-holding capacity of pork loin. *J Anim Sci* 80:108-112.
- Kim EJ, Lee SY, Park DH, Kim H, Choi MJ. 2020. Physicochemical properties of pork neck and chicken leg meat under various freezing temperatures in a deep freezer. *Food Sci Anim Resour* 40:444-460.
- Kim GD, Jung EY, Lim HJ, Yang HS, Joo ST, Jeong JY. 2013. Influence of meat exudates on the quality characteristics of fresh and freeze-thawed pork. *Meat Sci* 95:323-329.
- Kim HW, Kim JH, Seo JK, Setyabrata D, Kim YHB. 2018. Effects of aging/freezing sequence and freezing rate on meat quality and oxidative stability of pork loins. *Meat Sci* 139:162-170.
- Kim YHB, Liesse C, Kemp R, Balan P. 2015. Evaluation of combined effects of ageing period and freezing rate on quality attributes of beef loins. *Meat Sci* 110:40-45.
- King DA, Hunt MC, Barbut S, Claus JR, Cornforth DP, Joseph P, Kim YH, Lindahl G, Mancini RA, Nair MN, Merok KJ, Milkowski A, Mohan A, Pohlman F, Ramanathan R, Raines CR, Seyfert M, Sørheim O, Suman SP, Weber M. 2023.

- American meat science association guidelines for meat color measurement. *Meat Muscle Biol* 6:12473.
- Lagerstedt Å, Enfält L, Johansson L, Lundström K. 2008. Effect of freezing on sensory quality, shear force and water loss in beef *M. longissimus dorsi*. *Meat Sci* 80:457-461.
- Lan W, Zhao Y, Gong T, Mei J, Xie J. 2021. Effects of different thawing methods on the physicochemical changes, water migration and protein characteristic of frozen pompano (*Trachinotus ovatus*). *J Food Biochem* 45:e13826.
- Lang Y, Zhang S, Xie P, Yang X, Sun B, Yang H. 2020. Muscle fiber characteristics and postmortem quality of *longissimus thoracis, psoas major* and *semitendinosus* from Chinese Simmental bulls. *Food Sci Nutr* 8:6083-6094.
- Leygonie C, Britz TJ, Hoffman LC. 2012. Impact of freezing and thawing on the quality of meat: Review. *Meat Sci* 91:93-98.
- Li B, Sun DW. 2002. Novel methods for rapid freezing and thawing of foods: A review. *J Food Eng* 54:175-182.
- Lorentzen G, Hustad A, Lian F, Grip AE, Schrödter E, Medeiros T, Siikavuopio SI. 2020. Effect of freezing methods, frozen storage time, and thawing methods on the quality of mildly cooked snow crab (*Chionoecetes opilio*) clusters. *LWT-Food Sci Technol* 123:109103.
- Love RM, Haraldsson SB. 1961. The expressible fluid of fish fillets. XI.—Ice crystal formation and cell damage in cod muscle frozen before rigor mortis. *J Sci Food Agric* 12:442-449.
- Mancini RA, Hunt MC. 2005. Current research in meat color. *Meat Sci* 71:100-121.
- Marriott NG, Garcia RA, Pullen JH, Lee DR. 1980. Effect of thaw conditions on ground beef. *J Food Prot* 43:180-184.
- McMillin KW. 2008. Where is MAP going? A review and future potential of modified atmosphere packaging for meat. *Meat Sci* 80:43-65.
- Mercier Y, Gatellier P, Viau M, Remignon H, Renerre M. 1998. Effect of dietary fat and vitamin E on colour stability and on lipid and protein oxidation in turkey meat during storage. *Meat Sci* 48:301-318.
- Min SG, Hong GP, Chun JY, Park SH. 2016. Pressure ohmic thawing: A feasible approach for the rapid thawing of frozen meat and its effects on quality attributes. *Food Bioprocess Technol* 9:564-575.
- Morrissey PA, Sheehy PJA, Galvin K, Kerry JP, Buckley DJ. 1998. Lipid stability in meat and meat products. *Meat Sci* 49:S73-S86.
- Nakazawa N, Okazaki E. 2020. Recent research on factors influencing the quality of frozen seafood. *Fish Sci* 86:231-244.
- Ngapo TM, Babare IH, Reynolds J, Mawson RF. 1999. Freezing and thawing rate effects on drip loss from samples of pork. *Meat Sci* 53:149-158.
- Papuc C, Goran GV, Predescu CN, Nicorescu V, Stefan G. 2017. Plant polyphenols as antioxidant and antibacterial agents for shelf-life extension of meat and meat products: Classification, structures, sources, and action mechanisms. *Compr Rev Food Sci Food Saf* 16:1243-1268.
- Park J, Song S, Cheng H, Im C, Jung EY, Moon SS, Choi J, Hur SJ, Joo ST, Kim GD. 2022. Comparison of meat quality and muscle fiber characteristics between porcine skeletal muscles with different architectures. *Food Sci Anim Resour* 42:874-888.
- Park MH, Kwon JE, Kim SR, Won JH, Ji JY, Hwang IK, Kim MR. 2012. Physicochemical and microbiological properties of pork by various thawing methods. *J East Asian Soc Diet Life* 22:298-304.
- Patist A, Bates D. 2008. Ultrasonic innovations in the food industry: From the laboratory to commercial production. *Innov Food Sci Emerg Technol* 9:147-154.
- Pikul J, Leszczynski DE, Kummerow FA. 1985. Oxidation products in chicken meat after frozen storage, microwave and convection oven cooking, refrigerated storage, and reheating. *Poult Sci* 64:93-100.

- Pogorzelski G, Pogorzelska-Nowicka E, Pogorzelski P, Póltorak A, Hocquette JF, Wierzbicka A. 2022. Towards an integration of pre- and post-slaughter factors affecting the eating quality of beef. *Livest Sci* 255:104795.
- Poudyal RL, Maekawa R, Redo MA, Khanal R, Suzuki T, Watanabe M. 2023. Effect of supercooled freezing on the quality of pork tenderloin meat under different thawing conditions. *Food Control* 144:109331.
- Renner M, Poncet K, Mercier Y, Gatellier P, Métro B. 1999. Influence of dietary fat and vitamin E on antioxidant status of muscles of turkey. *J Agric Food Chem* 47:237-244.
- Ruusunen M, Puolanne E. 2004. Histochemical properties of fibre types in muscles of wild and domestic pigs and the effect of growth rate on muscle fibre properties. *Meat Sci* 67:533-539.
- Sanz PD, de Elvira C, Martino M, Zaritzky N, Otero L, Carrasco JA. 1999. Freezing rate simulation as an aid to reducing crystallization damage in foods. *Meat Sci* 52:275-278.
- Schudel S, Prawiranto K, Defraeye T. 2021. Comparison of freezing and convective dehydrofreezing of vegetables for reducing cell damage. *J Food Eng* 293:110376.
- Shanks BC, Wulf DM, Maddock RJ. 2002. The effect of freezing on Warner-Bratzler shear force values of beef *longissimus* steaks across several postmortem aging periods. *J Anim Sci* 80:2122-2125.
- Škrovánková S, Mišurcová L, Machů L. 2012. Antioxidant activity and protecting health effects of common medicinal plants. *Adv Food Nutr Res* 67:75-139.
- Song S, Ahn CH, Kim GD. 2020. Muscle fiber typing in bovine and porcine skeletal muscles using immunofluorescence with monoclonal antibodies specific to myosin heavy chain isoforms. *Food Sci Anim Resour* 40:132-144.
- Soyer A, Özalp B, Dalmış Ü, Bilgin V. 2010. Effects of freezing temperature and duration of frozen storage on lipid and protein oxidation in chicken meat. *Food Chem* 120:1025-1030.
- Su G, Ramaswamy HS, Zhu S, Yu Y, Hu F, Xu M. 2014. Thermal characterization and ice crystal analysis in pressure shift freezing of different muscle (shrimp and porcine liver) versus conventional freezing method. *Innov Food Sci Emerg Technol* 26:40-50.
- Suman SP, Joseph P. 2013. Myoglobin chemistry and meat color. *Annu Rev Food Sci Technol* 4:79-99.
- Tang J, Faustman C, Hoagland TA, Mancini RA, Seyfert M, Hunt MC. 2005a. Interactions between mitochondrial lipid oxidation and oxymyoglobin oxidation and the effects of vitamin E. *Food Chem* 53:6073-6079.
- Tang J, Faustman C, Hoagland TA, Mancini RA, Seyfert M, Hunt MC. 2005b. Postmortem oxygen consumption by mitochondria and its effects on myoglobin form and stability. *Food Chem* 53:1223-1230.
- Tang J, Faustman C, Mancini RA, Seyfert M, Hunt MC. 2005c. Mitochondrial reduction of metmyoglobin: Dependence on the electron transport chain. *J Agric Food Chem* 53:5449-5455.
- Valero Díaz A, Calleja CA, Fernández Escámez P, Manuel C, Abuín F, Sillué SM, Moragas GS. 2023. Report of the Scientific Committee of the Spanish Agency for Food Safety and Nutrition (AESAN) on the determination of the date of minimum durability for frozen meat in retail establishments. *Food Risk Assess Eur* 1:209-236.
- Vieira C, Diaz MT, Martínez B, García-Cachán MD. 2009. Effect of frozen storage conditions (temperature and length of storage) on microbiological and sensory quality of rustic crossbred beef at different states of ageing. *Meat Sci* 83:398-404.
- Von Seggern DD, Calkins CR, Johnson DD, Brickler JE, Gwartney BL. 2005. Muscle profiling: Characterizing the muscles of the beef chuck and round. *Meat Sci* 71:39-51.
- Wang Y, Liang H, Xu R, Lu B, Song X, Liu B. 2020. Effects of temperature fluctuations on the meat quality and muscle

- microstructure of frozen beef. *Int J Refrig* 116:1-8.
- Xanthakis E, Havet M, Chevallier S, Abadie J, Le-Bail A. 2013. Effect of static electric field on ice crystal size reduction during freezing of pork meat. *Innov Food Sci Emerg Technol* 20:115-120.
- Yun YC, Kim H, Ramachandraiah K, Hong GP. 2021. Evaluation of the relationship between freezing rate and quality characteristics to establish a new standard for the rapid freezing of pork. *Food Sci Anim Resour* 41:1012-1021.
- Zhang L, Li J, Zhou K. 2010. Chelating and radical scavenging activities of soy protein hydrolysates prepared from microbial proteases and their effect on meat lipid peroxidation. *Bioresour Technol* 101:2084-2089.
- Zhang M, Li F, Diao X, Kong B, Xia X. 2017. Moisture migration, microstructure damage and protein structure changes in porcine *longissimus* muscle as influenced by multiple freeze-thaw cycles. *Meat Sci* 133:10-18.
- Zhang R, Realini CE, Kim YHB, Farouk MM. 2023. Challenges and processing strategies to produce high quality frozen meat. *Meat Sci* 205:109311.
- Zhou GH, Xu XL, Liu Y. 2010. Preservation technologies for fresh meat: A review. *Meat Sci* 86:119-128.
- Zhu S, Le Bail A, Chapleau N, Ramaswamy HS, de Lamballerie-Anton M. 2004. Pressure shift freezing of pork muscle: Effect on color, drip loss, texture, and protein stability. *Biotechnol Prog* 20:939-945.