

ARTICLE

Effects of Edible Insect Powders as Meat Partial Substitute on Physicochemical Properties and Storage Stability of Pork Patties

 Nayoung Choi¹, Sanghun Park¹, Yunhwan Park¹, Gyutae Park¹, Sehyuk Oh¹, Yun-a Kim¹, Youngho Lim¹, Soyoung Jang¹, Youngjin Kim¹, Ki-Su Ahn², Xi Feng³, and Jungseok Choi^{1,*}
¹Department of Animal Science, Chungbuk National University, Cheongju 28644, Korea

²Chungcheongbuk-do Research and Extension Services, Cheongju 28130, Korea

³Department of Nutrition, Food Science, and Packaging, San Jose State University, San Jose, CA 95192, United States

 OPEN ACCESS

Received November 9, 2023

Revised February 11, 2024

Accepted February 13, 2024

***Corresponding author** : Jungseok Choi
 Department of Animal Science, Chungbuk National University, Cheongju 28644, Korea
 Tel: +82-43-261-2551
 Fax: +82-43-273-2240
 E-mail: jchoi@chungbuk.ac.kr

***ORCID**

Nayoung Choi
<https://orcid.org/0000-0002-4782-1098>
 Sanghun Park
<https://orcid.org/0000-0003-4804-0848>
 Yunhwan Park
<https://orcid.org/0000-0002-2239-6697>
 Gyutae Park
<https://orcid.org/0000-0003-1614-1097>
 Sehyuk Oh
<https://orcid.org/0000-0003-4105-2512>
 Yun-a Kim
<https://orcid.org/0000-0002-5505-030X>
 Youngho Lim
<https://orcid.org/0000-0002-0238-4736>
 Soyoung Jang
<https://orcid.org/0009-0001-1146-2695>
 Youngjin Kim
<https://orcid.org/0009-0002-6243-3250>
 Ki-Su Ahn
<https://orcid.org/0009-0007-7748-8349>
 Xi Feng
<https://orcid.org/0000-0001-7229-3752>
 Jungseok Choi
<https://orcid.org/0000-0001-8033-0410>

Abstract In this study, physicochemical and antioxidant properties, and storage stability (1, 3, and 7 days) of pork patties added with edible insect powders (EIP) of four species (Larvae of *Tenentobryonia molitor*, *Protaetia brevitarsis seulensis*, *Allomyrina dichotoma*, and *Gryllus bimaculatus*) as meat partial substitutes were investigated. Twenty percent of each EIP was added to pork patties, and four treatments were prepared. On the other hand, two control groups were set, one with 0.1 g of ascorbic acid and the other without anything. Adding EIP decreased water content but increased protein, fat, carbohydrate, and ash contents. In addition, the use of EIP increased the water holding capacity and texture properties as well as decreased the cooking loss. However, the sensory evaluation and storage stability were negatively affected by the addition of EIP. The 2,2-diphenyl-1-picrylhydrazyl radical scavenging activity had a positive effect on storage stability. It is believed that the addition of EIP resulted in high antioxidants due to the presence of polyphenol compounds in EIP. These results indicate that EIP has great potential to be used as meat partial substitute to improve the quality improvement and antioxidant in pork patties. However, in order to improve storage stability and consumer preference, further research is needed to apply it to patties by reducing the amount of EIP or adding auxiliary ingredients.

Keywords edible insects, partial substitute, phenolic compounds, cooking loss, antioxidant

Introduction

As the world population grows, meat consumption per capita also increases (Alexandratos and Bruinsma, 2012; van Huis et al., 2013). Meat is an important protein source in the human diet because it contains adequate and balanced amino acid composition (van der Weele et al., 2019; Wu, 2022). However, it is challenging to meet

the demand for meat (animal protein) due to climate change and the reduction of agricultural land (Abbasi and Abbasi, 2016; Premalatha et al., 2011; van Huis and Tomberlin, 2017). Therefore, there is a need to use an alternative protein source that can replace meat to meet the growing demand for proteins.

Protein is an essential macronutrient that must be obtained from animals (meat, dairy, etc.) and vegetables (legumes, etc.). Adequate intake of protein is necessary for health (WHO, 2007; Wu et al., 2014). Edible insects are often proposed as a substitute for animal proteins because they are known to be high in protein, more than 50% (Beets, 1997; Bukkens, 1997). Edible insects provide protein and energy and are high in monounsaturated fatty acids and polyunsaturated fatty acids. They are also rich in micronutrients, such as copper (Cu) and iron (Fe; Rumpold and Schlüter, 2013). They produce less greenhouse and ammonia gas as well as with efficient land use, and productive yield (Klunder et al., 2012; Oonincx et al., 2010). As a result, edible insects show great potential to be used as an environmentally friendly future food resource (Akhtar and Isman, 2018).

It has been reported that 10 species of insects are consumed in Korea and more than 1,900 species of insects are consumed worldwide (Lange and Nakamura, 2021). The representative insects that are eaten in Korea include *Tenebrio molitor* L., *Protaetia brevitarsis seulensis* L., *Gryllus bimaculatus* and *Allomyrin dichotoma* L. *T. molitor* L. is a type of *Carabidae* that is cultivated all over the world. It is mainly used as an animal feed or protein supplement because it is rich in protein and essential fatty acids (Kim et al., 2021; van Broekhoven et al., 2015). *P. brevitarsis seulensis* L. is a phytophagous insect belonging to *Cetoniinae* and has traditionally been used to treat inflammation, breast cancer, and liver disease. Its larvae have been proven to have various physiological effects such as antioxidant and antibacterial effects (Choi et al., 2019; Song et al., 2017; Suh and Kang, 2012; Yoon et al., 2003). *G. bimaculatus* belongs to *Orthoptera* and has the highest protein content and unsaturated fatty acid (UFA) content (Churchward-Venne et al., 2017; Wang et al., 2004). *A. dichotoma* L. belongs to *Scarabaeidae* and has anti-tumor, anti-hepatic fibrosis, and antibacterial effects (Miyanoshita et al., 1996; Sagisaka et al., 2001; Yoshikawa et al., 1999).

This study aimed to determine the physicochemical and antioxidant properties and storage stability of pork patties added with edible insect powder (EIP) of four species. The results of this study can pave a fundament for the future development of meat products with EIPs as protein substitutes.

Materials and Methods

Materials used in the research

Freeze-dried edible insects of four species (*T. molitor* L., *P. brevitarsis seulensis* L., *G. bimaculatus*, *A. dichotoma* L.) were provided by Chungcheongbukdo Agricultural Research and Extension Services. Edible insects were ground for 1 min using a mixer (Hanilelec, Cheongju, Korea) to prepare *T. molitor* L. powder (TMP), *P. brevitarsis seulensis* L. powder (PBP), *G. bimaculatus* powder (GBP), and *A. dichotoma* L. powder (ADP). The EIPs that were ground were stored frozen at -50°C until use. Minced pork (Manpyeong Livestock Products, Cheongju, Korea), ice, salt (Beksul, Haenam, Korea), pepper (Ottogi, Anyang, Korea), sodium tripolyphosphate (Samchun Chemical, Peongtak, Korea), and ascorbic acid (ES Food, Gunpo, Korea) was used to manufacture pork patties.

Phenolic analysis of edible insect powder by high-performance liquid chromatography

One mL of 95% methanol and 10 g of EIP were mixed. After stirring for 3 h in a constant temperature water bath at 37°C ,

the extraction was filtered using Whatman No. 2 filter paper (Advantec[®], Tokyo, Japan). Three replications were carried out.

Reversed-phase (RP) high-performance liquid chromatography (HPLC) analysis was performed using a 4.6×250 mm RP spherisorb ODS₂ column based on the method of Dimitrova et al. (2007) with minor modifications. Chromatographic analysis was performed using a Young Lin HPLC. The 20 mM potassium dihydrogen phosphate buffer (pH 2.92) was used as mobile phase A and methanol was used as mobile phase B. Elution was started with 3% methanol and allowed to reach 100% methanol in 65 min. The flow rate was at 1.0 mL/min. Phenol content was monitored at 220 nm and 280 nm. A total of seven phenolic acids (gallic acid, vanillic acid, caffeic acid, syringic acid, p-coumaric acid, phenylacetic acid, and benzoic acid) were determined with authentic standards.

Manufacturing of pork patties

Pork patties were prepared by mixing pork meat with EIPs at the ratio as shown in Table 1. Six types of pork patties were prepared with the addition of EIPs (T20, P20, G20, A20; respectively 20%), positive control (PC, 0.1% ascorbic acid), or negative control (NC, without any addition). Equal amounts of ice, salt, pepper, and sodium tripolyphosphate were added to all treatments. Finished patties were divided into 100 g each and molded with a molding machine (diameter of 10 cm, thickness of 1 mm). Manufactured patties were aged for 24 h in refrigeration at 4°C. Samples were wrapped and stored at 4°C for 7 days and evaluated on days 1, 3, and 7.

Proximate composition

Proximate composition was measured following AOAC (1990) methods. Water content was determined with a conventional oven at 105°C. Protein content was quantified with Kjeldahl method. Fat content was examined with Folch extraction method. Ash content was determined with a Muffle oven. The carbohydrate content was determined by subtracting the sum of moisture content, protein content, fat content, and ash content from the total sum of 100%.

Table 1. Formulation of pork patties

Ingredients [% (w/w)]		Treatment					
		PC	NC	T20	P20	G20	A20
Pork meat		90	90	70	70	70	70
Ice		10	10	10	10	10	10
Edible insect powder	TMP	-	-	20	-	-	-
	PBP	-	-	-	20	-	-
	GBP	-	-	-	-	20	-
	ADP	-	-	-	-	-	20
Total				100			
Ascorbic acid		0.1	-	-	-	-	-
Additive ¹⁾		1.5	1.5	1.5	1.5	1.5	1.5

¹⁾ Salt, 1.2%; pepper, 0.1%; sodium tripolyphosphate, 0.2%.

PC, positive control; NC, negative control; TMP, *Tenebrio molitor* L. powder; PBP, *Protaetia brevitarsis seulensis* L. powder; GBP, *Gryllus bimaculatus* powder; ADP, *Allomyrina dichotoma* L. powder.

pH measurements

Sample (5 g) was mixed with 50 mL of distilled water and homogenized with a stomacher (Stomacher[®] 400 Circulator, BioMed, London, UK) at 200 rpm for 30 s. Then, the pH of the mixture was determined with a pH meter (Orion STAR A211, Thermo Fisher Scientific, Waltham, MA, USA).

Water holding capacity

Water holding capacity (WHC) of sample was determined with a centrifugation method. After weighing 0.5 g of sample and placing the sample in an upper filter tube of a centrifuge tube (Union 55R, Hanil Science, Daejeon, Korea), the filter tube and sample were subjected to heating in a water bath at 80°C for 20 min. Following heating, the mixture was cooling at room temperature for 10 min. The filter tube was then placed at the bottom of the centrifuge tube and centrifuged at 447×g for 10 min. The upper filter tube was removed and weighed. The WHC value was calculated using the following formula:

$$[(\text{Total water} - \text{Free water}) / \text{Total water}] \times 100. \quad (1)$$

Cooking loss

After vacuum packaging, the sample was placed in a 70°C water bath and heated for 40 min. Water on the surface of the heated sample was wiped off, and the weight of sample was measured. The weight difference before and after heating was used to calculate the cooking loss (CL) with the following formula:

$$\text{CL} (\%) = [(\text{Initial weight (g)} - \text{Final weight (g)}) / \text{Initial weight (g)}] \times 100. \quad (2)$$

Color

The center portion of the surface of the uncooked patties was measured. Color parameters of CIE L*, CIE a*, and CIE b*-value were measured with a spectrophotometer (CM-26d, Konica Minolta, Tokyo, Japan) in the standards set of Commission Internationale de l'Éclairage (CIE). The average of measured values was obtained and recorded.

Texture profile analysis

To evaluate texture profile analysis (TPA), a rheometer (Model Compac-100, Sun Scientific, Dobbs Ferry, NY, USA) equipped with a probe (No. 3, ϕ 20 mm) of area 3.14 cm² was used. The sample size was 1 cm³ and two compression cycles were used to obtain the force versus time curve. The table speed was 60 mm/min, the crosshead speed was 200 mm/min, and the load cell was 2 kg (max. 4 kg). TPA is expressed as hardness, springiness, cohesiveness, chewiness, and gumminess.

Sensory evaluation

Sensory evaluation was performed with ten trained panelists (male and female, age range 20–29) in the Department of Animal Science at Chungbuk National University. Color, flavor, juiciness, umami, hardness, texture, and overall preference were measured. Sensory scores were assessed on a 5-point scale (1=extremely bad or undesirable, and very weak, and 5=extremely good or desirable, and very strong). The approved consent procedure for sensory evaluation is Institutional Review Board (IRB) of Chungbuk National University (No. CBNU-202302-HR-0017).

Total microbial count

A stomacher bag containing 5 g of the sample was combined with 45 mL of a 0.1% peptone solution. The mixture was homogenized in a stomacher (Stomacher® 400 Circulator, BioMed) at 200 rpm for 30 s. After serially diluting the homogeneous sample, it was inoculated into a plate count agar medium (Microgiene, Suwon, Korea) and incubated at 37°C for 48 h. The number of microorganisms was determined using a colony counter. It is expressed as Log CFU/g.

Thiobarbituric acid reactive substance

Sample (10 g) was combined with cold 10% perchloric acid 15 mL and tertiary distilled water 25 mL using a homogenizer (AM-7, Nissei, Izumichom, Tokyo). After homogenization at 10,000 rpm for 15 s, the homogenate was filtered using Whatman No. 2 filter paper (Advantech, Tokyo, Japan). Subsequently, 5 mL of 0.02 M thiobarbituric acid solution was mixed with 5 mL of the filtrate (5 mL of tertiary distilled water for blank). The mixture was then kept in a cool, dark location for 16 h. Absorbance at 529 nm was then measured using a spectrophotometer (DU-650, Beckman, Brea, CA, USA). Absorbance was converted to malonaldehyde content using a standard curve. The resulting thiobarbituric acid reactive substance (TBARS) level was expressed as mg malonaldehyde per 1,000 g of the sample (mg MDA/kg).

Peroxide value

One gram of minced sample added into an Erlenmeyer flask. Then, 10 mL of chloroform was added to dissolve the sample and mixed with 15 mL of CH₃COOH. To prepare a saturated KI solution, 99% potassium iodide was dissolved in tertiary distilled water at a ratio of 7:3. 1 mL of saturated KI solution was added into the Erlenmeyer flask, the sample was then homogenized for 1 min and kept in the dark for 10 min. After 10 min, 30 mL of distilled water was added, and the mixture was homogenized again. Then, 1 mL of 1% starch solution was added and the solution was titrated with 0.01 N Na₂S₂O₃ solution until the indicator color was disappeared. A blank sample (distilled water) was conducted with the same procedure.

2,2-Diphenyl-1-picrylhydrazyl radical scavenging activity

Sample (5 g) was homogenized with methanol (45 mL) with a homogenizer (AM-7, Nissei), followed by filtration with a Whatman No. 2 filter paper (Advantech). Blank sample was prepared with 5 mL methanol. Reference was prepared with 1 mL of 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity and 4 mL methanol. The testing sample was prepared with 2 mL filtrate, 1 mL of DPPH radical scavenging activity, and 2 mL methanol. After wrapping with an aluminum foil, blank sample, reference, and test samples were kept in a dark room for 20 to 30 min. Then, 250 µL was added to a 96-well plate and measured the absorbance at 517 nm using a microplate reader (DU-650, Beckman). The DPPH radical scavenging activity value was calculated using the following formula:

$$\text{DPPH radical scavenging activity (\%)} = [1 - (\text{sample average absorbance} / \text{reference absorbance})] \times 100. \quad (3)$$

Statistical analysis

All experiments were repeated three times. Statistical analysis was conducted by one-way analysis of variance using the generalized linear model of SAS program (SAS Institute, Cary, NC, USA, 1999). Duncan Multiple Range Test was used to evaluate the significant differences among treatments ($p < 0.05$).

Results and Discussion

Proximate composition, pH, and phenolic compounds in edible insect powder

Phenolic acids, including gallic acid, vanillic acid, caffeic acid, syringic acid, p-coumaric acid, phenylacetic acid, and benzoic acid, were analyzed and identified (Table 2). Phenolic compounds comprise one or more aromatic rings with hydroxyl groups (Balasundram et al., 2006). They exhibit antioxidant, anti-inflammatory, and antiviral effects (Benavente-García et al., 1997). TMP consisted of gallic acid (1.17 µg/L) and p-coumaric acid (18.17 µg/L). PBP included vanillic acid (33.18 µg/L), caffeic acid (10.88 µg/L), syringic acid (17.27 µg/L), phenylacetic acid (17.51 µg/L), and benzoic acid (0.82 µg/L). GBP contained gallic acid (18.17 µg/L) and benzoic acid (0.79 µg/L). ADP had gallic acid (4.46 µg/L), caffeic acid (3.52 µg/L), syringic acid (3.83 µg/L), and phenylacetic acid (5.27 µg/L). It has been proved that the antioxidant capacity of edible insects is mainly provided by phenolic compounds. Aiello et al. (2023) reported that edible insects' phenolic compounds can exert antioxidant biological activity with potential as bioactive sources. The phenolic chemicals from edible insects could improve food quality and provide antioxidant activity (Torres-Castillo and Olazarán-Santibáñez, 2023). Therefore, phenolic compounds of the four EIPs predicted the potential to be used as natural antioxidants in food.

Proximate composition

Fig. 1 displayed that proximate composition of pork patties added with EIPs. Moisture contents were significantly lower in treatments with EIP ($p < 0.05$). Protein contents were higher in groups added with T20, P20, and G20 compared to PC and NC ($p < 0.05$). However, there was no significant difference in the treatment with A20 than PC and NC ($p < 0.05$). As shown in Table 3, the protein contents of TMP (50.65%), PBP (60.35%), GBP (60.38%), and ADP (32.75%) influence the protein contents of patties. Kim et al. (2016) reported that emulsion sausages added with insects have lower moisture content but higher protein content. Carbohydrate content was significantly higher in the group added with A20 than those of PC and NC ($p < 0.05$). This might be due to chitin, a dietary fiber, found in the exoskeleton of insects (Kipkoech, 2023; Montowska et al., 2019). Fat content and ash content were significantly higher in treatments added with EIP than those of PC and NC ($p < 0.05$). Edible insects possess a wealth of UFAs and essential minerals, including copper, iron, magnesium, selenium, and zinc (Lu et al., 2024; Zielińska et al., 2015). These components could influence fat and ash contents in pork patties.

Table 2. High-performance liquid chromatography (HPLC) of phenolic compounds of edible insect powder

Phenolic acids (µg/L)	Treatments			
	TMP	PBP	GBP	ADP
Gallic acid	1.17±0.00	ND	18.17±0.00	4.46±0.15
Vanillic acid	ND	33.18±0.00	ND	ND
Caffeic acid	ND	10.88±0.00	ND	3.52±0.00
Syringic acid	ND	17.27±0.00	ND	3.83±0.18
p-Coumaric acid	6.24±0.05	ND	ND	ND
Phenylacetic acid	ND	17.51±0.00	ND	5.27±0.01
Benzoic acid	ND	0.82±0.00	0.79±0.00	ND

TMP, *Tenebrio molitor* L. powder; PBP, *Protaetia brevitarsis seulensis* L. powder; GBP, *Gryllus bimaculatus* powder; ADP, *Allomyrina dichotoma* L. powder; ND, not detected.




Fig. 1. Proximate composition of pork patties formulated with various edible insect powders. (A) Moisture, (B) protein, (C) carbohydrate, (D) fat, and (E) ash content. ^{a-d} Means with different letters on bars indicate significant differences at $p < 0.05$. PC, positive control; NC, negative control; T20, 20% *Tenebrio molitor* L. powder; P20, 20% *Protaetia brevitarsis seulensis* L. powder; G20, 20% *Gryllus bimaculatus* powder; A20, 20% *Allomyrina dichotoma* L. powder.

Table 3. Proximate composition and pH of edible insect powder itself

Traits	Treatments			
	TMP	PBP	GBP	ADP
Protein (%)	50.65±0.47 ^b	60.35±0.51 ^a	60.38±0.18 ^a	32.75±0.34 ^c
Carbohydrate (%)	36.16±0.73 ^b	36.01±0.49 ^b	31.47±0.31 ^c	64.17±0.36 ^a
Fat (%)	11.92±0.43 ^a	2.16±0.19 ^c	7.24±0.16 ^b	2.63±0.08 ^c
Ash (%)	1.27±0.34 ^{ab}	1.47±0.16 ^a	0.91±0.06 ^b	0.45±0.06 ^c
pH	6.59±0.01 ^d	7.47±0.00 ^a	6.84±0.02 ^c	7.19±0.00 ^b

^{a-d} Means in a row with different letters are significantly different ($p < 0.05$).

TMP, *Tenebrio molitor* L. powder; PBP, *Protaetia brevitarsis seulensis* L. powder; GBP, *Gryllus bimaculatus* powder; ADP, *Allomyrina dichotoma* L. powder.

pH, water holding capacity, cooking loss, and color

The results of pH, WHC, CL, and patty color with added EIP are presented in Table 4. Additionally, Fig. 2 displays the color (visual appearance) of the patties in photographs after the cooking process. It was found that pH values were significantly higher in treatments added with EIP than those of PC and NC ($p < 0.05$). TMP, PBP, GBP, and ADP had pH values of 6.59, 7.47, 6.84, and 7.19, respectively (Table 3). The pH of EIP might be affected by the pH of the patties. The isoelectric point of meat is 5.2–5.4. A pH higher than the isoelectric point of meat products can increase WHC and lower CL (Honikel, 2008). WHC indicates the ability of meat to retain water. It is an important criterion for evaluating meat quality (Honikel, 2004; Szymańko et al., 2021). Treatments added with EIP showed significantly ($p < 0.05$) higher WHC values than those of PC and NC. When insect protein was added to phosphate-free meat emulsion increased the WHC (Kim et al., 2022). Also, CL was significantly lower in treatments added with EIP than those of PC and NC (all $p < 0.05$). This result is attributed to decreased moisture and increased solid contents of meat emulsion formulation that contains insect powders (Kim et al., 2016). Bessa et al. (2023) have been reported that CL was decreased when *Hermetia illucens* L. was added as a meat substitute to burger patties.

Compared to PC and NC, all treatments with EIP added showed lower CIE L*-value ($p < 0.05$). CIE a*-value was the

Table 4. pH, water holding capacity (WHC), cooking loss (CL), color, texture profile analysis (TPA), and sensory evaluation of pork patties formulated with various levels of edible insect powder

Traits	Treatments					
	PC	NC	T20	P20	G20	A20
pH	5.91±0.03 ^f	5.98±0.01 ^e	6.35±0.01 ^c	6.70±0.01 ^a	6.27±0.01 ^d	6.51±0.01 ^b
WHC (%)	60.14±2.24 ^{bc}	55.31±1.20 ^e	71.11±8.37 ^{ab}	68.15±4.46 ^{ab}	71.92±13.22 ^{ab}	78.06±2.21 ^a
CL (%)	19.81±0.35 ^a	20.91±0.62 ^a	8.29±0.44 ^c	8.24±0.42 ^c	12.58±2.19 ^b	10.00±0.75 ^c
(Uncooked) color						
CIE L*	53.61±0.98 ^a	53.85±0.58 ^a	46.11±0.28 ^b	36.20±0.40 ^c	41.88±0.93 ^d	44.84±0.47 ^c
CIE a*	5.54±0.39 ^b	5.25±0.97 ^b	8.23±0.20 ^a	5.75±0.31 ^b	5.77±0.04 ^b	2.39±0.08 ^c
CIE b*	13.57±1.02 ^a	13.56±0.92 ^a	14.11±0.33 ^a	9.44±0.17 ^c	11.35±0.02 ^b	4.70±0.78 ^d
TPA						
Hardness (kg)	1.74±0.31 ^b	2.22±0.21 ^b	2.16±0.47 ^b	2.21±0.26 ^b	4.22±0.70 ^a	2.00±0.10 ^b
Springiness (%)	75.52±0.58 ^a	66.29±8.08 ^a	63.07±3.01 ^{ab}	54.15±5.49 ^{bc}	73.67±2.90 ^a	51.71±0.89 ^c
Cohesiveness (%)	71.31±0.62 ^a	59.39±8.91 ^a	41.58±7.20 ^b	30.62±5.93 ^b	60.54±4.38 ^a	36.96±5.35 ^b
Chewiness (kg)	0.84±0.70 ^{bc}	1.32±0.26 ^b	0.91±0.35 ^{bc}	0.68±0.18 ^c	2.54±0.34 ^a	0.74±0.14 ^{bc}
Gumminess (kg)	123.98±21.12 ^{bc}	132.15±26.25 ^b	91.31±35.21 ^{bc}	68.15±17.96 ^c	254.08±34.07 ^a	74.18±14.35 ^{bc}
Sensory evaluation						
Color	4.30±0.67 ^a	4.20±0.63 ^a	2.50±0.53 ^b	1.30±0.48 ^c	1.60±0.52 ^c	1.20±0.42 ^c
Flavor	4.00±0.47 ^a	3.90±0.32 ^a	2.50±0.85 ^b	1.50±0.53 ^c	2.40±0.84 ^b	1.40±0.84 ^c
Juiciness	3.30±0.67 ^a	3.60±0.52 ^a	2.30±0.48 ^b	1.90±0.57 ^{bc}	1.60±0.70 ^{cd}	1.20±0.42 ^d
Umami	3.90±0.57 ^a	3.85±0.75 ^a	2.80±0.92 ^b	1.60±0.52 ^c	1.90±0.74 ^c	1.40±0.70 ^c
Texture	3.30±0.48	3.30±0.48	3.90±0.57	3.50±1.35	3.70±1.16	3.20±1.23
Overall preference	4.30±0.67 ^a	4.20±0.92 ^a	2.60±0.97 ^b	1.40±0.52 ^{cd}	2.00±0.67 ^{bc}	1.10±0.32 ^d

All values are means±SD of three replicates.

Sensory scores were assessed on a 5-point scale base on 1=extremely bad or undesirable, and very weak, and 5=extremely good or desirable, and very strong.

^{a-f} Means in a row with different letters are significantly different ($p < 0.05$).

PC, positive control; NC, negative control; T20, 20% *Tenebrio molitor* L. powder; P20, 20% *Protaetia brevitarsis seulensis* L. powder; G20, 20% *Gryllus bimaculatus* powder; A20, 20% *Allomyrina dichotoma* L. powder.



Fig. 2. Visual appearance of pork patties after cooking with various edible insect powders. PC, positive control; NC, negative control; T20, 20% *Tenebrio molitor* L. powder; P20, 20% *Protaetia brevitarsis seulensis* L. powder; G20, 20% *Gryllus bimaculatus* powder; A20, 20% *Allomyrina dichotoma* L. powder.

highest in the group with T20 and it was the lowest in treatment with A20 compared to PC and NC ($p < 0.05$). CIE b^* -value was significantly lower in the groups added with P20, G20, and A20, and higher added with T20 ($p < 0.05$). Similarly, Choi et al. (2017) reported that the addition of yellow worms (*T. molitor* L.) to Frankfurt could reduce CIE L^* -value but increase CIE a^* and CIE b^* -value. Lemke et al. (2023) reported that the addition of 20% insects to sausage products reduced in CIE L^* -value compared to 10%. Cruz-López et al. (2022) reported that the CIE L^* -value of sausage decreases when insect powder. Edible insects have different melanin contents (Wittkopp and Beldade, 2009). It could contribute color differences in the pork patties.

Texture profile analysis

The TPA values of patties made with various EIPs are shown in Table 4. The treatment with G20 had the highest hardness compared to PC and NC ($p < 0.05$). This might be due to a decrease in water content when EIP was added. Kim et al. (2017) reported that hardness increased when meat was replaced with cricket powder, which was similar to this study. Springiness was significantly higher in the group of PC, NC, and G20 ($p < 0.05$). Ho et al. (2022) reported that an increase in springiness was observed in the sausage partially substituted with cricket powder. Cohesiveness was significantly highest in G20 among the EIP-added treatments ($p < 0.05$) and showed no significant difference from PC and NC. Chewiness and Gumminess were also significantly the highest in the group with G20 added compared to other treatments ($p < 0.05$). Damasceno et al. (2023) found that albumin that can also affect the texture properties of meat is the most highly distributed protein in *G. bimaculatus* powder. The addition of albumin to meat batter can cause a change in hardness (Pietrasik, 2003). Carballo et al. (1995) reported that the addition of albumin-rich egg white to Bologna sausage increased its hardness and chewiness. Thus, the albumin component of *G. bimaculatus* could influence the tissue characteristics of pork patties. Our findings indicate that adding EIP as a meat partial substitute can improve the WHC and TPA of patties and reduce CL. Therefore, using EIP could be beneficial in enhancing the quality and texture properties of pork patties.

Sensory evaluation

Table 4 displays the sensory evaluation results of patties added with EIPs. Color, flavor, juiciness, and umami were lower in all treatments added with EIP than in those of PC and NC ($p < 0.05$). The meat juice showed a trend opposite to the WHC results, this might be due to the high-fat content of edible insects with more fat released than those of PC and NC during cooking. Pinter et al. (2021) reported that more fat was released when cooking hamburger patties added with insects. Overall preference was lower for treatments added with EIP than those of PC and NC ($p < 0.05$), which meant that the addition of EIPs to patties decreased acceptability. Megido et al. (2016) reported that beef burger patties added with mealworms had a lower preference than pure beef patties. Also, Cruz-López et al. (2022) stated that pork sausage with locust (*Sphenarium purpurascens*) powder had low preference. According to the results of this study, adding EIP to patties can lower preference, but it is believed that this can be solved by manufacturing by reducing the amount of EIP or adding auxiliary ingredients.

Total microbial count, lipid oxidation (thiobarbituric acid reactive substance, peroxide value)

Total microbial count (TMC) increased with the storage time ($p < 0.05$; Table 5). Yong et al. (2023) reported that the number of microorganisms in Tteokgalbi added with edible insect extract increased with the storage time. All treatments added with EIPs had significantly higher TMC than PC and NC ($p < 0.05$). By adding EIP, not only protein but also other

Table 5. Results of storage stability (TMC, TBARS, PV, DPPH) of pork patties formulated with various levels of edible insect powder

Traits	Treatments	Storage (days)		
		1	3	7
TMC (Log CFU/g)	PC	3.33±0.46 ^{Bd}	4.15±0.21 ^{Ad}	5.81±0.02 ^{Ac}
	NC	3.88±0.04 ^{Cc}	4.90±0.07 ^{Bc}	5.81±0.11 ^{Ac}
	T20	6.15±0.01 ^{Ba}	6.18±0.02 ^{Ba}	6.80±0.02 ^{Ab}
	P20	6.27±0.01 ^{Ca}	6.38±0.01 ^{Ba}	6.95±0.00 ^{Aa}
	G20	3.63±0.21 ^{Ccd}	4.85±0.00 ^{Bc}	6.36±0.01 ^{Ad}
	A20	5.12±0.05 ^{Bb}	5.22±0.06 ^{Bb}	6.51±0.02 ^{Ac}
TBARS (mg MDA/kg)	PC	0.05±0.01 ^{Bd}	0.06±0.01 ^{ABe}	0.09±0.02 ^{Ac}
	NC	0.04±0.01 ^{Bd}	0.08±0.01 ^{Ad}	0.10±0.01 ^{Ac}
	T20	0.18±0.01 ^{Bc}	0.19±0.01 ^{Bc}	0.29±0.01 ^{Ab}
	P20	0.23±0.00 ^b	0.25±0.02 ^b	0.26±0.03 ^b
	G20	0.19±0.04 ^{Bc}	0.25±0.01 ^{Bb}	0.43±0.06 ^{Aa}
	A20	0.33±0.02 ^{Ba}	0.39±0.01 ^{Ba}	0.48±0.07 ^{Aa}
PV (meq/kg)	PC	14.92±0.06 ^{Ba}	16.62±0.00 ^{Ac}	14.26±0.05 ^{Cb}
	NC	13.66±0.02 ^{Cb}	33.84±0.08 ^{Aa}	31.52±0.13 ^{Ba}
	T20	2.64±0.00 ^{Bd}	7.62±0.00 ^{Af}	2.32±0.01 ^{Ce}
	P20	0.66±0.00 ^{Ce}	18.28±0.02 ^{Ab}	14.26±0.07 ^{Bb}
	G20	3.99±0.01 ^{Cc}	13.64±0.04 ^{Ad}	5.63±0.01 ^{Bc}
	A20	0.66±0.00 ^{Ce}	8.28±0.02 ^{Ac}	2.99±0.01 ^{Bd}
DPPH (%)	PC	86.45±0.94 ^{Abc}	86.42±0.88 ^{Aa}	85.23±1.31 ^{Aa}
	NC	63.96±4.07 ^{Ad}	46.34±1.41 ^{Bd}	42.60±1.72 ^{Bd}
	T20	83.20±1.02 ^{Ac}	82.23±1.12 ^{Ac}	74.39±1.77 ^{Bc}
	P20	88.29±4.70 ^{Ab}	82.11±1.22 ^{Bc}	80.62±1.64 ^{Bb}
	G20	95.40±0.73 ^{Aa}	85.23±1.64 ^{Bab}	83.20±1.02 ^{Bab}
	A20	94.58±0.93 ^{Aa}	83.88±0.85 ^{Bbc}	83.33±1.08 ^{Bab}

^{A-C} Means in a row with different letters are significantly different ($p < 0.05$).

^{a-f} Means in a column with different letters are significantly different ($p < 0.05$).

TMC, total microbial count; TBARS, thiobarbituric acid reactive substance; PV, peroxide value; DPPH, 2,2-diphenyl-1-picrylhydrazyl; PC, positive control; NC, negative control; T20, 20% *Tenebrio molitor* L. powder; P20, 20% *Protaetia brevitarsis seulensis* L. powder; G20, 20% *Gryllus bimaculatus* powder; A20, 20% *Allomyrina dichotoma* L. powder.

nutrients increased (Fig. 1). It is thought that the number of microorganisms has increased because this nutrient-rich environment serves as a growth medium for various microorganisms (Anas et al., 2019; Elsharawy et al., 2018). Among treatments added with EIP, the treatment added with G20 had the lowest TMC ($p < 0.05$). Malm and Liceaga (2021) found that the chitosan in the cricket had antibacterial effects. Thus, the reason why treatment with G20 had lower TMC value than other EIP-added treatments might be due to high antibacterial activity of crickets.

Since lipid oxidation causes a decrease in the quality of meat and meat products, peroxide value (PV) and TBARS were used as indicators of lipid oxidation to confirm this in this study (Love and Pearson, 1971; Turgut et al., 2017). PV measures the primary product of lipid oxidation (hydrogen peroxide) while TBARS measures the end product of lipid oxidation (Gan et al., 2022; Hadidi et al., 2022; Juntachote et al., 2007; Simic and Taylor, 1987). PV showed increasing until the 3rd day and then

decreased, and TBARS increased values in all treatments as the storage period increased ($p < 0.05$; Table 5). This is because hydroperoxides decompose into secondary products (Gunstone and Norris, 1983). All treatments added with EIP showed lower PV values but higher TBARS values than those of PC and NC ($p < 0.05$). Han et al. (2023) reported that the TBARS value of hybrid sausage added with cricket flour increased. Also, it has been reported that adding silkworm pupae flours to emulsion sausage increases the TBARS value (Kim et al., 2016). It is believed to be due to the high fat content of EIP.

2,2-Diphenyl-1-picrylhydrazyl radical scavenging activity

Table 5 showed DPPH radical scavenging activity of pork patties with EIPs. The treatments added with EIP showed higher antioxidant activities than those of NC ($p < 0.05$). Yong et al. (2023) reported that Tteokgalbi with edible insect extract had high DPPH radical scavenging activity, but it decreased as the storage period increased. It also reported that the antioxidant activities of *Dactylopius opuntiae* extract were confirmed when applied to beef patties (Aragon-Martinez et al., 2023). When compared with the PC, the treatment with G20 showed a similar DPPH radical scavenging activity value. This is because GBP had a higher gallic acid content than other EIPs (Table 2). This is similar to the report that cricket (*G. bimaculatus*) has phenolic compounds, showing excellent antioxidant activity (Baigts-Allende et al., 2021; Kurdi et al., 2021). Di Mattia et al. (2019) reported that water-soluble extracts of crickets showed the highest antioxidant capacities and other insect extracts also showed high antioxidant capacities. From these results, we can assume that when EIP is added to patties, the antioxidant capacity from EIP could be beneficial to pork patties.

Conclusion

This study investigated the effect of adding EIP as meat partial substitute on the physicochemical properties and storage stability of pork patties. With the addition of EIP, the moisture content of the patties decreased while the protein, carbohydrate, fat, and ash contents increased. Additionally, pH and WHC increased, and CL decreased. TPA showed that hardness, chewiness, and gumminess were higher compared to PC and NC, but overall preference decreased. As a result of storage stability with the addition of EIP, TMC, and TBARS were higher compared to PC and NC, while PV was low values. According to the addition of EIP, DPPH radical scavenging activity was higher than NC, and among patties with EIP added, G20 was similar to or higher than PC. Therefore, the addition of EIP as a meat partial substitute showed a positive effect on the physicochemical properties and antioxidant activities of pork patties. Among the 4 species of EIP, G20 was the most promising insect powder to improve the quality of patties and enhance their antioxidant activities. These findings indicated that EIP can serve as meat partial substitute. However, in order to improve storage stability and preference, further research is needed to apply it to patties by reducing the amount of EIP or adding auxiliary ingredients.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) through the High Value-added Food Technology Development Project, funded by Ministry of Agriculture, Food and

Rural Affairs (MAFRA) (Project No. 321028-5). This research was supported by “Regional Innovation Strategy (RIS)” through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (MOE) (2021RIS-001).

Author Contributions

Conceptualization: Choi N. Data curation: Choi N, Park Y. Formal analysis: Choi N, Park S, Oh S, Kim Y, Lim Y. Validation: Park G, Feng X, Choi J. Investigation: Choi N, Jang S, Kim Y, Ahn KS. Writing - original draft: Choi N. Writing - review & editing: Choi N, Park S, Park Y, Park G, Oh S, Kim Y, Lim Y, Jang S, Kim Y, Ahn KS, Feng X, Choi J.

Ethics Approval

The approved consent procedure for sensory evaluation is Institutional Review Board (IRB) of Chungbuk National University (No. CBNU-202302-HR-0017).

References

- Abbasi T, Abbasi SA. 2016. Reducing the global environmental impact of livestock production: The minilivestock option. *J Clean Prod* 112:1754-1766.
- Aiello D, Barbera M, Bongiorno D, Cammarata M, Censi V, Indelicato S, Mazzotti F, Napoli A, Piazzese D, Saiano F. 2023. Edible insects an alternative nutritional source of bioactive compounds: A review. *Molecules* 28:699.
- Akhtar Y, Isman MB. 2018. Insects as an alternative protein source. In *Proteins in food processing*. 2nd ed. Yada RY (ed). Woodhead, Sawston, UK. pp 263-288.
- Alexandratos N, Bruinsma J. 2012. *World agriculture towards 2030/2050: The 2012 revision*. Food and Agriculture Organization of the United Nations, Rome, Italy. p 71.
- Anas M, Ahmad S, Malik A. 2019. *Microbial escalation in meat and meat products and its consequences*. Springer, Cham, Switzerland. pp 29-49.
- AOAC. 1990. *Official methods of analysis*. 15th ed. Association of Official Analytical Chemists, Washington, DC, USA. pp 777-788.
- Aragon-Martinez OH, Martinez-Morales F, González-Chávez MM, Méndez-Gallegos SJ, González-Chávez R, Posadas-Hurtado JC, Isiordia-Espinoza MA. 2023. *Dactylopius opuntiae* [Cockerell] could be a source of antioxidants for the preservation of beef patties. *Insects* 14:811.
- Baigts-Allende D, Sedaghat Doost A, Ramírez-Rodrigues M, Dewettinck K, Van der Meeren P, de Meulenaer B, Tzompasosa D. 2021. Insect protein concentrates from Mexican edible insects: Structural and functional characterization. *LWT-Food Sci Technol* 152:112267.
- Balasundram N, Sundram K, Samman S. 2006. Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses. *Food Chem* 99:191-203.
- Beets WC. 1997. The need for an increased use of small and mini-livestock in integrated smallholder farming systems. *Ecol Food Nutr* 36:237-245.
- Benavente-García O, Castillo J, Marin FR, Ortuño A, Del Río JA. 1997. Uses and properties of *citrus* flavonoids. *J Agric Food Chem* 45:4505-4515.

- Bessa LW, Pieterse E, Marais J, Hoffman LC. 2023. Black soldier fly larvae (*Hermetia illucens*) as a meat replacer in a burger patty. *J Insects Food Feed* 9:1211-1222.
- Bukkens SGF. 1997. The nutritional value of edible insects. *Ecol Food Nutr* 36:287-319.
- Carballo J, Barreto G, Colmenero FJ. 1995. Starch and egg white influence on properties of bologna sausage as related to fat content. *J Food Sci* 60:673-677.
- Choi MH, Kim KH, Yook HS. 2019. Antioxidant activity and quality evaluation of the larvae of *Protaetia brevitarsis* after feeding with Korean *Panax ginseng*. *J Korean Soc Food Sci Nutr* 48:403-409.
- Choi YS, Kim TK, Choi HD, Park JD, Sung JM, Jeon KH, Paik HD, Kim YB. 2017. Optimization of replacing pork meat with yellow worm (*Tenebrio molitor* L.) for frankfurters. *Korean J Food Sci Anim Resour* 37:617-625.
- Churchward-Venne TA, Pinckaers PJM, van Loon JJA, van Loon LJC. 2017. Consideration of insects as a source of dietary protein for human consumption. *Nutr Rev* 75:1035-1045.
- Cruz-López SO, Álvarez-Cisneros YM, Domínguez-Soberanes J, Escalona-Buendía HB, Sánchez CN. 2022. Physicochemical and sensory characteristics of sausages made with grasshopper (*Sphenarium purpurascens*) flour. *Foods* 11:704.
- Damasceno BC, Nakajima M, Taarji N, Kobayashi I, Ichikawa S, Neves MA. 2023. Improvements in visual aspects and chemical, techno-functional and rheological characteristics of cricket powder (*Gryllus bimaculatus*) by solvent treatment for food utilization. *Foods* 12:1422.
- Di Mattia C, Battista N, Sacchetti G, Serafini M. 2019. Antioxidant activities *in vitro* of water and liposoluble extracts obtained by different species of edible insects and invertebrates. *Front Nutr* 6:106.
- Dimitrova B, Gevrenova R, Anklam E. 2007. Analysis of phenolic acids in honeys of different floral origin by solid-phase extraction and high-performance liquid chromatography. *Phytochem Anal* 18:24-32.
- Elsharawy NT, Ahmad AM, Abdelrahman HA. 2018. Quality assessment of nutritional value and safety of different meat. *J Food Microbiol Saf Hyg* 3:1-5.
- Gan J, Zhao M, He Z, Sun L, Li X, Feng Y. 2022. The effects of antioxidants and packaging methods on inhibiting lipid oxidation in deep fried crickets (*Gryllus bimaculatus*) during storage. *Foods* 11:326.
- Gunstone FD, Norris FA. 1983. Oxidation. In *Lipids in foods: Chemistry, biochemistry, and technology*. 1st ed. Gunstone FD, Norris FA (ed). Pergamon Press, Oxford, UK. p 58.
- Hadidi M, Orellana-Palacios JC, Aghababaei F, Gonzalez-Serrano DJ, Moreno A, Lorenzo JM. 2022. Plant by-product antioxidants: Control of protein-lipid oxidation in meat and meat products. *LWT-Food Sci Technol* 169:114003.
- Han X, Li B, Puolanne E, Heinonen M. 2023. Hybrid sausages using pork and cricket flour: Texture and oxidative storage stability. *Foods* 12:1262.
- Ho I, Peterson A, Madden J, Huang E, Amin S, Lammert A. 2022. Will it cricket? Product development and evaluation of cricket (*Acheta domesticus*) powder replacement in sausage, pasta, and brownies. *Foods* 11:3128.
- Honikel KO. 2004. Water-holding capacity of meat. In *Muscle development of livestock animals: Physiology, genetics and meat quality*. te Pas MFW, Everts ME, Haagsman HP (ed). CABI, Wallingford, UK. pp 389-400.
- Honikel KO. 2008. Moisture and water-holding capacity. 1st ed. CRC Press, Boca Raton, FL, USA. pp 335-354.
- Juntachote T, Berghofer E, Siebenhandl S, Bauer F. 2007. Antioxidative effect of added dried Holy basil and its ethanolic extracts on susceptibility of cooked ground pork to lipid oxidation. *Food Chem* 100:129-135.
- Kim HW, Setyabrata D, Lee YJ, Jones OG, Kim YHB. 2016. Pre-treated mealworm larvae and silkworm pupae as a novel protein ingredient in emulsion sausages. *Innov Food Sci Emerg Technol* 38:116-123.
- Kim HW, Setyabrata D, Lee Y, Jones OG, Kim YHB. 2017. Effect of house cricket (*Acheta domesticus*) flour addition on

- physicochemical and textural properties of meat emulsion under various formulations. *J Food Sci* 82:2787-2793.
- Kim TK, Kim YJ, Kim J, Yun HJ, Kang MC, Choi YS. 2022. Effect of grafted insect protein with palatinose on quality properties of phosphate-free meat emulsion. *Foods* 11:3354.
- Kim TK, Yong HI, Jung S, Sung JM, Jang HW, Choi YS. 2021. Physicochemical and textural properties of emulsions prepared from the larvae of the edible insects *Tenebrio molitor*, *Allomyrina dichotoma*, and *Protaetia brevitarsis seoulensis*. *J Anim Sci Technol* 63:417-425.
- Kipkoech C. 2023. Beyond proteins: Edible insects as a source of dietary fiber. *Polysaccharides* 4:116-128.
- Klunder HC, Wolkers-Rooijackers J, Korpela JM, Nout MJR. 2012. Microbiological aspects of processing and storage of edible insects. *Food Control* 26:628-631.
- Kurdi P, Chaowiwat P, Weston J, Hansawasdi C. 2021. Studies on microbial quality, protein yield, and antioxidant properties of some frozen edible insects. *Int J Food Sci* 2021:5580976.
- Lange KW, Nakamura Y. 2021. Edible insects as future food: Chances and challenges. *J Future Foods* 1:38-46.
- Lemke B, Siekmann L, Grabowski NT, Plötz M, Krischek C. 2023. Impact of the addition of *Tenebrio molitor* and *Hermetia illucens* on the physicochemical and sensory quality of cooked meat products. *Insects* 14:487.
- Love JD, Pearson AM. 1971. Lipid oxidation in meat and meat products: A review. *J Am Oil Chem Soc* 48:547-549.
- Lu M, Zhu C, Smetana S, Zhao M, Zhang H, Zhang F, Du Y. 2024. Minerals in edible insects: A review of content and potential for sustainable sourcing. *Food Sci Hum Wellness* 13:65-74.
- Malm M, Liceaga AM. 2021. Physicochemical properties of chitosan from two commonly reared edible cricket species, and its application as a hypolipidemic and antimicrobial agent. *Polysaccharides* 2:339-353.
- Megido RC, Gierts C, Blecker C, Brostaux Y, Haubruge É, Alabi T, Francis F. 2016. Consumer acceptance of insect-based alternative meat products in Western countries. *Food Qual Prefer* 52:237-243.
- Miyanoshta A, Hara S, Sugiyama M, Asaoka A, Taniai K, Yukuhiro F, Yamakawa M. 1996. Isolation and characterization of a new member of the insect defensin family from a beetle, *Allomyrina dichotoma*. *Biochem Biophys Res Commun* 220:526-531.
- Montowska M, Kowalczewski PŁ, Rybicka I, Fornal E. 2019. Nutritional value, protein and peptide composition of edible cricket powders. *Food Chem* 289:130-138.
- Ooninx DGAB, van Itterbeeck J, Heetkamp MJW, van den Brand H, van Loon JJA, van Huis A. 2010. An exploration on greenhouse gas and ammonia production by insect species suitable for animal or human consumption. *PLOS ONE* 5:e14445.
- Pietrasik Z. 2003. Binding and textural properties of beef gels processed with κ -carrageenan, egg albumin and microbial transglutaminase. *Meat Sci* 63:317-324.
- Pinter R, Molnar E, Hussein KN, Toth A, Friedrich L, Pasztor-Huszar K. 2021. Effect of refrigerated storage on the technological characteristics of meat stick made of insect and pork: Alternative burger meat. *Prog Agric Eng Sci* 16:117-125.
- Premalatha M, Abbasi T, Abbasi T, Abbasi SA. 2011. Energy-efficient food production to reduce global warming and ecodegradation: The use of edible insects. *Renew Sustain Energy Rev* 15:4357-4360.
- Rumpold BA, Schlüter OK. 2013. Nutritional composition and safety aspects of edible insects. *Mol Nutr Food Res* 57:802-823.
- Sagisaka A, Miyanoshta A, Ishibashi J, Yamakawa M. 2001. Purification, characterization and gene expression of a glycine and proline-rich antibacterial protein family from larvae of a beetle, *Allomyrina dichotoma*. *Insect Mol Biol* 10:293-302.
- Simic MG, Taylor KA. 1987. Free radical mechanisms of oxidation reactions. In *Warmed-over flavor of meat*. St. Angelo AJ, Bailey ME (ed). Academic Press, London, UK. pp 69-72.

- Song MH, Han MH, Lee S, Kim ES, Park KH, Kim WT, Choi JY. 2017. Growth performance and nutrient composition in the white-spotted flower chafer, *Protaetia brevitarsis* (Coleoptera: Scarabaeidae) fed agricultural by-product, soybean curd cake. *J Life Sci* 27:1185-1190.
- Suh HJ, Kang SC. 2012. Antioxidant activity of aqueous methanol extracts of *Protaetia brevitarsis* Lewis (Coleoptera: Scarabaeidae) at different growth stages. *Nat Prod Res* 26:510-517.
- Szmańko T, Lesiów T, Górecka J. 2021. The water-holding capacity of meat: A reference analytical method. *Food Chem* 357:129727.
- Torres-Castillo JA, Olazarán-Santibáñez FE. 2023. Insects as source of phenolic and antioxidant entomochemicals in the food industry. *Front Nutr* 10:1133342.
- Turgut SS, Işıklı F, Soyer A. 2017. Antioxidant activity of pomegranate peel extract on lipid and protein oxidation in beef meatballs during frozen storage. *Meat Sci* 129:111-119.
- van Broekhoven S, Oonincx DGAB, van Huis A, van Loon JJA. 2015. Growth performance and feed conversion efficiency of three edible mealworm species (Coleoptera: Tenebrionidae) on diets composed of organic by-products. *J Insect Physiol* 73:1-10.
- van der Weele C, Feindt P, van der Goot AJ, van Mierlo B, van Boekel M. 2019. Meat alternatives: An integrative comparison. *Trends Food Sci Technol* 88:505-512.
- van Huis A, Tomberlin JK. 2017. Insects as food and feed: From production to consumption. Wageningen Academic, Wageningen, The Netherlands. pp 24-41.
- van Huis A, Van Itterbeeck J, Klunder H, Mertens E, Halloran A, Muir G, Vantomme P. 2013. Edible insects: Future prospects for food and feed security. Food and Agriculture Organization of the United Nations, Rome, Italy. pp 1-156.
- Wang D, Bai Y, Li J, Zhang C. 2004. Nutritional value of the field cricket (*Gryllus testaceus* Walker). *Insect Sci* 11:275-283.
- Wittkopp PJ, Beldade P. 2009. Development and evolution of insect pigmentation: Genetic mechanisms and the potential consequences of pleiotropy. *Semin Cell Dev Biol* 20:65-71.
- World Health Organization [WHO]. 2007. Protein and amino acid requirements in human nutrition: Report of a joint FAO/WHO/UNU expert consultation. WHO, Geneva, Switzerland.
- Wu G, Fanzo J, Miller DD, Pingali P, Post M, Steiner JL, Thalacker-Mercer AE. 2014. Production and supply of high-quality food protein for human consumption: Sustainability, challenges, and innovations. *Ann NY Acad Sci* 1321:1-19.
- Wu J. 2022. Emerging sources and applications of alternative proteins: An introduction. Academic Press, Cambridge, MA, USA. pp 1-15.
- Yong HI, Kim TK, Cha JY, Lee JH, Kang MC, Jung S, Choi YS. 2023. Effects of edible insect extracts on the antioxidant, physiochemical, and microbial properties of Tteokgalbi during refrigerated storage. *Food Biosci* 52:102377.
- Yoon HS, Lee CS, Lee SY, Choi CS, Lee IH, Yeo SM, Kim HR. 2003. Purification and cDNA cloning of inducible antibacterial peptides from *Protaetia brevitarsis* (Coleoptera). *Arch Insect Biochem Physiol* 52:92-103.
- Yoshikawa K, Umetsu K, Shinzawa H, Yuasa I, Maruyama K, Ohkura T, Yamashita K, Suzuki T. 1999. Determination of carbohydrate-deficient transferrin separated by lectin affinity chromatography for detecting chronic alcohol abuse. *FEBS Lett* 458:112-116.
- Zielińska E, Baraniak B, Karaś M, Rybczyńska K, Jakubczyk A. 2015. Selected species of edible insects as a source of nutrient composition. *Food Res Int* 77:460-466.