

ARTICLE

Determination of Flunixin and 5-Hydroxy Flunixin Residues in Livestock and Fishery Products Using Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)

 Dahae Park^{1,#}, Yong Seok Choi², Ji-Young Kim^{1,*}, Jang-Duck Choi¹, and Gui-Im Moon¹
 OPEN ACCESS

Received January 29, 2024

Revised March 12, 2024

Accepted March 12, 2024

¹Pesticides & Veterinary Drug Residues Division, National Institute of Food & Drug Safety Evaluation, Cheongju 28159, Korea

²College of Pharmacy, Dankook University, Cheonan 31116, Korea

***Corresponding author** : Ji-Young Kim
 Pesticides & Veterinary Drug Residues
 Division, National Institute of Food & Drug
 Safety Evaluation, Cheongju 28159, Korea
 Tel: +82-43-719-4218
 Fax: +82-43-719-4200
 E-mail: jykim98@korea.kr

#Current affiliation: Department of
 Environmental Science, Kangwon National
 University, Chuncheon 24341, Korea

***ORCID**
 Dahae Park
<https://orcid.org/0000-0002-1774-7413>
 Yong Seok Choi
<https://orcid.org/0000-0001-6740-3160>
 Ji-Young Kim
<https://orcid.org/0000-0002-6330-6731>
 Jang-Duck Choi
<https://orcid.org/0000-0002-8576-2754>
 Gui-Im Moon
<https://orcid.org/0000-0002-3726-6748>

Abstract Flunixin is a veterinary nonsteroidal anti-inflammatory agent whose residues have been investigated in their original form within tissues such as muscle and liver. However, flunixin remains in milk as a metabolite, and 5-hydroxy flunixin has been used as the primary marker for its surveillance. This study aimed to develop a quantitative method for detecting flunixin and 5-hydroxy flunixin in milk and to strengthen the monitoring system by applying to other livestock and fishery products. Two different methods were compared, and the target compounds were extracted from milk using an organic solvent, purified with C₁₈, concentrated, and reconstituted using a methanol-based solvent. Following filtering, the final sample was analyzed using liquid chromatography-tandem mass spectrometry. Method 1 is environmentally friendly due to the low use of reagents and is based on a multi-residue, multi-class analysis method approved by the Ministry of Food and Drug Safety. The accuracy and precision of both methods were 84.6%–115% and 0.7%–9.3%, respectively. Owing to the low matrix effect in milk and its convenience, Method 1 was evaluated for other matrices (beef, chicken, egg, flatfish, and shrimp) and its recovery and coefficient of variation are sufficient according to the Codex criteria (CAC/GL 71-2009). The limits of detection and quantification were 2–8 and 5–27 µg/kg for flunixin and 2–10 and 6–33 µg/kg for 5-hydroxy flunixin, respectively. This study can be used as a monitoring method for a positive list system that regulates veterinary drug residues for all livestock and fisheries products.

Keywords flunixin, veterinary drug, residues, food safety, liquid chromatography-tandem mass spectrometry (LC-MS/MS)

Introduction

Non-steroidal anti-inflammatory drugs (NSAIDs) are extensively used as veterinary drugs to prevent inflammation or to treat allergies, fever, and pain. Flunixin is the only

NSAID in the U.S. labeled for use in beef and dairy cattle to treat respiratory tract disease and mastitis (Smith et al., 2008). However, it has adverse side effects, such as gastrointestinal bleeding, intestinal ulceration, aplastic anemia, and the inhibition of platelet aggregation (Kari et al., 1995). Daeseleire et al. (2003) proposed a withdrawal time of 12 h owing to the presence of trace residues of 5-hydroxy flunixin after 10 h. According to the U.S. Food and Drug Administration (FDA), the withdrawal times of flunixin are 24 h at a dose of 2.2 mg/kg and 12 h at 1.1 mg/kg for intravenous administration. Moreover, the slaughter and milk withdrawal times are 4 d and 36 h, respectively (FDA, 2004). And tolerance/Safe Level of 5-hydroxy flunixin is 2 µg/kg in milk (FDA, 2020) and the final acceptable daily intake for carcinogenicity was established at 0.72 µg/kg bw/day for human exposure (FDA, 2017).

Flunixin is easily metabolized, and approximately 90% is excreted in the urine (Levionnois et al., 2018). 5-Hydroxy flunixin, one of its metabolites, is a marker in bovine milk, and the residue of flunixin in bovine muscle can be identified in its original form (Chen et al., 2019; Jedziniak et al., 2016; Ngoh et al., 2003). In 2011, the U.S. FDA presented plans to surveil 5-hydroxy flunixin in milk beginning with residue violations of flunixin in dairy cows. This event increased concerns about the presence of drug residues in milk. The European Union (EU), under Community Regulation 2377/90, set the maximum residue level (MRL) for flunixin and its metabolite (5-hydroxy flunixin) at <40 µg/kg in milk produced for human consumption, while the MRL of flunixin is 20 µg/kg in bovine muscle (EU, 1990). In South Korea, flunixin in animal-based foods is regulated by the MRL (MFDS, 2023a). Flunixin residue is defined as its original form, and the MRLs are 0.02, 0.05, 0.01, and 0.02 mg/kg for beef, pork, and horse muscles and milk, respectively. In 2022, the definition was amended for 5-hydroxy flunixin in milk by the Ministry of Food and Drug Safety (MFDS). Therefore, a method for analyzing 5-hydroxy flunixin in milk is required to protect public health.

5-Hydroxy flunixin has a fluorescent property and can be detected at excitation and emission wavelengths of 360 and 495 nm, respectively, using fluorescence detection (FLD; Gallo et al., 2010). Analyzing residues in agricultural products is challenging due to the matrix effects caused by co-extractives, such as pigments, fatty acids, sugars, and other interrupting substances (Fialkov et al., 2007). For the treatment of food samples, a clean-up process is required to remove interfering substances. Solid-phase extraction (SPE) using OASIS HLB, Strata X, and ABN has been used for deproteinization with trichloroacetic acid; however, it causes the loss of certain target compounds of NSAIDs (Malone et al., 2009). Plasma samples have been acidified using inorganic acids, such as hydrochloric acid, to precipitate proteins (Jedziniak et al., 2016). Quick, Easy, Cheap, Effective, Rugged, and Safe (QuEChERS) is the most common method for food sample preparation and is widely used for multi-class, multi-residue analyses (Hajrulai-Musliu et al., 2021; Jang et al., 2022; Wilkowska and Biziuk, 2011). To determine flunixin and 5-hydroxy flunixin contents, high-performance liquid chromatography (HPLC)-ultraviolet spectrometry (Asea et al., 2001; Jedziniak et al., 2016), liquid chromatography (LC)-tandem mass spectrometry (MS/MS; Daeseleire et al., 2003; Krueve et al., 2008; Malone et al., 2009), and other methods have been used. Flunixin is a substance classified and managed as group B (pharmacologically active substances authorised for use in food-producing animals), and the FLD method was suggested in the European Council decision 2002/657/EC (European Commission, 2002). This study attempted to analyze using QuEChERS, which has a simple pretreatment process, and LC-MS/MS, which can analyze various residual animal medicines.

The number of regulated veterinary drugs has continually increased, and the surveillance of residues is becoming stricter with systems such as the positive list system (PLS; MFDS, 2023a). The PLS regulates all veterinary drugs in all types of animal-based foods at a specific concentration (0.01 mg/kg). Therefore, this study aimed to develop a quantitative method for the detection of 5-hydroxy flunixin in milk. By applying 5-hydroxy flunixin, which is not included in the multi-class, multi-

residue analysis method established by the MFDS, the applicability was assessed to improve monitoring efficiency and prepare for PLS.

Materials and Methods

Materials

Standards (flunixin and 5-hydroxy flunixin) were obtained from Toronto Research Chemicals (TRC; Windsor, CT, USA) and stored at -80°C . HPLC-grade acetonitrile, dichloromethane, *n*-hexane, methanol, and water were purchased from Merck (Rahway, NJ, USA). The ethylenediaminetetraacetic acid (EDTA) disodium salt solution and formic acid were purchased from Sigma-Aldrich (St. Louis, MO, USA). C_{18} sorbent was obtained from CHROMAtific (Heidenrod, Germany), and polytetrafluoroethylene (PTFE) and nylon syringe filters were obtained from Teknokroma (Barcelona, Spain). All standard solutions were individually prepared at 1,000 mg/L in methanol and stored at -80°C .

Instrumental conditions

The analysis was performed using a Shimadzu LCMS-8060 system (Shimadzu, Osaka, Japan) connected to a Waters X select HSS C_{18} column (150 mm \times 2.1 mm id, 3.5 μm). The mobile phases A and B were 0.1% formic acid in water (A) and 0.1% formic acid in acetonitrile (B), respectively. The LC gradient conditions were as follows: 5% B (0 min), hold at 5% B (0.5 min), increase to 60% B (5.5 min), increase to 100% B (6.0 min), hold at 100% B (10.0 min), rapidly decrease B to 5% (10.2 min), and hold for stabilization (12.00 min). The flow rate, injection volume, and temperature were 0.3 mL/min, 5 μL , and 40°C , respectively. The ESI (electrospray ionization) mode was used, the capillary temperature and voltage were 500°C and 3.6 kV, respectively, and argon gas was used as the impact gas. The specific multiple-reaction monitoring (MRM) conditions are listed in Table 1.

Sample preparation

Method 1

The applicability of the multi-residue, multi-class analysis method developed by the MFDS (2023b) was confirmed as follows. Homogenized samples (2 g) were prepared in a 50-mL centrifuge tube, and 10 mL of water: acetonitrile (1:4, v/v) was added to the sample. The mixture was shaken for 5 min and centrifuged at $4,700\times g$ for 10 min at 4°C . The supernatant was decanted into a second 50-mL centrifuge tube containing 500 mg of C_{18} and 10 mL of *n*-hexane saturated with acetonitrile. The mixture was shaken for 10 min and centrifuged at $4,700\times g$ for 5 min at 4°C . Acetonitrile extract (5 mL) was transferred to a 15-mL centrifuge tube and concentrated at 40°C to dry under nitrogen gas. The concentrate was dissolved with 1 mL of water: methanol (1:1, v/v) and filtered through a 0.2- μm PTFE syringe filter. The final sample was analyzed

Table 1. Multiple reaction monitoring (MRM) conditions

Target	Retention time (min)	Ionization	Exact mass (g/mol)	Precursor ion (m/z)	Product ion (m/z)	Collision energy (eV)
Flunixin	7.19	$[\text{M}+\text{H}]^+$	296.1	297.1	279.0, 263.9, 281.0	22, 33, 14
5-Hydroxy flunixin	6.78	$[\text{M}+\text{H}]^+$	312.2	312.9	279.9, 295.0	34, 23

using LC-MS/MS.

Method 2

The analysis methods developed in Korea by Shin and Choi (2022) for flunixin and 5-hydroxy flunixin were compared with Method 1. Milk samples (2 g) were prepared in a 50-mL centrifuge tube. EDTA solution (100 μ L) was added, and the mixture was shaken for 3 min. Subsequently, 8.5 mL of acetonitrile containing 0.1% formic acid and 1.5 mL of dichloromethane were added. The mixture was shaken for 15 min and centrifuged at 4,500 \times g for 10 min at 4°C. The supernatant was decanted into a 15-mL centrifuge tube containing 150 mg of C₁₈. The mixture was shaken for 10 min and centrifuged at 4,500 \times g for 10 min at 4°C. The supernatant was transferred to a second 15-mL centrifuge tube and concentrated to dryness under gentle nitrogen gas at 40°C. The concentrate was reconstituted with methanol (1 mL), sonicated for 5 min, and vigorously mixed for 1 min. After centrifuging for 10 min (at 16,000 \times g and 4°C) and filtering through a 0.2- μ m nylon syringe filter, the final sample was analyzed using LC-MS/MS.

Method confirmation

According to the Codex guidelines (CAC/GL 71-2009), the method was validated for linearity, accuracy, precision, and detection limits (Codex Alimentarius, 2014). Linearity was represented as the coefficient of determination (r^2), and the five points of the calibration curve were at 2.5, 5, 10, 20, and 40 μ g/kg. Accuracy and precision were confirmed by repeating the analysis seven times at three level and were represented as the recovery and coefficient of variation (CV), respectively. The detection limits were expressed as limits of detection (LOD) and quantification (LOQ), calculated as 3.3 and 10 times the slope SD, respectively. To confirm inter-lab reproducibility, method validation was performed in two other laboratories.

The matrix effects of the two methods were estimated by calculating the calibration slope (Kwon et al., 2012). Target compounds were spiked into each matrix at six points within the range of 0–80 μ g/kg to prepare calibration curves and calculate slope factors. Then, the slope factor was obtained from the calibration curve created by spiking the same amount into the solvent. The matrix effects (%) of the analytes were calculated using the following Eq. (1):

$$\%ME = \left[\frac{\text{Slope of the matrix-matched calibration}}{\text{Slope of the reagent-only calibration}} \right] \times 100. \quad (1)$$

Ferrer et al. (2011) classified the matrix effects as no effect (<20%), medium effect (20%–50%), or strong effect (>50%). Matrix effects were evaluated by referring to the classification.

Results and Discussion

Accuracy and precision of the two methods

This study presented a quantitative analysis method for 5-hydroxy flunixin residues in milk using flunixin and compared the two methods to increase monitoring efficiency (Tables 2 and 3). Both methods exhibited good linearity (>0.99). In Method 1, the recoveries and CVs were 110%–115% and 2.2%–5.4% for flunixin and 94.0%–108% and 3.1%–9.3% for 5-hydroxy flunixin, respectively. In Method 2, the recoveries and CVs were 97.2%–99.6% and 2.2%–3.9% for flunixin and 84.6%–101% and 0.7%–8.4% for 5-hydroxy flunixin, respectively. The results of both methods met the performance criteria

Table 2. Inter-lab and intra-lab validation of Method 1 for quantification of flunixin and 5-hydroxy flunixin in milk

Target	Linearity (r ²)	Fortification (µg/kg)	Inter-lab		Inter-lab		LOD (µg/kg)	LOQ (µg/kg)	
			Recovery (%)	CV (%)	Recovery (%)				CV (%)
					Lab A	Lab B			
Flunixin	0.9943	10	112	3.7	85.0	74.5	21	4	15
		20	115	5.4	82.4	76.1	22		
		40	110	2.2	79.8	94.0	8.9		
5-Hydroxy flunixin	0.9979	10	108	3.1	82.7	73.6	20	5	15
		20	99.6	5.9	82.7	71.2	18		
		40	94.0	9.3	77.4	74.8	14		

CV, coefficient of variation; LOD, limits of detection; LOQ, limits of quantification.

Table 3. Inter-lab validation of Method 2 for quantification of flunixin and 5-hydroxy flunixin in milk

Target	Linearity (r ²)	Fortification (µg/kg)	Recovery (%)	Intra-lab CV (%)
Flunixin	0.9991	10	99.6	3.0
		20	99.1	3.9
		40	97.2	2.2
5-Hydroxy flunixin	0.9993	10	101	4.1
		20	93.6	0.7
		40	84.6	8.4

CV, coefficient of variation.

set by Codex for quantitative analytical methods for veterinary drug residues in foods. Therefore, Methods 1 and 2 were suitable for identifying and quantifying flunixin residues in milk products.

However, Method 1 is an easy and efficient method owing to its simpler sample preparation compared with Method 2. Method 1 is a multi-class, multi-residue analysis method used in Korea to determine the residues of 157 veterinary drugs in livestock products. In addition, Method 1 does not interrupt the existing multi-class, multi-residue analysis method. The structures of the analytes and their selectivities are shown in Figs. 1 and 2.

Milk analysis

In Korea, the MRL of milk is 20 µg/kg for 5-hydroxy flunixin. In Method 1, the LODs and LOQs were 4 and 15 µg/kg for flunixin and 5 and 15 µg/kg for 5-hydroxy flunixin, respectively. In Method 2, the LODs and LOQs were 2 and 5 µg/kg for flunixin and 3 and 9 µg/kg for 5-hydroxy flunixin, respectively. Both methods can be used to quantify concentrations lower than the MRL. The matrix effects of flunixin and 5-hydroxy flunixin were -15% and -24.2% in Method 1 and 78.5% and 65.9% in Method 2, respectively (Table 4). Method 1 had no matrix effect on flunixin and a medium matrix effect on 5-hydroxy flunixin, whereas Method 2 had a strong matrix effect on both compounds. Considering the importance of quantifying 5-hydroxy flunixin in milk, both methods satisfied the Codex validation criteria. However, Method 1 has a simpler sample preparation and less matrix effect, and can be applied to multi-class, multi-residue analysis. Therefore, the present study proceeded with additional validation using other livestock and fishery products.




Fig. 1. Chemical structures of flunixin and 5-hydroxy flunixin.




Fig. 2. Chromatograms of flunixin and 5-hydroxyflunixin in standard solutions ($0.2 \mu\text{g/mL}$) and spiked samples (milk, 0.02 mg/kg) using Methods 1 and 2.

The inter-lab validation of Method 1 was assessed (Table 2). For flunixin and 5-hydroxy flunixin, the recoveries were 74.5%–94.0% and 71.2%–94.0%, and the inter-lab CVs were 8.9%–22% and 14%–20%, respectively. Method 1 was validated using intra- and inter-lab validation results that met the Codex guidelines (CAC/GL 71-2009). According to the guidelines, the recovery should be within the range of 60%–120% at $1\text{--}10 \mu\text{g/kg}$ and 70%–120% at $10\text{--}100 \mu\text{g/kg}$. And the

Table 4. Matrix effect (ME) of flunixin and 5-hydroxy flunixin for 4 livestock and 2 fishery products by Method 1

Variable	Flunixin			5-Hydroxy flunixin		
	Slope	ME%	Class	Slope	ME%	Class
Solvent	993,371.5			136,638.3		
Milk	844,357.0	-15.0	No effect	103,543.5	-24.2	Medium
Beef	187,881.6	-81.1	Strong	4,302.4	-96.9	Strong
Chicken	1,665,518.6	67.7	Strong	258,806.7	89.4	Strong
Egg	1,966,821.5	98.0	Strong	227,537.8	66.5	Strong
Flatfish	1,801,178.7	81.3	Strong	275,312.2	101.5	Strong
Shrimp	1,805,141.4	81.7	Strong	223,522.6	63.6	Strong

intra-lab and inter-lab CV should be less than 30% and 45% at 1–10 µg/kg and 20% and 32% at 10–100 µg/kg, respectively. In previous studies for milk or bovine muscle, ultrasound-assisted extraction showed 0.5 µg/kg of CC α for both flunixin and 5-hydroxy flunixin (Daeseleire et al., 2003) and 22.85 and 27.28 µg/kg of CC α and CC β for flunixin (Lugoboni et al., 2014), respectively. Extraction with acetonitrile and wash with *n*-hexane showed 45.04 and 50.08 µg/kg of CC α and CC β for 5-hydroxy flunixin, respectively (Malone et al., 2009). The present method used sorbents and washed the extract with *n*-hexane, as described in Method 1. These processes are necessary to improve purification, and filtration is required to eliminate particulates from the final sample (Malone et al., 2009). Acidic organic solvents, such as hydrochloric acid, were introduced for extraction, and the LODs were 2.98 µg/kg of flunixin in bovine muscle and 0.78 µg/L of 5-hydroxy flunixin in milk (Chen et al., 2019). For the bovine muscle sample, potassium acetate buffer (adjusted to pH 4.5 with acetic acid) and enzymatic hydrolysis using β -glucuronidase were also used with 6 mg/kg of LOD (Asea et al., 2001). Method 1 of the present study enabled the quantification of flunixin residues down to the MRL using the usual QuEChERS method, without the use of acids or buffers.

Application to other livestock and fishery products

Method 1 was validated using other animal-based food products (Table 5). Five livestock and fishery products (beef, chicken, eggs, flatfish, and shrimp) were evaluated. All matrices showed good linearity (>0.99), and the recoveries and CVs were 82.4%–110% and 1.2%–8.9% for flunixin and 83.3%–106% and 2.2%–20% for 5-hydroxy flunixin, respectively. The matrix effects of Method 1 are listed in Table 4. For all matrices, except milk, Method 1 was estimated to have a strong matrix effect. Flunixin exhibited the strongest matrix effect in eggs, followed by shrimp, flatfish, beef, and chicken. 5-Hydroxy flunixin showed the strongest matrix effect in flatfish, followed by beef, chicken, eggs, and shrimp. When using LC-MS/MS, the matrix effect depends on the sample properties, ionization interface, mobile phase additives, stationary phase, and other characteristics. The ionization intensity can be reduced by nonvolatile substances via increasing the viscosity and surface tension and decreasing the solvent evaporation rate (Silvestro et al., 2013). Therefore, the strong matrix effect is interpreted as a phenomenon caused by ion suppression. Simple purification with *n*-hexane and C₁₈ is not sufficient to reduce matrix effects, however, this can be addressed using a matrix-matched calibration.

Recently, the strict monitoring of veterinary drug residues has been emphasized with the introduction of the PLS system in Korea. Consistent with this situation, analysis methods for all animal-based foods are required. A previous FDA study

Table 5. Validation of other livestock and fishery products using Method 1

Matrix	Target	Linearity (r^2)	Fortification ($\mu\text{g}/\text{kg}$)	Recovery (%)	CV (%)	LOD ($\mu\text{g}/\text{kg}$)	LOQ ($\mu\text{g}/\text{kg}$)
Beef	Flunixin	0.9952	10	93.1	6.9	8	27
			20	82.4	7.7		
			40	88.4	2.3		
	5-Hydroxy flunixin	0.9928	10	83.3	20	10	33
			20	93.0	16		
			40	100	18		
Chicken	Flunixin	0.9981	5	97.5	8.9	2	7
			10	90.6	1.7		
			20	99.1	4.5		
	5-Hydroxy flunixin	0.9904	5	100	9.7	5	16
			10	88.4	4.5		
			20	88.0	8.4		
Egg	Flunixin	0.9992	5	104	1.2	3	8
			10	103	4.5		
			20	108	2.8		
	5-Hydroxy flunixin	0.9955	5	104	6.8	3	11
			10	102	8.1		
			20	103	2.2		
Flatfish	Flunixin	0.9984	5	105	7.9	2	5
			10	110	2.4		
			20	96.6	2.4		
	5-Hydroxy flunixin	0.9989	5	98.7	8.3	2	6
			10	96.1	4.8		
			20	95.6	7.1		
Shrimp	Flunixin	0.9967	5	106	5.5	2	6
			10	107	6.5		
			20	99.5	7.2		
	5-Hydroxy flunixin	0.9989	5	101	7.2	2	7
			10	106	7.7		
			20	92.8	8.5		

CV, coefficient of variation; LOD, limits of detection; LOQ, limits of quantification.

optimized a multi-class, multi-residue LC-MS/MS method to determine 25 veterinary drug residues (Clark et al., 2011). In the abovementioned method, acetonitrile was added to extract target compounds and precipitate protein, and SPE using OASIS[®] HLB was also employed; however, this method is only for milk and requires additional confirmation to apply other animal-based food. Food analysis using SPE not only has the disadvantage of clogging the adsorbent but also requires conditioning and cleaning steps for activation.

The detection limits of Method 1 were assessed for six animal-based food products (milk, beef, chicken, egg, flatfish, and

shrimp) and are described in Tables 2 and 5. The LODs and LOQs were 2–8 and 5–27 µg/kg for flunixin and 2–10 and 6–33 µg/kg for 5-hydroxy flunixin, respectively. However, Method 1 showed poor recovery of 5-hydroxy flunixin from pork muscle and eels. Recoveries and CVs of flunixin were 96.0%–98.9% and 2.9%–4.1% at 25, 50, and 100 µg/kg in pork and 91.0%–99.0% and 1.5%–9.1% at 5, 10, and 20 µg/kg in eel, respectively. In contrast, the recovery of the same concentration of 5-hydroxy flunixin from pork and eel was <40%. Therefore, Method 1 could be used for the quantification of flunixin or the identification of 5-hydroxy flunixin in pork and eels in a limited manner.

Most studies of flunixin residues have been conducted on beef or milk (Daeseleire et al., 2003; Douglas et al., 2012; Gallo et al., 2010; Kissell et al., 2013; Malone et al., 2009) and only a few have investigated swine and chicken (Chou et al., 2022; Liu et al., 2015). Flunixin is approved for swine at a dosage of 2.2 mg/kg with a withdrawal time of 12 d (Sidhu et al., 2017). In a previous flunixin study, recovery and repeatability were 76.7%–86.8% and 4.1%–12.5% in swine tissues and 73.8%–84.0% and 4.1%–9.9% in chicken tissues, respectively, and the LOQ ranged from 0.05 to 0.50 µg/kg in both matrices (Liu et al., 2015). However, the previous study was performed only for flunixin and used hydrolysis with hydrochloric acid and heating. Although the LOQs in the present study were higher than those of previous studies, the method is beneficial in terms of its simple process and application to other matrices for two compounds (flunixin and 5-hydroxy flunixin) in livestock and fishery products.

Conclusion

This study assessed the linearity, accuracy, precision, detection limits, and matrix effect of two methods for the analysis of flunixin and 5-hydroxy flunixin in milk. Two methods involve QuEChERS-based extraction and purification followed by analysis by LC-MS/MS. Both methods can quantify the two compounds at a lower level than their MRL in milk. Method 1 is the existing multi-class, multi-residue method used in Korea, and Method 2 is a little more complicated than Method 1, including the use of EDTA, formic acid, and dichloromethane. Applicability of Method 1 to four livestock and two fishery products (milk, beef, chicken, egg, shrimp, and flatfish) has been confirmed. The results of this study should improve the efficiency of the monitoring system and emphasize the importance of improving multi-class, multi-residue analysis methods. In future studies, improvements through additional purification processes are needed to obtain low detection limits and MEs even in simultaneous analysis methods.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This study was supported by a grant (22191MFDS317) from the Ministry of Food and Drug Safety in 2022.

Author Contributions

Conceptualization: Choi YS, Kim JY, Choi JD, Moon GI. Data curation: Park D, Choi YS. Formal analysis: Park D, Choi YS. Methodology: Choi YS. Validation: Park D, Choi YS. Investigation: Park D, Choi YS. Writing - original draft: Park D.

Writing - review & editing: Park D, Choi YS, Kim JY, Choi JD, Moon GI.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Asea PA, Patterson JR, Korsrud GO, Dowling PM, Boison JO. 2001. Determination of flunixin residues in bovine muscle tissue by liquid chromatography with UV detection. *J AOAC Int* 84:659-665.
- Chen X, Peng S, Liu C, Zou X, Ke Y, Jiang W. 2019. Development of an indirect competitive enzyme-linked immunosorbent assay for detecting flunixin and 5-hydroxyflunixin residues in bovine muscle and milk. *Food Agric Immun* 30:320-332.
- Chou WC, Tell LA, Baynes RE, Davis JL, Maunsell FP, Riviere JE, Lin Z. 2022. An interactive generic physiologically based pharmacokinetic (igPBPK) modeling platform to predict drug withdrawal intervals in cattle and swine: A case study on flunixin, florfenicol, and penicillin G. *Toxicol Sci* 188:180-197.
- Clark SB, Storey JM, Turnipseed SB. 2011. Optimization and validation of multi-class, multi-residue LC-MS/MS screening and confirmation method for drug residues in milk. *J AOAC Int* 94:1-27.
- Codex Alimentarius. 2014. Guidelines for the design and implementation of national regulatory food safety assurance programme associated with the use of veterinary drugs in food producing animals. CAC/GL 71-2009. Available from: <https://www.fao.org/fao-who-codexalimentarius/codex-texts/guidelines/en/>. Accessed at Jan 24, 2024.
- Daeseleire E, Mortier L, De Ruyck H, Geerts N. 2003. Determination of flunixin and ketoprofen in milk by liquid chromatography–tandem mass spectrometry. *Anal Chim Acta* 488:25-34.
- Douglas D, Banaszewski K, Juskelis R, Al-Taher F, Chen Y, Cappozzo J, McRobbie L, Salter RS. 2012. Validation of a rapid lateral flow test for the simultaneous determination of β -lactam drugs and flunixin in raw milk. *J Food Prot* 75:1270-1277.
- European Commission. 2002. 2002/657/EC: Commission decision of 12 August 2002 implementing council directive 96/23/EC concerning the performance of analytical methods and the interpretation of results (text with EEA relevance) (notified under document number C(2002) 3044). Available from: <https://op.europa.eu/en/publication-detail/-/publication/ed928116-a955-4a84-b10a-cf7a82bad858/language-en>. Accessed at Jan 24, 2024.
- European Union [EU]. 1990. Council regulation (EEC) no. 2377/90 laying down a community procedure for the establishment of maximum residue limits of veterinary medicinal products in foodstuffs of animal origin. Available from: <https://www.fao.org/faolex/results/details/en/c/LEX-FAOC018519/>. Accessed at Jan 24, 2024.
- Food and Drug Administration [FDA]. 2004. Freedom of information summary: Supplemental new animal drug application. Available from: <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/355>. Accessed at Jan 24, 2024.
- Food and Drug Administration [FDA]. 2017. Freedom of information summary: Original new animal drug application. NADA 141–450: Banamine transdermal, flunixin transdermal solution. Available from: <https://animaldrugsatfda.fda.gov/adafda/app/search/public/document/downloadFoi/1944>. Accessed at Mar 13, 2024.
- Food and Drug Administration [FDA]. 2020. Drug residues. Available from: <https://www.fda.gov/animal-veterinary/compliance-enforcement/drug-residues>. Accessed at Dec 9, 2023.

- Ferrer C, Lozano A, Agüera A, Jiménez Girón A, Fernández-Alba AR. 2011. Overcoming matrix effects using the dilution approach in multiresidue methods for fruits and vegetables. *J Chromatogr A* 1218:7634-7639.
- Fialkov AB, Steiner U, Lehotay SJ, Amirav A. 2007. Sensitivity and noise in GC–MS: Achieving low limits of detection for difficult analytes. *Int J Mass Spectrom* 260:31-48.
- Gallo P, Fabbrocino S, Dowling G, Salini M, Fiori M, Perretta G, Serpe L. 2010. Confirmatory analysis of non-steroidal anti-inflammatory drugs in bovine milk by high-performance liquid chromatography with fluorescence detection. *J Chromatogr A* 1217:2832-2839.
- Hajrulai-Musliu Z, Uzunov R, Jovanov S, Jankuloski D, Stojkovski V, Pendovski L, Sasanya JJ. 2021. A new LC–MS/MS method for multiple residues/contaminants in bovine meat. *BMC Chem* 15:62.
- Jang S, Seo H, Kim H, Kim H, Ahn J, Cho H, Hong S, Lee S, Na T. 2022. Development of a quantitative method for detection of multiclass veterinary drugs in feed using modified QuPPE extraction and LC–MS/MS. *Molecules* 27:4483.
- Jedziniak P, Szprengier-Juskiewicz T, Olejnik M, Jaroszewski JJ. 2016. Determination of flunixin and 5-hydroxyflunixin in bovine plasma with HPLC-UV-method development, validation and verification. *Bull Vet Inst Pulawy* 51:261-266.
- Kari F, Bucher J, Haseman J, Eustis S, Huff J. 1995. Long-term exposure to the anti-inflammatory agent phenylbutazone induces kidney tumors in rats and liver tumors in mice. *Japanese J Cancer Res* 86:252-263.
- Kissell LW, Baynes RE, Riviere JE, Smith GW. 2013. Occurrence of flunixin residues in bovine milk samples from the USA. *Food Addit Contam A* 30:1513-1516.
- Krueve A, Künnapas A, Herodes K, Leito I. 2008. Matrix effects in pesticide multi-residue analysis by liquid chromatography–mass spectrometry. *J Chromatogr A* 1187:58-66.
- Kwon H, Lehotay SJ, Geis-Asteggiane L. 2012. Variability of matrix effects in liquid and gas chromatography–mass spectrometry analysis of pesticide residues after QuEChERS sample preparation of different food crops. *J Chromatogr A* 1270:235-245.
- Levionnois OL, Fosse TK, Ranheim B. 2018. PK/PD modeling of flunixin meglumine in a kaolin-induced inflammation model in piglets. *J Vet Pharmacol Ther* 41:314-323.
- Liu ZY, Yang K, Chen FH, Long XM, Deng YB, Kuang GW, Sun ZL. 2015. Development of a rapid method for the confirmatory analysis of flunixin residue in animal tissues using liquid chromatography–tandem mass spectrometry. *Food Anal Methods* 8:352-362.
- Lugoboni B, Barbarossa A, Gazzotti T, Zironi E, Farabegoli F, Pagliuca G. 2014. A quick LC–MS-MS method for the determination of flunixin in bovine muscle. *J Anal Toxicol* 38:80-85.
- Malone EM, Dowling G, Elliott CT, Kennedy DG, Regan L. 2009. Development of a rapid, multi-class method for the confirmatory analysis of anti-inflammatory drugs in bovine milk using liquid chromatography tandem mass spectrometry. *J Chromatogr A* 1216:8132-8140.
- Ministry of Food and Drug Safety [MFDS]. 2023a. Standard for the maximum residue level of animal drugs in food [11-1471000-0001261-10]. Available from: <https://residue.foodsafetykorea.go.kr/vd/mrl>. Accessed at Jan 24, 2024.
- Ministry of Food and Drug Safety [MFDS]. 2023b. Food code. Available from: <https://various.foodsafetykorea.go.kr/fsd/#/>. Accessed at Jan 24, 2024.
- Ngoh MA, Wislocki PG, Thompson K, Katz T, Weingarten A, Terhune T, Hurshman B. 2003. Residue depletion study and withdrawal period for flunixin-*N*-methyl glucamine in bovine milk following intravenous administration. *J Agric Food Chem* 51:4701-4707.

- Shin HC, Choi YS. 2022. A study of animal drug residue detection methods improvement among the Korean food code (20162MFDS621). Available from: <https://rnd.mfds.go.kr/RDCAC08F01View>. Accessed at Jan 24, 2024.
- Sidhu PK, Gehring R, Mzyk DA, Marmulak T, Tell LA, Baynes RE, Vickroy TW, Riviere JE. 2017. Avoiding violative flunixin meglumine residues in cattle and swine. *J Am Vet Med Assoc* 250:182-189.
- Silvestro L, Tarcomnicu I, Rizea Savu S. 2013. Matrix effects in mass spectrometry combined with separation methods: Comparison HPLC, GC and discussion on methods to control these effects. In *Tandem mass spectrometry: Molecular characterization*. Coelho AV, de Matos Ferraz Franco C (ed). IntechOpen, London, UK.
- Smith GW, Davis JL, Tell LA, Webb AI, Riviere JE. 2008. Extralabel use of nonsteroidal anti-inflammatory drugs in cattle. *J Am Vet Med Assoc* 232:697-701.
- Wilkowska A, Biziuk M. 2011. Determination of pesticide residues in food matrices using the QuEChERS methodology. *Food Chem* 125:803-812.