

RESEARCH NOTE

First Report of Anthracnose Caused by *Colletotrichum nymphaeae* on Strawberry Fruits in Korea

Je Hyeok Yoo¹, Myung Soo Park², Doo Hee Yi¹, and Myeong Hyeon Nam^{3,*}

¹Strawberry Research Institute, Chungnam ARES, Nonsan 32914, Korea

²Department of Crops and Forestry, Korea National University of Agriculture and Fisheries, Jeonju 54874, Korea


³Industrial Entomology Research Institute, Chungnam ARES, Gongju 32418, Korea

*Corresponding author: namtel7@korea.kr

ABSTRACT

Anthracnose fruit rot (AFR) has been observed in greenhouses during the harvesting period in the Chungnam Province of South Korea. Fruits infected inside the greenhouse show black or brown spots, orange conidial masses and in some areas of the infected parts, white fungal growths are visible. The size of these spots gradually expands, leading to the necrosis of the fruits and flowers. Three isolates were obtained from infected fruits and identified as strains of the *Colletotrichum acutatum* species complex based on morphological characteristics. Multilocus sequence analysis of actin, chitin synthase, glyceraldehyde-3-phosphate dehydrogenase genes, and internal transcribed spacer rDNA regions revealed that the isolates belong to a monophyletic group with the type strain of *C. nymphaeae*. This is the first time *C. nymphaeae* has been confirmed in strawberry fruit in Korea.

Keywords: Anthracnose fruit rot, *Colletotrichum nymphaeae*, Strawberry



OPEN ACCESS

pISSN : 0253-651X
eISSN : 2383-5249

Kor. J. Mycol. 2024 June, 52(2): 109-114
<https://doi.org/10.4489/kjm.520203>

Received: May 14, 2024

Revised: May 29, 2024

Accepted: June 10, 2024

© 2024 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Strawberries (*Fragaria x ananassa*) are a high income crop in Korea. In 2022, they were ranked as the third-highest income crop among greenhouse vegetables, follow cucumbers and eggplants, they also ranked fourth in terms of cultivation area among fruit and vegetable crops [1]. In Chungnam Province, various strawberry varieties such as Seolhyang [2] and Kingsberry [3] are cultivated.

In November 2021, anthracnose symptoms were observed on strawberries (cv. Kingsberry) in Nonsan, Chungnam Province. In the early stages of anthracnose fruit rot (AFR), the symptoms include blackening of certain tissues of the strawberry (Fig. 1A), followed by the enlargement of lesions, hindering proper fruit development, and the formation of orange-colored spores within the lesions (Fig. 1B). Similar symptoms were observed on the petioles, with sunken and blackened tissues similar to those on the fruit, accompanied by the formation of an orange conidial mass (Fig. 1C).




Fig. 1. Anthracnose symptoms of *Colletotrichum nymphaeae* on strawberry. (A) The initial symptom of anthracnose fruit rot on the surface of strawberry fruit. (B) The exacerbated symptom on the surface of strawberry fruit. (C) The symptom of anthracnose fruit rot on the peduncle. (D) Symptoms of artificial inoculation test with an isolate.

In Korea, AFR caused by *Colletotrichum acutatum* (J. H. Simmonds) was first reported in 2008 [4]. However, the primary concern here is anthracnose crown rot (ACR), not AFR [5]. Initially reported as caused by *C. gloeosporioides*, ACR was reclassified as resulting from *C. fructicola* based on molecular phylogenetic analyses [6].

To identify the fungus causing the AFR, infected fruit tissues displaying symptoms were first surface-sterilized with 70% alcohol for 2 min and then rinsed three times with sterilized water. Subsequently, the surface sterilized tissues were air-dried on sterilized tissue paper, placed onto water agar, and incubated at 25°C. Following incubation, the edges of the fungal mycelia grown from the tissue were transferred to potato dextrose agar (PDA; Difco, Becton Dickinson) under aseptic conditions. Single spore isolation was performed as previously described [7], and pure cultures were stored at 4°C.

Morphological characteristics, such as the size and shape of conidia, were examined using a compound microscope (Olympus BX46, Tokyo, Japan). Based on the morphological features as reported previously [8], the isolates were identified as belonging to the *C. acutatum* species complex. All isolates initially formed white mycelia on PDA, which later turned light brown starting from the center of the front side (Fig. 2A and 2B). The conidia were hyaline, single-celled, and cylindrical, with pointed ends (Fig. 2C). The size range of the conidia was 11.0-15.0 × 4.5-5.0 μm (Table 1), with appressoria being ovate or globose and brown in color (Fig. 2D), with a size range of 7.0-11.3 × 5.3-7.5 μm.




Fig. 2. Cultural and morphological characteristics of *Colletotrichum nymphaeae*. (A and B) Front and reverse colony morphologies of CAN211201 grown on potato dextrose agar after 7 days. (C) Conidia. (D) Appressoria. Scale bar=10 μ m.

Table 1. Morphological characteristics of the strain used in this study compared with a previous report on *Colletotrichum nymphaeae*.

Characteristic	<i>C. nymphaeae</i> (CAN211201)	<i>C. nymphaeae</i> ^a
Colony morphology	White mycelia, which later turned light brown starting from the center of the front side	Pinkish-orange conidial masses, which turned gray
Conidia size (μ m)	11.0-15.0 \times 4.5-5.0	10.6-15.9 \times 4.0-5.8
Conidia shape	One-celled, and cylindrical in shape with pointed ends	Cylindrical, one-celled, aseptate, hyaline

^a Described by [16].

For phylogenetic classification, genomic DNA was extracted from fungal isolates grown on PDA using a modified CTAB extraction protocol [9]. The ITS, *ACT*, *CHS-1*, and *GAPDH* gene regions were amplified using a previously described method [10]. DNA sequencing was performed on an ABI PRISM 3730XL analyzer (Thermo Fisher Scientific, Waltham, MA, USA) at Macrogen (Seoul, Korea). Each sequence was assembled and proofread using MEGA v.7 [11]. The sequences generated in this study were deposited in GenBank (PP839268-PP839270 for ITS, PP908669-PP908671 for *ACT*, PP908672-PP908674 for *CHS-1*, PP908675-PP908677 for *GAPDH*). Multiple alignments were performed using MAFFT ver. 7 [12]. Maximum likelihood phylogenetic analyses were conducted using RAxML [13,14] implemented on the CIPRES web portal using the GTR+G model with 1,000 bootstrap replicates. Phylogenetic analysis revealed that three isolates formed a monophyletic group with *C. nymphaeae* CBS 515.78 (ex-type), CBS 526.77, CBS 129936, and CBS 126366 (bootstrap support=89%; Fig. 3).




Fig. 3. Maximum likelihood phylogenetic tree based on the concatenated datasets (ITS, *ACT*, *CHS-1*, and *GAPDH*) used to identify *Colletotrichum* strains isolated from strawberry in Korea. Bootstrap scores greater than 70 are presented at the nodes. The scale bar indicates the number of nucleotide substitutions per site and the letter ^T indicates ex-type strains. The strains originating from strawberry are indicated in bold.

The isolate, CAN211201, was utilized for artificial inoculation for pathogenicity testing and a suspension of 1×10^5 conidia/mL was sprayed onto 1 mL of each of the five fruits of the 'Kingsberry' cultivar. Five fruits treated with sterilized water were used as controls. The inoculated fruits were kept in a moist box (100% relative humidity) at 25°C for 5 days. During pathogenicity testing, the isolated strain exhibited typical symptoms of AFR on fruits, initially showing a black lesion. Eventually, due to high humidity, the lesion was covered with white fungi (Fig. 1D), while the controls remained symptom-free. The fungus was successfully re-isolated and identified as *C. nymphaeae* through morphological and phylogenetic analysis.

Morphological and phylogenetical analyses revealed that these three isolates were identified as *C. nymphaeae*, and pathogenicity tests indicated that the isolate showed typical symptoms of AFR.

Previously, *C. acutatum*, not *C. nymphaeae*, has been reported to cause AFR in strawberries in Korea. However, *C. nymphaeae* has been reported in strawberry plants in Egypt [15] and Argentina [16]. Additionally, anthracnose caused by *C. nymphaeae* in Korea has been reported in kiwiberries [17], Japanese plums [18], and persimmons [19]. Given these findings, it is necessary to investigate whether *C. nymphaeae* originated from other plants or countries. In this study, *C. nymphaeae* inducing AFR in strawberry fruits was confirmed based on morphological features and molecular phylogenetic analyses.

Therefore, continuous monitoring should be conducted to determine if *C. nymphaeae* causing AFR occurs in other strawberry cultivation regions.

CONFLICT OF INTERESTS

No conflict of interest was reported by the authors

ACKNOWLEDGEMENT

This study was funded by a grant under the Rural Development Administration (Project No. PJ01533202).

REFERENCES

1. Korea Statistical Information Service. Cultivated area of fruit and vegetable crops and Income of greenhouse crops [Internet]. Daejeon: Statistics Korea; 2024 [cited 2024 May 20]. Available from: <https://kosis.kr>.
2. Kim TI, Jang WS, Nam MH, Lee WK, Lee SS. Breeding of strawberry Seolhyang for forcing culture. *IHC* 2006;231.
3. Kim HS, Jang WS, Kim TI, Nam MH, Lee IH, Lee HC. Breeding of big size strawberry 'Kingsberry' for forcing culture. *Kor J Hort Sci Technol* 2016;34:167.
4. Nam MH, Kim TI, Gleason ML, Song JY, Kim HG. First report on anthracnose fruit rot caused by *Colletotrichum acutatum* on strawberry in Korea. *Plant Dis* 2008;92:1247.
5. Nam MH, Yoo JH, Yun TS, Kim HH, Kim HG. Anthracnose of strawberry: Etiological and ecological characteristics and management. *Res Plant Dis* 2023;29:205-19.
6. Nam MH, Park MS, Lee HD, Yu SH. Taxonomic re-evaluation of *Colletotrichum gloeosporioides* isolated from strawberry in Korea. *Plant Pathol J* 2013;29:317-22.
7. Prihastuti H, Cai L, Chen H, McKenzie EHC, Hyde KD. Characterization of *Colletotrichum* species associated with coffee berries in northern Thailand. *Fungal Divers* 2009;39:89-109.
8. Damm U, Cannon PF, Woudenberg JHC, Crous PW. The *Colletotrichum acutatum* species complex. *Stud Mycol* 2012;73:37-113.
9. Zhang L, Song L, Xu X, Zou X, Duan K, Gao Q. Characterization and fungicide sensitivity of *Colletotrichum* species causing strawberry anthracnose in eastern China. *Plant Dis* 2020;104:1960-8.
10. Rogers SO, Bendich AJ. Extraction of total cellular DNA from plants, algae and fungi. In: Gelvin SB, Schilperoort RA, editors. *Plant molecular biology manual*. Dordrecht: Springer Netherlands; 1994. p. 183-90.
11. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Bio Evol* 2011;28:2731-9.
12. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Mol Biol Evol* 2013;30:772-80.
13. Stamatakis A. RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 2006;22:2688-90.

14. Miller MA, Pfeiffer W, Schwartz T. Creating the CIPRES science gateway for inference of large phylogenetic trees. LA: SC10 workshop on gateway computing environments (GCE10); 2010. pp. 1-8.
15. Abo-Elwafa TM, Ragab SSM, Nehela Y, Essa TA. First report of strawberry anthracnose caused by *Colletotrichum nymphaeae* in Egypt. *New Dis Rep* 2023;48:e12205.
16. Fernandez LM, Maumary RL, Seimandi GM, Pernuzzi C, Derita MG, Favaro MA. First report of anthracnose caused by *Colletotrichum nymphaeae* on strawberry fruits in central Argentina. *Plant Dis* 2024;108:221.
17. Kim GH, Choi DH, Park SY, Koh YJ. First report of anthracnose caused by *Colletotrichum nymphaeae* on kiwiberry in Korea. *Plant Dis* 2018;102:1455.
18. Chang TH, Hassan O, Lee YS. First report of anthracnose of Japanese plum (*Prunus salicina*) caused by *Colletotrichum nymphaeae* in Korea. *Plant Dis* 2018;102:1461.
19. Hassan O, Lee DW, Chang T. First report of anthracnose of persimmon caused by *Colletotrichum nymphaeae* in Korea. *Plant Dis* 2019;103:1772.