



# Korean treatment recommendations for patients with axial spondyloarthritis

Mi Ryoung Seo<sup>1</sup>, Jina Yeo<sup>1</sup>, Jun Won Park<sup>2</sup>, Yeon-Ah Lee<sup>3</sup>, Ju Ho Lee<sup>4</sup>, Eun Ha Kang<sup>4</sup>, Seon Mi Ji<sup>5</sup>, Seong-Ryul Kwon<sup>6</sup>, Seong-Kyu Kim<sup>7</sup>, Tae-Jong Kim<sup>8</sup>, Tae-Hwan Kim<sup>9</sup>, Hye Won Kim<sup>10</sup>, Min-Chan Park<sup>11</sup>, Kichul Shin<sup>12</sup>, Sang-Hoon Lee<sup>13</sup>, Eun Young Lee<sup>2</sup>, Hoon Suk Cha<sup>14</sup>, Seung Cheol Shim<sup>15</sup>, Youngim Yoon<sup>2</sup>, Seung Ho Lee<sup>16</sup>, Jun Hong Lim<sup>17</sup>, Han Joo Baek<sup>1</sup>, and on behalf of the Korean Society of Spondyloarthritis Research

<sup>1</sup>Division of Rheumatology, Department of Internal Medicine, Gil Medical Center, Gachon University College of Medicine, Incheon; <sup>2</sup>Division of Rheumatology, Department of Internal Medicine, Seoul National University College of Medicine, Seoul; <sup>3</sup>Division of Rheumatology, Department of Internal Medicine, Kyung Hee University College of Medicine, Seoul; <sup>4</sup>Division of Rheumatology, Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam; <sup>5</sup>National Health Insurance Service, Wonju; <sup>6</sup>Division of Rheumatology, Department of Internal Medicine, Inha University College of Medicine, Incheon; <sup>7</sup>Division of Rheumatology, Department of Internal Medicine, Daegu Catholic University School of Medicine, Daegu; <sup>8</sup>Division of Rheumatology, Department of Internal Medicine, Chonnam National University Medical School and Hospital, Gwangju; <sup>9</sup>Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases, Seoul; <sup>10</sup>Division of General Internal Medicine, Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam; <sup>11</sup>Division of Rheumatology, Department of Internal Medicine, Yonsei University College of Medicine, Seoul; <sup>12</sup>Division of Rheumatology, Department of Internal Medicine, Seoul Metropolitan Government-Seoul National University Boramae Medical Center, Seoul; <sup>13</sup>Department of Rheumatology, Kyung Hee University Hospital at Gangdong, Kyung Hee University College of Medicine, Seoul; <sup>14</sup>Division of Rheumatology, Department of Internal Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul; <sup>15</sup>Division of Rheumatology, Daejeon Rheumatoid & Degenerative Arthritis Center, Chungnam National University Hospital, Daejeon; <sup>16</sup>Korea Ankylosing Spondylitis Society, Seoul; <sup>17</sup>Korea Ankylosing Spondylitis Corporation, Daejeon, Korea

Korean J Intern Med 2023;38:620-640  
<https://doi.org/10.3904/kjim.2023.194>

In the article “Korean treatment recommendations for patients with axial spondyloarthritis” by Seo MR et al. (Korean J Intern Med 2023;38:620-640), there is a mistake in the first author’s affiliation. The affiliation 18 should be added and corrected as follows:

## BEFORE

Mi Ryoung Seo<sup>1</sup>

<sup>1</sup>Division of Rheumatology, Department of Internal Medicine, Gil Medical Center, Gachon University College of Medicine, Incheon, Korea

## AFTER

Mi Ryoung Seo<sup>1,18</sup>

<sup>1</sup>Division of Rheumatology, Department of Internal Medicine, Gil Medical Center, Gachon University College of Medicine, Incheon, Korea

<sup>18</sup>Department of Medicine, Inha University Graduate School, Incheon, Korea

We apologize for any inconvenience that this may have caused.