



ERRATUM

비침습적 간섬유화 검사의 적용

인제대학교 의과대학 인제대학교 해운대백병원 내과

허내윤

Non-Invasive Evaluation of Liver Fibrosis

Nae-Yun Heo

*Department of Internal Medicine, Inje University Haeundae Paik Hospital, Inje University College of Medicine, Busan, Korea*Korean J Med 2023;98:84-87. <https://doi.org/10.3904/kjm.2023.98.2.84>

다음과 같이 본문 내 오류가 있어 수정합니다.

Table 1. Diagnostic performance of transient elastography in chronic liver diseases

Etiology	Liver fibrosis					
	≥ F2		≥ F3		F4	
	Cut-off (kPa)	AUROC	Cut-off (kPa)	AUROC	Cut-off (kPa)	AUROC
HCV [8]	7.1	0.83	9.5	0.90	12.5	0.95
HBV [9]	7.2	0.88	9.4	0.91	12.2	0.93
NAFLD [10]	8.2	0.77	9.7	0.80	13.6	0.89
Alcohol [11]	9.0	0.86	12.1	0.90	18.6	0.91

Table 1. Diagnostic performance of transient elastography in chronic liver diseases

Etiology	Liver fibrosis					
	≥ F2		≥ F3		F4	
	Cut-off (kPa)	AUROC	Cut-off (kPa)	AUROC	Cut-off (kPa)	AUROC
HCV [12]	7.1	0.83	9.5	0.90	12.5	0.95
HBV [13]	7.2	0.88	9.4	0.91	12.2	0.93
NAFLD [14]	8.2	0.77	9.7	0.80	13.6	0.89
Alcohol [15]	9.0	0.86	12.1	0.90	18.6	0.91

1. Wai CT, Greenson JK, Fontana RJ, et al. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology* 2003;38:518-526.
2. Lin ZH, Xin YN, Dong QJ, et al. Performance of the aspartate aminotransferase-to-platelet ratio index for the staging of hepatitis C-related fibrosis: an updated meta-analysis. *Hepatology* 2011;53:726-736.
3. Xiao G, Yang J, Yan L. Comparison of diagnostic accuracy of aspartate aminotransferase to platelet ratio index and fibrosis-4 index for detecting liver fibrosis in adult patients with chronic hepatitis B virus infection: a systemic review and meta-analysis. *Hepatology* 2015;61:292-302.
4. Sterling RK, Lissen E, Clumeck N, et al. Development of a simple noninvasive index to predict significant fibrosis in patients with HIV/HCV coinfection. *Hepatology* 2006;43:1317-1325.
5. Kim BK, Kim DY, Park JY, et al. Validation of FIB-4 and comparison with other simple noninvasive indices for predicting liver fibrosis and cirrhosis in hepatitis B virus-infected patients. *Liver Int* 2010;30:546-553.
6. Shah AG, Lydecker A, Murray K, et al. Comparison of noninvasive markers of fibrosis in patients with nonalcoholic fatty liver disease. *Clin Gastroenterol Hepatol* 2009;7:1104-1112.
7. Sandrin L, Fourquet B, Hasquenoph JM, et al. Transient elastography: a new noninvasive method for assessment of hepatic fibrosis. *Ultrasound Med Biol* 2003;29:1705-1713.
8. Yu JH, Lee HA, Kim SU. Noninvasive imaging biomarkers for liver fibrosis in nonalcoholic fatty liver disease: current and future. *Clin Mol Hepatol* 2023;29 Suppl:S136-S149.
9. Kim BK, Tamaki N, Imajo K, et al. Head-to-head comparison between MEFIB, MAST, and FAST for detecting stage 2 fibrosis or higher among patients with NAFLD. *J Hepatol* 2022;77:1482-1490.
10. Kang SH, Lee HW, Yoo JJ, et al. KASL clinical practice guidelines: management of nonalcoholic fatty liver disease. *Clin Mol Hepatol* 2021;27:363-401.
11. Korean Association for the Study of the Liver (KASL). KASL clinical practice guidelines for management of chronic hepatitis B. *Clin Mol Hepatol* 2022;28:276-331.
1. Wai CT, Greenson JK, Fontana RJ, et al. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology* 2003;38:518-526.
2. Lin ZH, Xin YN, Dong QJ, et al. Performance of the aspartate aminotransferase-to-platelet ratio index for the staging of hepatitis C-related fibrosis: an updated meta-analysis. *Hepatology* 2011;53:726-736.
3. Xiao G, Yang J, Yan L. Comparison of diagnostic accuracy of aspartate aminotransferase to platelet ratio index and fibrosis-4 index for detecting liver fibrosis in adult patients with chronic hepatitis B virus infection: a systemic review and meta-analysis. *Hepatology* 2015;61:292-302.
4. Sterling RK, Lissen E, Clumeck N, et al. Development of a simple noninvasive index to predict significant fibrosis in patients with HIV/HCV coinfection. *Hepatology* 2006;43:1317-1325.
5. Kim BK, Kim DY, Park JY, et al. Validation of FIB-4 and comparison with other simple noninvasive indices for predicting liver fibrosis and cirrhosis in hepatitis B virus-infected patients. *Liver Int* 2010;30:546-553.
6. Shah AG, Lydecker A, Murray K, et al. Comparison of noninvasive markers of fibrosis in patients with nonalcoholic fatty liver disease. *Clin Gastroenterol Hepatol* 2009;7:1104-1112.
7. Sandrin L, Fourquet B, Hasquenoph JM, et al. Transient elastography: a new noninvasive method for assessment of hepatic fibrosis. *Ultrasound Med Biol* 2003;29:1705-1713.
8. Yu JH, Lee HA, Kim SU. Noninvasive imaging biomarkers for liver fibrosis in nonalcoholic fatty liver disease: current and future. *Clin Mol Hepatol* 2023;29 Suppl:S136-S149.
9. Kim BK, Tamaki N, Imajo K, et al. Head-to-head comparison between MEFIB, MAST, and FAST for detecting stage 2 fibrosis or higher among patients with NAFLD. *J Hepatol* 2022;77:1482-1490.
10. Kang SH, Lee HW, Yoo JJ, et al. KASL clinical practice guidelines: management of nonalcoholic fatty liver disease. *Clin Mol Hepatol* 2021;27:363-401.
11. Korean Association for the Study of the Liver (KASL). KASL clinical practice guidelines for management of chronic hepatitis B. *Clin Mol Hepatol* 2022;28:276-331.
12. Castéra L, Vergniol J, Foucher J, et al. Prospective comparison of transient elastography, Fibrotest, APRI, and liver biopsy for the assessment of fibrosis in chronic hepatitis C. *Gastroenterology* 2005;128:343-350.
13. Li Y, Huang YS, Wang ZZ, et al. Systematic review with meta-analysis: the diagnostic accuracy of transient elastography for the staging of liver fibrosis in patients with chronic hepatitis B. *Aliment Pharmacol Ther* 2016;43:458-469.
14. Eddowes PJ, Sasso M, Allison M, et al. Accuracy of FibroScan controlled attenuation parameter and liver stiffness measurement in assessing steatosis and fibrosis in patients with nonalcoholic fatty liver disease. *Gastroenterology* 2019;156:1717-1730.
15. Nguyen-Khac E, Thiele M, Voican C, et al. Non-invasive diagnosis of liver fibrosis in patients with alcohol-related liver disease by transient elastography: an individual patient data meta-analysis. *Lancet Gastroenterol Hepatol* 2018;3:614-625.