

The solar chromosphere can be observed well through strong absorption lines. We infer the physical parameters of chromospheric plasmas from these lines using a multilayer spectral inversion. This is a new technique of spectral inversion. We assume that the atmosphere consists of a finite number of layers. In each layer the absorption profile is constant and the source function is allowed to vary with optical depth. Specifically, we consider a three-layer model of radiative transfer where the lowest layer is identified with the photosphere and the two upper layers are identified with the chromosphere. This three-layer model is fully specified by 13 parameters. Four parameters can be fixed to prescribed values, and one parameter can be determined from the analysis of a satellite photospheric line. The remaining eight parameters are determined from a constrained least-squares fitting. We applied the multilayer spectral inversion to the spectral data of the H α and the Ca II 854.21 nm lines taken in a quiet region by the Fast Imaging Solar Spectrograph (FISS) of the Goode Solar Telescope (GST). We find that our model successfully fits most of the observed profiles and produces regular maps of the model parameters. We conclude that our multilayer inversion is useful to infer chromospheric plasma parameters on the Sun.

[7 SS-05] Detection of Opposite Magnetic Polarity in a Light Bridge : Its Emergence and Cancellation in association with LB Fan-shaped Jets

Eun-Kyung Lim¹, Heesu Yang¹, Vasyl Yurchyshyn², Jongchul Chae³, Donguk Song⁴, Maria S. Madjarska⁵

¹KASI, ²BBSO, ³SNU, ⁴NAOJ, ⁵MPIS

Light bridges (LBs) are relatively bright structures that divide sunspot umbrae into two or more parts. Chromospheric LBs are known to be associated with various activities including fan-shaped jet-like ejections and brightenings. Although magnetic reconnection is frequently suggested to be responsible for such activities, not many studies presented firm evidence to support the scenario. We carry out magnetic field measurements and imaging spectroscopy of a LB where fan-shaped jet-like ejections occur with co-spatial brightenings at their footpoints. We study their fine photospheric structures and magnetic field changes using TiO images, Near-InfraRed Imaging Spectropolarimeter data, and H α data taken by the 1.6 m Goode Solar Telescope. As a result, we detect magnetic flux

emergence in the LB that is of opposite polarity to that of the sunspot. The new flux cancels with the pre-existing flux at a rate of 5.6×10^{18} Mx hr⁻¹. Both recurrent jet-like ejections and their footpoint brightenings are initiated at the vicinity of the magnetic cancellation, and show apparent horizontal extension along the LB at a projected speed of 4.3 km s⁻¹ to form the fan-shaped appearance. Based on these observations, we suggest that the fan-shaped ejections may have resulted due to slipping reconnection between the new flux emerging in the LB and the ambient sunspot field.

[7 SS-06] Surface exposure age of (25143) Itokawa estimated from the number of mottles on the boulder

Sunho Jin, Masateru Ishiguro
Seoul National University

Various processes, such as space weathering and granular convection, are occurring on asteroids' surfaces. Estimation of the surface exposure timescale is essential for understanding these processes. The Hayabusa mission target asteroid, (25143) Itokawa (Sq-type) is the only asteroid whose age is estimated from remote sensing observations as well as sample analyses in laboratories. There is, however, an unignorable discrepancy between the timescale derived from these different techniques. The ages estimated based on the solar flare track density and the weathered rim thickness of regolith samples range between 102 and 104 years [1][2]. On the contrary, the ages estimated from the crater size distributions and the spectra cover from 106 to 107 years [3][4].

It is important to notice that there is a common drawback of both age estimation methods. Since the evidence of regolith migration is found on the surface of Itokawa [5], the surficial particles would be rejuvenated by granular convection. At the same time, it is expected that the erasure of craters by regolith migration would affect the crater size distribution.

We propose a new technique to estimate surface exposure age, focusing on the bright mottles on the large boulders. Our technique is less prone to the granular convection. These mottles are expected to be formed by impacts of mm to cm-sized interplanetary particles. Together with the well-known flux model of interplanetary dust particles (e.g., Grün, 1985 [6]), we have investigated the timescale to form such mottles before they become dark materials again by the space weathering. In this work, we used three AMICA (Asteroid Multi-band Imaging Camera) v-band

images. These images were taken on 2005 November 12 during the close approach to the asteroid. As a result, we found the surface exposure timescales of these boulders are an order of 106 years. In this meeting, we will introduce our data analysis technique and evaluate the consistency among previous research for a better understanding of the evolution of this near-Earth asteroid.

References

- [1] Keller and Berger (2014), *Earth, Planets and Space*, Volume 66, article id.71
- [2] Noguchi et al. (2014), *Meteoritics & Planetary Science*, Volume 49, Issue 2, p. 188-214.
- [3] Bonal et al. (2015), *Meteoritics & Planetary Science*, Volume 50, Issue 9, p. 1562-1576.
- [4] Tatsumi et al. (2018), *Icarus*, Volume 300, p. 227-248.
- [5] Miyamoto et al. (2007), *Science*, Volume. 316, Issue 5827, p. 1011-1014.
- [6] Grün (1985), *Icarus*, Volume 62, Issue 2, p. 244-272.

[ㄱ SS-07] DEEP-South: 2nd phase of observations for small Solar System bodies

Myung-Jin Kim¹, Young-Jun Choi^{1,2}, Hongu Yang¹, Hee-Jae Lee^{1,3}, Dong-Heun Kim^{1,3}, Youngmin JeongAhn¹, Dong-Goo Roh¹, Hong-Kyu Moon¹, Chan-Kao Chang⁴, Josef Durech⁵, Miroslav Broz⁵, Josef Hanus⁵, Joseph Masiero⁶, Amy Mainzer⁷, James Bauer⁸

¹*Korea Astronomy and Space Science Institute, Korea*, ²*Univ of Science and Technology, Korea*, ³*Chungbuk National Univ, Korea*, ⁴*National Central Univ, Taiwan*, ⁵*Charles Univ, Czech Republic*, ⁶*NASA/JPL, USA*, ⁷*Univ of Arizona*, ⁸*Univ of Maryland*

DEEP-South (DEep Ecliptic Patrol of the Southern Sky) team will start the 2nd phase of KMTNet observation in Oct 2020. The DEEP-South observation mainly consists of three survey modes: (1) Activity survey (AS) that aims at finding active phenomena of small Solar System bodies. (2) Light curve survey (LS) targets to discover and characterize light variations of asteroids. And (3) Deep drilling survey (DS) focuses on the objects beyond the orbit of Jupiter (Centaurus and trans-Neptunian objects) as well as near Earth asteroids. For asteroid family (AF) studies and target of opportunity (TO) observations for urgent photometric follow-up, targeted mode will also be used.

DEEP-South team is awarded 7.0% of the telescope time at each site every year from Oct 2020 to Sep 2023 in the 2nd phase of KMTNet

operation which corresponds to about 75 full nights a year for the network. In this presentation, we will introduce our survey strategy and observation plan.

[ㄱ SS-08] ISudden brightness enhancements on main-belt objects

Hongu Yang¹, Hee-Jae Lee¹, Mingyeong Lee^{2,1}, Dong-Heun Kim^{3,1}, Masateru Ishiguro⁴, Hong-Kyu Moon¹, Youngmin JeongAhn¹, Young-Jun Choi^{1,2}
¹*Korea Astronomy and Space Science Institute*, ²*University of Science and Technology, Korea*, ³*Chungbuk National University*, ⁴*Seoul National University*

Dust ejection activities have been discovered from a few tens of asteroids since the first confirmation in 2006. Those objects are known as active asteroids. They provide good observational chances to study ongoing phenomena in the solar system such as sublimation of icy volatiles, mutual collisions among asteroids, rotational disintegrations, thermal fatigue, etc. Although dust ejection mechanisms of individual cases have been investigated through observations, the frequencies of the events and their connection to the overall evolutionary budget of the solar system have not yet been studied thoroughly, mainly because previous studies were based on serendipitous discoveries without any systematic surveys of these objects. In this work, we made wide-field monitoring observations of asteroids using Korea Microlensing Telescope Network (KMTNet) during the 2018/2019 winter season. Among 3,644 asteroids in the field-of-view, we detected nine candidates of brightness enhancements which we suspect as possible activities. It is still possible that some of those brightness increases have caused by long-term rotations. However, our observed frequency and brightness enhancements size-frequency distribution agrees with the expectations from impacts with decimeter sized objects, when the main belt objects size-frequency distribution observed down to decameter sized bodies are extrapolated to decimeter size.

[ㄱ SS-09] DEEP-South: Asteroid Light-Curve Survey Using KMTNet

Hee-Jae Lee^{1,2}, Hongu Yang¹, Dong-Heun Kim^{1,2}, Myung-Jin Kim¹, Hong-Kyu Moon¹, Chun-Hwey Kim², Young-Jun Choi^{1,3}
¹*Korea Astronomy and Space Science Institute*, ²*Chungbuk National University*, ³*University of Science and Technology*

Variations in the brightness of asteroids are